


DANGER—POISON—PRECAUTIONS 

CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist or dust may be fatal if swallowed.



CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

FOR CONTROL OF WIREWORMS IN NEWLY SET TOBACCO PLANTS: Apply carefully and uniformly as a broadcast application at the rate of 50 lbs. per acre, one to two weeks prior to setting. Disc into soil immediately.

DISPOSAL OF EMPTY CONTAINER

DO NOT re-use the container. Completely empty the contents and bury the unused or spilled chemical at least 18 inches deep in an isolated location away from water supplies. Burn the empty container completely. Exercise extreme caution and stay well away from the smoke. Do not use container in any connection with feed, food, or drinking water.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose.

Stop — Read the Label Before Use

HELENA

BRAND

4% Parathion Granules




CAN KILL YOU

POISON

See side panels for antidote and precautions

DANGER

Keep out of reach of children



ACTIVE INGREDIENT:

Parathion* 4%

INERT INGREDIENT 96%

TOTAL 100%

*O, O-diethyl O-p-nitrophenyl thiophosphate

NOT FOR HOME USE

USE ONLY WHEN WEARING THE FOLLOWING:

- (1) Wear water proof pants, coat, hat,
- (2) Wear goggles,
- (3) Wear mask or respirator approved for use with organophosphorus compounds,
- (4) Wear heavy duty natural rubber gloves.

Keep all unprotected persons and children away. Do not rub eyes or mouth with hands. If exposed to parathion and doctor cannot come, take victim to clinic immediately. Call a doctor (physician), clinic or hospital.

IMPORTANT! Before removing gloves, wash arms with soap and water before smoking.

AFTER WORK, take off all work clothes and wash thoroughly. Do not wear work clothes when leaving job. Do not wear work equipment with soap and water after work. See instructions included with reagent.

POISON

Parathion is a very dangerous poison. It can be absorbed through the skin and eyes. Clothing wet with this material should be removed and washed. Prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: blurred vision, pin-point pupils, tightening of eyes, drooling, twitching of mouth, nausea, vomiting, diarrhea, and convulsions.

FIRST AID

If swallowed and victim is awake (conscious), give water to drink. Do not give parathion and do not induce vomiting.

If breathing has stopped, start artificial respiration.

If swallowed and victim is awake (conscious), give water to drink. Do not give parathion and do not induce vomiting. If breathing has stopped, start artificial respiration.

In case of contact, immediately flush skin with plenty of water for 15 minutes and shower.

Active ingredient is O, O-diethyl O-p-nitrophenyl thiophosphate. It is a very dangerous poison. It can be absorbed through the skin and eyes. Clothing wet with this material should be removed and washed. Prompt medical treatment or they may die.

CAUTIONS 

SWALLOWED
If swallowed even in dust may be fatal if

SKIN CONTACT
If touched by hands or eyes or on clothing

BREATHED
If in fumes, spray mist or dust

POISON TO WILDLIFE
This is poisonous to wildlife. Birds and other animals may be killed. Keep away from water. Do not apply when weather is calm. Do not apply in protected areas. Do not apply near water. Do not apply near fish. Do not apply near birds. Do not apply near mammals. Do not apply near amphibians. Do not apply near reptiles. Do not apply near insects. Do not apply near plants. Do not apply near fungi. Do not apply near bacteria. Do not apply near viruses. Do not apply near parasites. Do not apply near predators. Do not apply near prey. Do not apply near humans.

AVOID SET TOBACCO
Do not broadcast apply to tobacco for two weeks prior to harvest.

CONTAINER
Always empty the container completely. Wash the container at least 18 inches from water supplies. Exercise extreme caution. Do not use container for drinking water. The container conforms to its label for the purpose of the purpose.

Stop Read the Label Before Use


HELENA

BRAND

4% Parathion Granules

POISON

See side panels for antidote and precautions



CAN KILL YOU

ACTIVE INGREDIENT:

Parathion* 4%

INERT INGREDIENT 96%

TOTAL 100%

*O, O-diethyl O-pnitrophenyl thiophosphate

NOT FOR HOME USE

DANGER

Keep out of reach of children

[Faint, illegible text, likely bleed-through from the reverse side of the page]

POISON SIGNS (Symptoms)

Inhalation, ingestion, or absorption of small amounts may cause irritation of the eyes, nose, throat, and skin. Ingestion of larger amounts may cause dizziness, weakness, nausea, vomiting, diarrhea, sweating, watering of eyes, and difficulty breathing. In severe cases, it may cause respiratory depression and coma.

FIRST AID TREATMENT

Call a doctor, physician, or hospital immediately. Explain that the victim has been exposed to parathion and describe the situation.

If breathing has stopped, start artificial respiration immediately, and maintain until doctor sees victim.

If swallowed and victim is awake, force him to vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes, while removing contaminated clothing and shoes. See doctor immediately.

NOTE TO PHYSICIAN

Antidote: Administer at a dose of 10 to 20 mg (1 mL) intravenously or intramuscularly at 10 to 15 minute intervals. Repeat at 15 to 30 minute intervals until signs of atropinization appear. **DO NOT USE MORPHINE OR TRANQUILIZERS**, which are strong central nervous system inhibitors affecting the central and peripheral nervous systems and tend to cause respiratory depression. At first sign