

CAUTION

Keep Out of Reach of Children


Harmful if swallowed.

- MAY CAUSE IRRITATION OF NOSE, THROAT, AND SKIN
- AVOID SKIN CONTACT.
- DO NOT BREATHE DUST.

Observe the Following Use Precautions

Do not contaminate any body of water; not apply in conditions favor drift from the crop area.

Apply only warm foliage when temperatures are above 60°F; do not apply sulphur when shade temperature is likely to exceed 90°F Fahrenheit.



ACTIVE INGREDIENTS:

Copper, expressed as metallic	4.0%
Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
Sulphur	75.0%
INERT INGREDIENTS	11.0%
TOTAL	100.0%

The copper is in the form of copper oxide.

NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so without acting with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and not include consequential, commercial damages such as loss of profit or value, etc.

DIRECTIONS FOR USE

Apply as a dust using ground equipment, aircraft or other means as otherwise specified, apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to control pest. Observe use limitations given for specific crops.

OBSERVE FOLLOWING USE PRECAUTIONS

Do not use to control insects if used with lime or other soil amendments.

Do not use on plants with sensitive roots and buds. If water is applied to plants with sensitive roots and buds, do not use on plants with sensitive roots and buds.

Do not use on plants with sensitive roots and buds.

Do not use on plants with sensitive roots and buds. Do not apply when crops are in bloom. Do not use on plants with sensitive roots and buds. Do not apply when crops are in bloom.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as timing, number and rate of applications needed will vary with local conditions.

10% SEVIN* — 4% COPPER — 75% SULPHUR DUST

*Trade Name of Cyanamid Corp. for the active ingredient carbaryl (1-naphthyl N-methylcarbamate)

U.S. Patent No. 2,907,476

RECOMMENDATIONS

PEANUTS: Potato Leafhopper, Thrips, Velvet Bean Caterpillar; Cercospora Leafspot — Apply at the rate of 10 pounds per acre when signs of insects first appear. If a heavy rain occurs within 24 hours of the first application repeat the application within three days. Repeat applications at 10 to 12 day intervals.

USE ONLY AS RECOMMENDED.

E.P.A. Reg. No. 5905-352

Completely empty container and bury unused chemicals at least 18 inches deep in isolated location away from water supplies. Burn empty container completely. Exercise extreme caution. Stay away from smoke. Bury ashes, residue.