

# WARNING

DO NOT BREATHE SPRAY MIST!  
 MAY BE ABSORBED THROUGH SKIN.  
 DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING.  
 IN CASE OF SKIN CONTACT, WASH WITH SOAP AND WATER.  
 AVOID CONTAMINATION OF FEED AND FOOD-STUFFS.  
 WASH THOROUGHLY AFTER USING AND BEFORE EATING OR SMOKING.

## OBSERVE FOLLOWING USE PRECAUTIONS

Do not apply during bloom when honey bees and other pollinators are active.  
 Avoid drift to adjoining food and forage crops.  
 Do not apply on feed or forage crops to be fed to dairy animals or to animals being finished for slaughter.

## POISONOUS TO FISH AND WILDLIFE.

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

### NOTICE OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so, except in writing with proper authorization from the company. Any damages resulting from a breach of this warranty shall be limited to the actual purchase price of the product. No consequential or special damages shall be recoverable.

**DESTROY EMPTY CONTAINER — Never RE-USE**



# TOXAPHENE 8-E

AN EMULSIFIABLE LIQUID  
 (Contains 8 lb. Toxaphene per Gallon)

**WARNING** Keep Out of Reach of Children

See side panel for additional required warning statements.

**MAY BE FATAL IF SWALLOWED**

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

ACTIVE INGREDIENTS:	BY WEIGHT
* Toxaphene .....	71.04%
Xylene .....	23.95%
INERT INGREDIENTS: .....	5.01%
	100.00%

\* Contains Chlorinated Camphene (Chlorine 67% to 92%)

MANUFACTURED BY

HELENA CHEMICAL COMPANY



# TOXAPHENE 8-E

AN EMULSIFIABLE LIQUID  
(Contains 8 lb. Toxaphene per Gallon)

**WARNING** Keep Out of Reach of Children  
See side panel for additional required warning statements.

**MAY BE FATAL IF SWALLOWED**

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

INGREDIENTS:	BY WEIGHT
.....	71.04%
.....	23.95%
.....	5.01%
.....	100.00%

Polychlorinated Camphene (Chlorine 67% to 69%)

## HELENA CHEMICAL COMPANY

### RECOMMENDATIONS


Unless otherwise indicated, dosage rates are given in terms of pints per acre of TOXAPHENE 8-E in sufficient water to thoroughly cover the plants.

**CELERY:** Armyworms, Cutworms — Use 1 to 2½ pts. per acre. Do not apply after the plant is half grown or after it starts to bunch. Apply to soil surface before planting if cutworms are known to be present and do not disturb the soil for 3 to 5 days.

**CORN:** Fall Armyworms, Corn Earworms — Use 1½ to 2 pts. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON:** Armyworms, Cotton Leafworms, Climbing Cutworms, Flea Beetles, Flea Shoppers, Grasshoppers, Plant Bugs, Thrips — Use 2 to 4 pts. per acre. DO NOT GRAZE DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER ON FIELDS TREATED LATE IN THE SEASON.

**TOMATOES:** Cabbage Loopers, Climbing Cutworms, Tomato Hornworms, Tomato Fruitworms, Armyworms, Tomato Pinworms, Flea Beetles — Use 1½ to 4 pts. per acre. Do not apply within 3 days of harvest.



FOR AIRCRAFT APPLICATION — Use with 1 to 3 gallons minimum of water per acre consistent with crop growth and good coverage.

USE ONLY AS RECOMMENDED!

EPA Reg. No. 5905-31

CONTENTS \_\_\_\_\_ GALLONS

## MEMPHIS, TENNESSEE