

CAUTION

Keep Out of Reach of Children
Harmful if Swallowed
Do not Breathe Dust
Avoid Contact With Skin and Clothing
Wash Thoroughly After Handling
Do not Contaminate Feed and Foodstuffs

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals, or onto adjoining food and fiber crops about to be harvested.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

Apply during early morning or evening when there is little or no wind.


Completely empty contents and bury unused chemical at least 18 inches deep in an isolated location away from water supplies. Burn empty container completely. Exercise extreme caution and stay well away from the smoke. Bury ashes and residues.

CAUTION: Do not use within 2 weeks of an oil spray treatment.



10% TOXAPHENE - 3.4%
COPPER - 70% SULPHUR DUST

An Agricultural Insecticide & Fungicide



CAUTION KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.

ACTIVE INGREDIENTS:

*Toxaphene	10.0%
Sulphur, as elemental	70.0%
Cupric oxide (CuO) ...	2.475%
(copper expressed as metallic) ...	1.98%
Cuprous oxide (Cu ₂ O) ..	1.323%
(copper expressed as metallic) ...	1.20%
Metallic Copper	0.22%
INERT INGREDIENTS:	16.6%
Total ...	100.0%

*Technical Chlorinated Camphene (67% to 69% chlorine)

EPA Reg No.

5905-359



GENERAL DIRECTIONS

Apply at first sign of infestation and infection, and repeat at 5 to 10 day intervals, depending upon severity of disease and insect condition, but observe use limits. For the control of diseases persisting after prescribed limits have expired, an effective fungicide should be used alone in such a manner which will not leave residues in excess of established tolerances.

CAUTION: Sulphur may burn foliage when temperatures are high. Do not apply sulphur when shade temperatures exceed or are likely to exceed 90° F.

PEANUTS: For control of Corn earworm and Velvetbean caterpillar - Apply 30 to 40 lb. per acre. For control of Leafhoppers, Fall armyworm, Southern armyworm and Thrips - Apply 20 to 25 lb. per acre. For control of Peanut leaf spot - Apply 10 to 24 lb. per acre. Begin when plants are 40 days old or when spots appear on lower leaves. Repeat at 10- to 14-day intervals and after rains. Consult your State Agricultural Experiment Station and County Agent for complete details on the peanut dusting program best suited to your local conditions.

Do not feed forage to dairy animals or to animals being finished for slaughter.

STATEMENT OF WARRANTY

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and not include consequential commercial damages such as loss of profit or value, etc.

NET WEIGHT

50 POUNDS