

NET CONTENTS:

USE AND APPLICATION DIRECTIONS

SMCP VAPONA INSECTICIDE 50% CONCENTRATE SOLUTION is a specially prepared concentrate for use in fogging devices.

TOBACCO WAREHOUSES: To control cigarette beetles and tobacco moths use a 2% fogging solution; dilute 1 quart of this product with 10 1/2 gallons of odorless oil or other oil solvents commonly used for diluting insecticidal poisons. Solvents must be non-flammable and known to be safe for use in tobacco warehouses. Apply the 2% fogging solution at a rate of 19 to 38 fl. ozs. per 10,000 cu. ft. (10 to 20 grams of VAPONA INSECTICIDE per 10,000 cu. ft.) Make application as often as necessary to maintain control. **TO BE APPLIED BY, OR UNDER THE DIRECTIONS OF TRAINED PERSONNEL** in a closed warehouse storing unfinished tobacco only.

Prepare the warehouse for treatment by closing all doors, windows and ventilators; lock or barricade all entrances, post warning signs and take whatever other precautions are necessary under the circumstances to prevent unprotected humans or domestic animals from entering the treated area. It is generally desirable to make the application in the late afternoon in order that the warehouse can be closed all night. All work in the warehouse must be suspended while warehouse remains closed. Workmen may enter the warehouse the morning after treatment.

The fogging apparatus and operators should remain outdoors and the fog directed into the warehouse. Operator must wear a full face mask of a type passed by the U. S. Bureau of Mines for DDVP (VAPONA Insecticide) protection. Operator must be clothed to protect hands, neck and all other skin surfaces.

DANGER


Human Cautions: Do not get on skin or in eyes. Do not breathe fumes. Avoid exposure to fumes. Wear clean natural rubber gloves, waterproof protective clothing and goggles when handling the concentrate to prepare dilutions. During application indoors wear a full face mask of the type passed by the U. S. Bureau of Mines for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational areas.

In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted solution. Mix the concentrate outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away. It is recommended that only the amount of diluted solution be prepared from the concentrate that will be used immediately. Do not contaminate feed, water or foodstuffs. Not for use or storage in or around the home.

Environmental Cautions: This product is toxic to fish, birds or other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DO NOT apply in any manner not specified on the label.



ACTIVE INGREDIENTS:

- 2,2-Dichlorovinyl dimethyl phosphate.....
- Related Compounds.....
- Aromatic Petroleum Derivative Solvent.....

INERT INGREDIENTS:.....

TOTAL



**DANGER
POISON**

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH REACH OF CHILDREN

See Antidote Statement and other Precautions

NET CONTENTS:



ACTIVE INGREDIENTS:	
2,2-Dichlorovinyl dimethyl phosphate.....	46.5%
Related Compounds.....	3.5%
Aromatic Petroleum Derivative Solvent.....	50.0%
INERT INGREDIENTS:.....	0.0%
TOTAL.....	100%

ACCEPTED
 OCT 7 1975
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.



**DANGER
 POISON**



POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES. KEEP OUT OF REACH OF CHILDREN

See Antidote Statement and other Precautions on side panel

ANTIDOTE

ATROPINE is the emergency antidote. Administer 1/100-grain atropine tablets for emergency cases of SUSPECTED POISONING.

INTERNAL: If the material has been in contact with the mouth, introduce a finger into the throat or induce vomiting. DO NOT INDUCE VOMITING BY ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been in contact with the skin, remove all contaminated clothing. If the material gets into the eyes, wash immediately with large amounts of water.

IF WARNING SYMPTOMS APPEAR: Administer atropine (see below). Keep patient prone and quiet. Transport the patient immediately to a hospital.

WARNING SYMPTOMS: Symptoms include non-reactive pin-point pupils, salivation, and blurred vision.

TREATMENT: Atropine is the specific antidote for nervous stimulation. If there are signs of respiratory depression, inject at 15 minute intervals, in doses of 2 mg. Morphine is contraindicated. Clear the airway if necessary. Observe patient continuously. Do not give any food or drink until the patient may eat without warning, cause prolo- nged vomiting, or cause prolo- nged inhibition. Allow no further exposure to the material until a blood test by blood test.

2-PAM is also antidotal and may be used if atropine is not available.

NOTICE: Seller's guaranty shall be void if the purchaser assumes any risk to persons or property.