

## CAUTION

### Keep Out of Reach of Children

**CAUTION:** Harmful if swallowed. Avoid inhalation of vapors or spray mist.

For foliage applications, the spray tank must be free of any sulfur residue before using OMNI Supreme Spray. Do not combine Sulfur sprays with OMNI Supreme Spray on foliage and do not follow a previous application of Sulfur or apply Sulfur following OMNI Supreme Spray until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulfur, using 1 quart to 25 gallons water. If sufficient time has not elapsed, burn will show in several days. Do not use OMNI Supreme Spray when temperatures are above 90° Fahrenheit. When temperatures drop below 32° F. (usually night-time lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° F. and are free from moisture due to melting frost or heavy dews. Do not use on foliage closer than 60 days after or 90 days before using Captan or Phaltan products. Do not use in combination with SEVIN\* on crops other than citrus. Application to trees weakened by disease, drought, driving winds, or high nitrogen applications may result in oil injury.

When this product is combined with Parathion or Cythion,\*\* it is toxic to fish. When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, limitations for the products involved. Keep out of any body of water.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.**

Do not reuse container for any substance that may be consumed by animals or humans.

OMNI Supreme Spray is an especially refined Phytonomic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage. When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, limitations for the products involved.

**AMOUNTS GIVEN ARE FOR 100 GALLONS OF DILUTED SPRAY.**

**ALFALFA (Seed Crop Only):** Use 1 quart per acre in 20 to 30 gallons of water by ground application or 10 to 15 gallons by air when used in combination and at the recommended label rates with either Cythion\*\* or Toxaphene for control of the pests listed on the product labels. Spider Mites: Use 1 quart per acre at the dilutions noted above with Kelthane\*\*\* at the recommended label rates.

**ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES:** Dormant or Delayed Dormant: Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale (Parlatoria Scale), Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Parlatoria, Soft Brown Scale, Certain Other Scale Insects, Eggs of Aphids, Red Spider Mites - 1 1/2 to 2 gals. Summer Use - Red Spider Mites - 3 to 4 qts.; Summer Stages of Brown Apricot Scale, San Jose Scale - 1 gal.

# OMNI

## Supreme Spray

### [INSECTICIDE-MITICIDE]

### LIQUID

### A Superior Oil Spray

## CAUTION

**KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS ON SIDE PANEL**

#### ACTIVE INGREDIENT

Petroleum Oil ..... 98%

#### INERT INGREDIENTS

Petroleum Oil Min. Unsaponated Residue-92%  
Petroleum Oil Classification - Unclassified.

ACCEPTED

10

5905-368

5 [ **FIGS:** Fig Scale - Dormant or Spray. Severe Infestation - Use at the recommended label rate.

6 [ **GRAPES:** Mealy bug - Dormant Parathion. Use 250 to 400 ga

7 [ **LEMONS, GRAPEFRUIT:** Red Spider Mites - 1 to 1 1/2 gals. Cythion\*\*. For best results, Scale are more easily controlled. Do not use within 8 weeks of

8 [ **OLIVES:** Parlatoria Scale. Spider Mites. OMNI Supreme Spray at recommended rate of Ethyl pre-bloom or in post-bloom

9 [ **NAVEL ORANGES:** Red Scale, Mites - 3 qts. OMNI Supreme. Do not use on Navel Orange

10 [ **VALENCIA ORANGES:** Red Scale, Spider Mites - 1 gal. applying Spray plus the recommended

11 [ **PEARS:** Pear Psylla - Delayed Blister Mite - Delayed Dormant

12 [ **ORNAMENTALS and SHADE:** Sycamores, Hawthorne, Flowering Maple, Dogwood, Oak, Privet, Brown Apricot Scale, Black Scale, etc. 100 to 1000 gals. water. Dormant Spray. NOTE: May cause discoloration not tolerant to oil sprays.

13 [ **WALNUTS:** Late Spring to Mid European Fruit Scale, San Jose Scale to 4 quarts. The addition of oil scale control: Trithion, Ethyl F dosages. Do not apply after

**MIXING DIRECTIONS:** With agitator. When combining with Wettable Powders, add these first when tank is 1/3 full, then add oil and while there is sufficient emulsification.

**WARRANTY AND LIMITATIONS:** This material is warranted that this material is reasonably fit for the purposes stated in the Conditions of Sale, subject to the Chemical Company makes no other express or implied warranty. The agent of Helena Chemical Company makes no specific reference to this warranty shall be limited to direct commercial damages such as

children

vapors or spray mist.

of any sulfur residue  
sulfur sprays with OMNI  
application of Sulfur or  
time has elapsed for  
of one tree previously  
sufficient time has not  
OMNI Supreme Spray when  
temperatures drop below 32° F.  
no crops until daytime  
F. and are free from  
on foliage closer than  
products. Do not use in  
Application to trees  
previous applications may

on,\*\* it is toxic to fish.  
icides as directed on this  
products involved. Keep

ANCE WITH CAUTIONS  
E STATE AND FEDERAL

PUT CONCENTRATE OR

consumed by animals or

hytonomic Oil Contact  
ormant stage.  
icides as directed on this  
e products involved.

D SPRAY.

to 30 gallons of water by  
d in combination and at  
Toxaphene for control of  
1 quart per acre at the  
mended label rates.

ES, PLUMS, PRUNES-Dor  
Scale, San Jose Scale,  
Scale, Italian Pear Scale,  
Soft Brown Scale, Certain  
1/2 to 2 gals. Summer  
Brown Apricot Scale, San

# OMNI

## Supreme Spray

### [INSECTICIDE-MITICIDE]

### LIQUID

### A Superior Oil Spray

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS ON SIDE PANEL

#### ACTIVE INGREDIENT

Petroleum Oil ..... 98%

#### INERT INGREDIENTS

Petroleum Oil Min. Unsulfonated Residue-92%  
Petroleum Oil Classification - Unclassified.

ACCEPTED

5905-368

- 3 [ **FIGS:** Fig Scale - Dormant or Delayed Dormant - Use 3 gals. OMNI Supreme Spray. Severe Infestation - Use 2 gals. OMNI Supreme Spray plus Ethyl Parathion at the recommended label rates.
- 6 [ **GRAPES:** Mealy bug - Dormant: 1 gal. plus the recommended rate of Ethyl Parathion. Use 250 to 400 gals. spray per acre, depending upon size of vines.
- 7 [ **LEMONS, GRAPEFRUIT:** Red Scale, Yellow Scale, Purple Scale, Black Scale, Red Spider Mites - 1 to 1 1/2 gals. OMNI Supreme Spray plus the recommended rate of Cythion \*\*. For best results, spray during winter and spring months when Red Scale are more easily controlled and trees generally less reactive to oil sprays. Do not use within 8 weeks of harvest on Lemons.
- 8 [ **OLIVES:** Parlatoria Scale, Spider Mites, Oleander Scale, Black Scale - Use 1 1/2 gals. OMNI Supreme Spray or combine 1 gal. OMNI Supreme Spray with the recommended rate of Ethyl Parathion per 100 gals. water. Apply in the pre-bloom or in post-bloom up to buckshot.
- 9 [ **NAVEL ORANGES:** Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites - 3 qts. OMNI Supreme Spray plus the recommended rate of Cythion \*\*. Do not use on Navel Oranges after Sept. 1.
- 10 [ **VALENCIA ORANGES:** Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites - 1 gal. applying not later than Sept. 1, or 3 qts. OMNI Supreme Spray plus the recommended rate of Cythion \*\*, applied not later than Sept. 15.
- 11 [ **PEARS:** Pear Psylla - Delayed Dormant: 2 gals. Summer Use: 1 to 1 1/2 gals. Blister Mite - Delayed Dormant and Fall Spray - 2 gals.
- 12 [ **ORNAMENTALS and SHADE TREES** (Conifers (see Note), Holly, Pyracantha, Sycamores, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch): Soft Brown Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites and Eggs - 1 gal. to 100 gals. water. Dormant Spray on Deciduous Varieties - 2 gals. per 100 gals. water. Dormant Spray on Other Varieties - 1 gal. per 100 gals. water. NOTE: May cause discoloration of blue spruce. Do not use on conifers or ferns not tolerant to oil sprays.
- 13 [ **WALNUTS:** Late Spring to Mid-Summer Only: Summer Stages of Frosted Scale, European Fruit Scale, San Jose and other Soft-shelled Scale, Red Spider Mites - 2 to 4 quarts. The addition of one of the following phosphate insecticides aids in scale control: Trithion, Ethyl Parathion, or Ethion. See label for recommended dosages. Do not apply after husk split.
- MIXING DIRECTIONS:** With agitator running start filling spray tank with water. When combining with Wettable Powder insecticide or fungicide products, add these first when tank is 1/3 filled. Add OMNI Supreme Spray when tank is about 1/2 full and while there is sufficient agitation of the spray to insure complete emulsification.
- WARRANTY AND LIMITATION OF DAMAGES.** Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.