

CAUTION

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90° or above). Do not make sulfur applications at such times. Do not use sulfur on Eastern American Type Grapes, Cucumbers, Apricots or other sulfur susceptible plants. Do not make applications within two weeks before or after an oil spray. On citrus do not apply within 60 days prior to or after an oil spray.

Workers are not to enter treated areas for any activity requiring substantial contact with treated foliage for 24 hours after application.

Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Burn or bury empty containers, never reuse. Stay out of smoke or fumes when burning containers.

DIRECTIONS

ALFALFA (Seed): Spider Mites - 20 to 25 lbs. per acre. Do not apply during blooming period in desert areas.

ALMONDS, CURRANTS, FIGS, HOPS, PLUMS, PRUNES, WALNUTS: Almond Mite, European Red Mite, Common Red Spiders and Pacific Mite - 50 lbs. per acre. Dust when Spiders begin to multiply in large numbers. Since the effectiveness of sulfur for these Red Spiders varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.

ASPARAGUS: Rust - 15 to 20 lbs. per acre. Dust after cutting stops. Irrigate, cultivate, then apply the sulfur. Repeat if disease is severe.


BEANS, PEAS: Common Red Spiders and Pacific Mite - 15 to 20 lbs. per acre. Dust when spiders begin to multiply in large numbers. Since the effectiveness of sulfur for these Red Spiders varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.



90% SULFUR DUST

CAUTION KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTIONS ON SIDE PANEL.

ACTIVE INGREDIENT :	BY WEIGHT
Sulfur as elemental	90.0%
INERT INGREDIENTS	10.0%
TOTAL	100.0%



MANUFACTURED BY
HELENA CHEMICAL COMPANY

BEANS, CURRANTS, DAHILAS, PEAS, PEACHES, ROSES, STRAWBERRIES: Powdery Mildew - 15 to 20 lbs. per acre on beans, dahilas, peas, and strawberries. 30 to 40 lbs. per acre on Roses. 50 lbs. per acre on currants and peaches. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

CHERRIES: Leaf Spot and or Brown Rot - 40 to 45 lbs. per acre. Make first application at Pink Bud Stage, repeat at petal fall and shuck fall stages. Repeat as necessary through growing season.

CITRUS: Citrus Red Spider, Two-Spotted Mite, Thrips - 80 to 100 lbs. per acre. Dust when infestation threatens.

COTTON: Atlantic Mite, Pacific Mite, Two-Spotted Mite - 25 to 35 lbs. per acre.

GRAPES: Powdery Mildew - 7 to 15 lbs. per acre. Dust first when shoots are 6 inches long. Follow a dusting program of application each 7 to 10 days.

MELONS: Powdery Mildew - 10 to 20 lbs. per acre. May be controlled by dusting, but the sulfur often injures the vines, particularly if the temperature reaches 95°F. Treat when disease first appears and re-treat as necessary. Do not use on sulfur sensitive varieties.

PEACHES, PLUMS, PRUNES: Brown Rot - 50 lbs. per acre. Put on a preventive dusting when the fruit starts to ripen. This often reduces the severity of the disease. Rust - Apply 50 lbs. per acre during summer when disease appears. Repeat after 2 weeks if necessary.

SUGAR BEETS: Spider Mites - 30 to 40 lbs. per acre.

TOMATOES: Russet Mite, Two-Spotted Spider Mite - 20 to 50 lbs. per acre. Because of sulfur, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

NOTICE OF WARRANTY

WARRANTY AND LIMITATION OF DAMAGES. Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other expressed or implied warranty, including any other expressed or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or value, etc.

NET WEIGHT: 50 lbs.

EPA Reg. No. 5905-422