

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Swallowing, inhaling, or getting on skin or clothes may irritate. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water. If you get it on your skin, wash with soap and water. If you get it in your eyes, wash with clean water for 15 minutes. If you get it on your clothes, wash with soap and water.

ENVIRONMENTAL HAZARDS

Do not apply to water bodies. Do not apply to crops that are used for animal feed. Do not apply to crops that are used for human consumption.

DIRECTIONS FOR USE

Use only as directed. Do not use more than the amount specified.

STORAGE AND DISPOSAL

Store in original container. Do not use if container is damaged. Do not use if contents are empty. Do not use if contents are solid. Do not use if contents are liquid. Do not use if contents are gas. Do not use if contents are other.

PRODUCT INFORMATION

This product is a liquid sulfur. It is used to control plant diseases and insects. It is safe for use on most crops. It is safe for use on most animals. It is safe for use on most humans.

**DO NOT USE WITHIN FOUR (4) WEEKS OF AN OIL SPRAYING
SULFUR IS HIGHLY CORROSIVE AND EQUIPMENT SHOULD BE
CLEANED THOROUGHLY BEFORE OIL SPRAYING.**



LIQUID SULFUR SIX

ACTIVE INGREDIENT

Sulfur 52.0%

INERT INGREDIENTS

48.0%

TOTAL

100.0%

This product contains 6 lbs. of sulfur per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SEE LEFT SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.**

ACCEPTED

APR 22 1981

NOT FOR HOME USE

DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE

**DIRECTIONS
BEFORE USING, SEE**

Label for full directions. Do not use if container is damaged. Do not use if contents are empty. Do not use if contents are solid. Do not use if contents are liquid. Do not use if contents are gas. Do not use if contents are other.

RECOMMEN

**WHEN GROWING CROPS
CONSULT THE PROCESSOR BE**

For specific directions, consult the processor. Do not use more than the amount specified. Do not use if container is damaged. Do not use if contents are empty. Do not use if contents are solid. Do not use if contents are liquid. Do not use if contents are gas. Do not use if contents are other.

APPLE PEAR

Scab

Use 1.5 to 2.0 gallons per acre of water.

Powdery Mildew

Use 1.5 to 2.0 gallons per acre of water.

ALMOND CHERRY PLUM AND PRUNE

Scab

Use 1.5 to 2.0 gallons per acre of water.

AVOCADOS

Use 1.5 to 2.0 gallons per acre of water.

BEANS PEAS, COLE CROPS, PEPPERS

Use 1.5 to 2.0 gallons per acre of water.

CITRUS (Oranges, Lemons, Grapefruit, Tangerines)

Use 1.5 to 2.0 gallons per acre of water.

THIRPS (any species) Treat with this product

Use 1.5 to 2.0 gallons per acre of water.



HELENA CHEMICAL COMPANY
Memphis, TN 38118

LENA

SULFUR SIX

WEIGHT: 52.0%
 Purity: 48.0%
 100.0%

Contains 6 lbs. of Sulfur per gallon

**REACH OF CHILDREN
 CAUTION
 LABEL PANEL FOR ADDITIONAL
 HAZARDARY STATEMENTS.**

FOR HOME USE
 CONTAINER FOR ANY PURPOSE

Liters) 1.9 L (0.509 US GAL)

DIRECTIONS FOR MIXING
 BEFORE USING STIR UNTIL SMOOTH
 DO NOT ALLOW MIXTURE TO STAND

RECOMMENDATIONS
 WHEN GROWING CROPS FOR PROCESSING
 CONSULT THE PROCESSOR BEFORE APPLYING SULFUR

- APPLE PEAR
- SOY
- FRUITING MALLOW
- ALMOND CHERRY PLUM AND PEACH
- AVOCADO
- BEANS PEAS COLL CROPS PEPPERS
- LETTUCE
- SPINACH
- TURNIP
- WATERMELON

LEACH ALLIUMS (Garlic, Shallots, Onions, etc.) Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

FIELD AND VINE CROPS Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

COTTON Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

GRAPES Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

FRUITING MALLOW Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

POTATOES Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

SUGAR BEETS Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

STRAWBERRIES Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

TURNIP Apply 1.0 to 1.5 lbs. of Sulfur Six per 100 gallons of water. Repeat at 7 to 10 day intervals. Do not apply to plants in flower. Do not apply to plants in fruit.

WARRANTY AND LIMITATION OF DAMAGES

