



US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 5905-437	DATE OF ISSUANCE MAY 14 1986
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT Helena Liquid Sulfur Six	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
<div style="display: flex; justify-content: space-between;"> <div style="width: 60%;"> <p>Helena Chemical Company 5100 Poplar Suite 3200 Memphis, TN 38137</p> </div> <div style="width: 35%; text-align: right;"> <p>686: 169862 17 1</p> </div> </div>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>Based on your response to the Registration Standard Guidance Package, EPA has reregistered the product listed above. This registration is granted under the authority of section 3(g) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended.</p> <p>The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.</p> <ol style="list-style-type: none"> Change the human hazard statement as follows: "Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse." Add the following physical hazard statement: "Keep away from heat, sparks, or flame. Do not smoke while applying this product." 		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL 		DATE 5/15/86

- 3. Change the environmental hazard statement to:
 - Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.
- 4. Add "Do not contaminate water, food, or feed by storage or disposal" directly under the Storage & Disposal heading.
- 5. Add "If burned, stay out of smoke" to the container disposal statement.
- 6. Change the Pesticide Disposal statement to read, "Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility."
- 7. The Storage & Disposal heading must be in 12-point type.
- 8. Add the appropriate language from PR Notice 83-2, Farmworker Safety.


 Henry M. Jacoby
 Product Manager (21)
 Fungicide-Herbicide Branch
 Registration Division (TS-767C)

00000000

ACCEPTED
with COMMENTS
In EPA Letter Dated:
REREGISTERED
MAY 14 1986



Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5905-437

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION
 Irritation to eyes. Avoid breathing spray mist. May cause irritation of nose and throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT
 If Swallowed: Flush mouth with glasses of water and induce vomiting if not already done.
 If on Skin: Wash thoroughly with soap and water.
 If in Eyes: Wash eyes with plenty of clear water for at least 15 minutes.

ENVIRONMENTAL HAZARDS
 Use of this product in a manner not in accordance with the information on this label may injure domestic animals, wildlife, aquatic life or produce other undesirable results. Do not apply directly to lakes, streams or ponds.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Workers are not to enter treated areas for any activity requiring substantial contact with treated foliage for 24 hours after application.

STORAGE AND DISPOSAL
 Product may be stored and if frozen may generally be restored after thawing and mixing. Repeated freezing-thawing cycles may impair utility.

CONTAINER DISPOSAL — Triple rinse or equivalent and offer for recycling or reconditioning or dispose of in a sanitary landfill or by incineration if permitted by state and local authorities.

PESTICIDE DISPOSAL — Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State, or local procedures under the Resource Conservation and Recovery Act.

PRODUCT INFORMATION
 Helena Liquid Sulfur Six is a flowable sulfur formulation that can be mixed with liquid fertilizers or water and applied by ground or air application. Consult your state or local extension specialist for additional treatment information.

Sulfur may cause injury if used improperly or under unfavorable weather conditions. During periods of high temperature sulfur may burn foliage. Do not make Sulfur applications at such times.

NOTE: Sulfur will cause severe and leaf injury to sulfur sensitive crops. Do not apply or allow to drift to apricots, orange and comice pears, cranberries, cucurbits (cucumber, cantaloupe, melons, squash), figs, sunchokes, tung trees, walnuts or other sensitive plants.

DO NOT USE WITHIN FOUR (4) WEEKS OF AN OIL SPRAYING
SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.

LIQUID SULFUR SIX


ACTIVE INGREDIENT:	
Sulfur	52.0%
INERT INGREDIENTS	48.0%
TOTAL	100.0%
This product contains 6 lbs. of Sulfur per gallon	

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NOT FOR HOME USE
DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE

NET CONTENTS 5 Gals (18.92 Liters) EPA REG NO 5905-4
SN-0180/0883-M EPA EST NO 37429-GA-





DIRECTIONS FOR MIXING BEFORE USING. STIR UNTIL SMOOTH

Pour recommended amount in partially filled spray tank. Keep agitator running during filling and spraying operation. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause Liquid Sulfur Six to settle and may necessitate manual stirring to redispense.

RECOMMENDATIONS:

WHEN GROWING CROPS FOR PROCESSING.

CONSULT THE PROCESSOR BEFORE APPLYING SULFUR.

Unless otherwise specified for specific crops, dosage rates are given as pints of Helena Liquid Sulfur Six per 100 gallons water to use in a thorough coverage spray. The total amount of Helena Liquid Sulfur Six per acre should be 2 to 4 gallons depending on the size of the trees. Because of variations in the types of spray equipment use, a range of low and high rates is listed. For high volume sprayers (output 600 to 1000 gallons spray per acre) use the low rate. For low volume sprayers (output 20 to 300 gallons spray per acre) use the high rate.

APPLE, PEAR

Scab
Pre-bloom through calyx sprays 1 to 3 pints
Cover Sprays 1/2 to 2 pints

Powdery Mildew

Pre-bloom through calyx sprays 3 pints
Cover Sprays 2 pints

ALMOND, CHERRY, PLUM AND PRUNE

Brown Rot leafspot
Pink and Bloom Sprays 1 to 3 pints
Petal fall, shuck & cover sprays 2/3 to 1 1/2 pints

AVOCADOS

Brown Mite Use 3.5 pints. Apply in a minimum of 8 gallons of water by air.

BEANS, PEAS, COLE CROPS, PEPPERS

Powdery Mildew, Red Spider Spp. Use 3-4 pints in 8 gallons of water by air. Apply at first sign of infection and repeat at 3 week intervals.

CITRUS: Oranges, Lime, Grapefruit, Lemons, Tangelos and Tangerines. To control Rust mites, Clover mite - Use 1/2 to 4 pints. Apply spray November thru May.

THRIPS Early spring treatment for the reduction of Thrips population. Apply 1 to 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

PEACH, NECTARINE To control Powdery Mildew, Brown Rot, Scabs - Use 1 to 3 pints per 100 gallons of water applied at pink and bloom. Use 2/3 to 1 1/2 pints per 100 gallons of water at petal fall, shuck split, shuck fall and at 10 to 14 day intervals as long as necessary. For Brown rot of fruit, apply when fruit starts to ripen. Application to mature nectarines may cause discoloration.

FIELD AND VINE CROPS

Unless otherwise specified for specific crops, dosage rates are given as pints of Helena Liquid Sulfur Six per acre. This product should be applied by ground equipment or aircraft in sufficient water to provide thorough coverage.

COTTON, Atlantic Mite. 2 pints for early season control and up to 1 gallon for mid to late season control. Do not apply when temperature exceeds 90 F.

GRAPES Powdery Mildew Control - Use 1-2 pints per 100 gallons of water, maximum of 8 pints LIQUID SULFUR SIX in spring and summer. Applications in California use 2-3 quarts per acre in enough water for thorough coverage or per 100 gallons of water in spring and summer applications. Make the first application when shoots are 6-8 inches long, the second when shoots are 12-15 inches long and the third at about bloom stage. Timing rates and number of applications vary with local conditions. Consult your state Agricultural Extension Service for specific recommendations.

PEANUTS For Rust - Use 2 to 4 pints per acre of LIQUID SULFUR SIX in enough water for good coverage. Apply when disease threatens. Repeat at 7 to 10 day intervals as necessary.

POTATOES Powdery Mildew. Use 3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

SUGAR BEETS: Powdery Mildew Control - Apply 2-4 quarts, depending on disease pressure, for 2-4 applications at 2 week intervals when disease symptoms first appear. For aerial application, use sufficient water to obtain adequate coverage.

STRAWBERRIES, Powdery Mildew. Use 2 pints. Apply at first sign of infection and repeat at 3 week intervals. For Strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canners' recommendations.

TOMATOES Tomato russet mite. Use 1 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

WARRANTY AND LIMITATION OF DAMAGES

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein. Helena Chemical Company makes no other expressed or implied warranty, including expressed or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Helena Chemical Company is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.