



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 5905-494	2. EPA Product Manager	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Setre Fluometuron 80WP Herbicide	PM#	
5. Name and Address of Applicant (Include ZIP Code) Helena Chemical Company 225 Schilling Boulevard, Suite 300 Collierville, Tennessee 38017 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  MAR 11 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Other Revisions  
Notification of Other Revisions per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information  
 Label  Container

4. Size(s) Retail Container

5. Location of Label Directions  
 On Label

6. Manner in Which Label is Affixed to Product  
Self Adhesive  Lithograph Paper glued Stenciled  Other \_\_\_\_\_

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Mandy K. Styles	Title Product Registration Supervisor	Telephone No. (Include Area Code) (901) 752-4470
-------------------------	--	---

**Certification**  
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Product Registration Supervisor	6. Date Application Received <b>(Stamped)</b>
4. Typed Name Mandy K. Styles	5. Date 3/1/2005	

MAR 11 2005

# FLUOMETURON 80WP HERBICIDE

For Preemergence and Postemergence Weed Control in Cotton

## ACTIVE INGREDIENT:

Fluometuron (1,1-dimethyl-3-(2,2,2-trifluoro-M-tolyl) urea)..... 80.0%

INERT INGREDIENTS\*..... 20.0%

TOTAL..... 100.0%

### KEEP OUT OF REACH OF CHILDREN

### WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you.)

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### WARNING - AVISO

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing spray mist.

### FIRST AID

**IF IN EYES:**

- Hold eyelid open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor immediately for treatment advice.

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin with plenty of water for 15-20 minutes.
- Call a poison control center or doctor immediately for treatment advice.

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- If instructed to induce vomiting, first have person drink promptly a large quantity of milk, egg whites, or gelatin solution; if these are not available, drink large quantities of water. Then, induce vomiting. Avoid alcohol.
- Do not give anything to an unconscious or convulsing person.

**IF INHALED:**

- Move victim to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth- to-mouth if possible.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency, call ChemTrec at 1-800-424-9300.

**Note to Physician:** There is no specific antidote for fluometuron. Induce emesis and treat symptomatically.

### SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 5905-494  
EPA EST. NO.

NET CONTENTS:

MANUFACTURED BY  
HELENA CHEMICAL COMPANY  
225 SCHILLING BOULEVARD, SUITE 300  
COLLIERVILLE TENNESSEE 38017

### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes Plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering controls statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal

protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: if product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a dry place.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**

**FLUOMETURON 80WP** is effective for both annual grass and annual broadleaf weed control in cotton.

When applied according to directions, **FLUOMETURON 80WP** will selectively control annual weeds in cotton such as:

Annual Grass Weeds

- |                   |              |
|-------------------|--------------|
| Barnyardgrass     | Fall panicum |
| <i>Brachiaria</i> | Foxtail      |
| Crabgrass         | Goosegrass   |
| Crowfootgrass     | Ryegrass     |

Annual Broadleaf Weeds

- |                       |                       |
|-----------------------|-----------------------|
| Buttonweed            | Pigweed               |
| Cocklebur             | Prickly sida          |
| Florida pusley        | (ironweed or teaweed) |
| Goathead              | Purslane              |
| Groundcherry (Wright) | Ragweed               |
| Jimsonweed            | Sesbania              |
| Lambsquarters         | Sicklepod             |
| Momingglory           | Smartweed             |
|                       | Tumbleweed            |

At rates up to 2.5 lbs. per acre, **FLUOMETURON 80WP** is not effective for the control of established perennial weeds, such as johnsongrass from rhizomes.

**MIXING INSTRUCTIONS**

**FLUOMETURON 80WP** is highly concentrated and should be thoroughly mixed as a paste or thick slurry in water in a separate

container before adding to the spray tank. The spray tank should contain 1/3-1/2 the total amount of water or nitrogen solution to be used before adding the slurry.

To determine compatibility, add one tablespoon product to one pint of nitrogen solution (equals 2.5 lbs. in 25 gals.; if using other application rates, adjust compatibility test amounts proportionally). Stir or shake thoroughly. Let stand 5 minutes. If it remains mixed or can be remixed readily, the mixture is compatible and can be used. If product settles out when mixed with nitrogen solution, a suspensibility agent, such as Compex® at 1-2 pts. per 100 gals. fertilizer may improve compatibility. Add 1/8-1/4 teaspoon of Compex to the compatibility test to determine its effectiveness.

Use bypass or mechanical agitation in the spray tank. Avoid leaving spray mixture without constant agitation. If bypass agitation is used, it should terminate at the bottom of the tank to minimize foaming. Wash sprayer thoroughly with clean water immediately after use.

**EQUIPMENT**

Thoroughly clean sprayer prior to use. Do not use a sprayer contaminated with 2,4-D or other materials as crop damage or sprayer clogging may result.

For uniform distribution, use fan-type nozzles. Use 50-mesh or coarser screens in nozzles and strainers. Use a suitable pump with capacity to deliver (1) the necessary volume to the nozzles at 35-40 psi, and (2) an additional volume to maintain the mixture in suspension in the spray tank. Apply 20-40 gals. of spray mixture per acre.

**BAND TREATMENT**

For band treatment, calculate the amount of **FLUOMETURON 80WP** needed by the formula:

band width		broadcast		amount
<u>in inches</u>	X	rate	=	needed
row width		per acre		per acre
in inches				
of field				

**COTTON**

**FLUOMETURON 80WP** may be applied as a preemergence treatment to the surface of the soil, or lightly incorporated immediately after planting, or as a directed, semi-directed, or over-the-top postemergence application. For preemergence treatment only, product may be tank-mixed with most liquid nitrogen fertilizers. Product may be combined with other herbicides in cotton as directed in the section, "Combinations".

**FLUOMETURON 80WP** is most effective when used as follows:

1st Application: Preemergence broadcast. See Preemergence Treatment Below.

2nd Application: Postemergence, directed, semi-directed, or over-the-top when cotton is 3-6 inches high. See Postemergence Treatments below.

3rd Application: If needed, postemergence, directed, or semi-directed, at lay-by (last normal cultivation), using no more than 1.25 lbs. per acre.

Preemergence Treatment

Apply at a rate of 1.25-2.5 lbs. per acre. On light silt and sandy soils, low in organic matter, use the lower rate.

Product may be applied to the surface of the soil, or lightly incorporated into the soil, after planting cotton.

Preplant Incorporated Use (Arizona, California, and New Mexico)

May be applied preplant as a broadcast spray at 2 lbs. per acre after the beds for preplant irrigation have been formed. Apply before the beds are dragged off (using a bed shaper set to remove 1-2 inches from beds) in preparation for planting. Furrow irrigation should precede the application and planting should follow as soon as possible. Plant into untreated soil after drag-off and return treated soil to the beds when furrows are reformed. Do not allow furrow irrigation water to pond over the beds.

#### Postemergence Treatments

May be applied postemergence to cotton, but preemergence to weeds, or postemergence to both cotton and weeds. Apply at rates of 1.25-2.5 lbs. per acre. Apply as a directed, semi-directed, or over-the-top spray. May be applied to cotton plants at any stage of growth from 3 inches high to lay-by. Use the higher rate when applying postemergence to weeds; weeds should be 2 inches high or less. Use of a surfactant that is approved by FDA for use on food and feed crops and is compatible with **FLUOMETURON 80WP** when applied to cotton may enhance the postemergence herbicidal activity of product. The surfactant rate should not exceed 0.5% of the carrier volume (1 pt. per 25 gals. of water).

Precautions: Rainfall that is sufficient to germinate seeds or for growing crops is adequate to activate the herbicidal activity; where dry weather conditions prevail, the herbicidal activity may be delayed or reduced.

Do not make more than 3 applications of **FLUOMETURON 80WP** to the same crops or field in any one year. Do not plant crops other than cotton within 6 months of the last application.

The use of **FLUOMETURON 80WP** as a preemergence or early postemergence application, following the use of a systemic insecticide at planting, may result in injury to cotton.

West Texas Only: Do not use on sandy, loamy sand, or fine sandy loam soils. Do not use on cotton planted in furrows.

Arkansas, Louisiana, and Mississippi Only: Use 1 lb. per acre on sandy loam soils low in organic matter.

Arizona, California, and New Mexico Only: Cotton, corn, and grain sorghum may be planted the next spring. Do not plant treated areas to other crops until one year after last application. Do not use on sandy or coarse-textured soil of less than 1% organic matter.

Note: Do not apply within 60 days of harvest. Do not feed foliage from treated cotton plants or gin trash to livestock.

#### Combinations

Prowl®: Apply **FLUOMETURON 80WP** as a preemergence surface treatment as recommended on this label at 1.25-2.5 lbs. per acre after a preplant soil-incorporated treatment of Prowl for control of all weeds

listed on both labels. Apply Prowl as recommended on its label at 1-3 pts. per acre.

DSMA or MSMA: **FLUOMETURON 80WP** may be tank-mixed and applied as a directed spray to cotton from 3 inches high to first bloom. Use the broadcast rate of 1.25-2.5 lbs. of **FLUOMETURON 80WP** per acre and the necessary amount of DSMA or MSMA plus 0.5% (1 pt. per 25 gals.) of a surfactant that is approved by FDA for use on food and feed crops and is compatible with **FLUOMETURON 80WP + MSMA** when applied to cotton. When using this mixture, do not allow spray to contact cotton foliage. Do not graze treated areas or feed treated forage or gin trash to livestock. Adhere strictly to the dosage and limitations on the labels of the products.

#### CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2005.

**COMPEX®** is a registered trademark of Kalo Laboratories, Inc.

**PROWL®** is a registered trademark of BASF Ag.



HELENA CHEMICAL COMPANY  
225 Schilling Blvd., Suite 300  
Collierville, Tennessee 38017  
Telephone: 901/761-0050

March 1, 2005

U.S. Environmental Protection Agency  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
1801 South Bell Street  
Crystal Mall #2, Room 266A  
Arlington, VA 22202-4501

RE: Notification to Add Other Revisions

To Whom It May Concern,

Enclosed, you will find Helena Chemical Company's Notification for numerous products. Helena has updated the copyright company name, all trademarks, and registered trademarks from Helena Chemical Company to Helena Holding Company. This has been done on the following products:

- Helena Wettable Sulphur 5905-289
- Fyfanon 25WP 5905-293
- Sol Oil 97 5905-294
- Sol Oil Plus 5905-302
- Helena Citrusperse Sulfur 5905-350
- Omni Supreme Spray 5905-368
- Helena 90% Sulfur Dust 5905-422
- Helena Liquid Sulfur Six 5905-437
- 6lb. Sodium Chlorate Defoliant Desiccant 5905-461
- Atrazine 4L 5905-470
- Helena Bravo S 5905-472
- Helena 4.8lb. Diuron Flowable Herbicide 5905-482
- Helena 3.5lb. Liquid Copper Concentrate 5905-483
- Copper Z 4/4 Algicide/Herbicide 5905-486
- Dimethoate 4EC 5905-493
- Setre Fluometuron 80WP Herbicide 5905-494
- Setre Prowl Herbicide + Propanil 5905-495
- Setre 1.5lb. Benfluramin EC 5905-496
- 5lb. Dimethoate Systemic Insecticide 5905-497
- Brush Rhap Low Volatile 4-D Herbicide 5905-498

This should also be noted on all alternate brand names of the master labels.

In support of these Notifications, you will find the following:

EPA Form 8570-1  
1 copy of the revised label highlighted

If you have any questions, do not hesitate to call me at (901) 752-4420 or by fax at (901) 758-1694. Thank you for your assistance in this matter.

Sincerely,

Mandy K. Styles  
Product Registration Supervisor

