

5905-495

03/11/2005

1/7

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 5905-495	2. EPA Product Manager	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Setre Prowl Herbicide + Propanil	PM#	
5. Name and Address of Applicant (Include ZIP Code) Helena Chemical Company 225 Schilling Boulevard, Suite 300 Collierville, Tennessee 38017 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION MAR 11 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Other Revisions
Notification of Other Revisions per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input checked="" type="checkbox"/> On Label	
6. Manner in Which Label is Affixed to Product Self Adhesive		<input checked="" type="checkbox"/> Lithograph Paper glued Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Mandy K. Styles	Title Product Registration Supervisor	Telephone No. (include Area Code) (901) 752-4420
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registration Supervisor	
4. Typed Name Mandy K. Styles	5. Date 3/1/2005	

Prowl® Herbicide + Propanil

FOR DRY-SEEDED RICE

MANUFACTURED BY
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE TENNESSEE 38017

ACTIVE INGREDIENT:

- Propanil: 3', 4'-dichloropropionanilide 33.70%
- Pendimethalin: N-(1-Ethylpropyl)-3,4-dimethyl-2, 6-Dinitrobenzenamine..... 11.25%

INERT INGREDIENTS:..... 55.05%
TOTAL100.00%

This product contains the toxic inert ingredient monochlorobenzene.
This product contains 3 pounds propanil and 1 pound pendimethalin.

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Harmful if swallowed or absorbed through skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

- | | |
|----------------------|---|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eyelid open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor immediately for treatment advice. |
| IF ON SKIN: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor immediately for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have a person sip a glass of water if able to swallow. • Do not give anything to an unconscious or convulsing person. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of emergency, call ChemTrec at 1-800-424-9300.

NOTE TO PHYSICIAN: This product contains monochlorobenzene.

SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO.: 5905-495
EPA EST. NO.:

NET CONTENTS:

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves, such as butyl rubber 14 mils, neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils. Gloves should be replaced when permeation or degradation is detected.
- Shoes plus socks
- Protective eyewear (goggles, face shield or safety glasses)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows

workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves, such as butyl rubber 14 mils, neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils. Gloves should be replaced when permeation or degradation is detected.
- Shoes plus socks
- Protective eyewear (goggles, face shield or safety glasses)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and shrimp. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use, pour, spill or store near heat or open flame.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE

Do not use, pour, spill or store near heat or open flame. Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. If entire contents are not used, protect remaining material from moisture and heat. Resealing by rolling top down and storage under roof are recommended. In the event of spillage, scrape up material and dispose of in accordance with **DISPOSAL**. Repackage and relabel usable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

DISPOSAL

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

PROWL® HERBICIDE + PROPANIL is a postemergence treatment which combines the direct contact action of propanil and the residual activity of pendimethalin (PROWL). Since the residual activity of pendimethalin provides pre-emergence control for certain annual grasses which can germinate after this pre-mix treatment is applied, flooding after application can be delayed.

For maximum weed control with this treatment, it is important to carefully follow the directions below for (1) adequate spray coverage of weeds and soil and (2) proper timing of application, when barnyard grass (watergrass) is in the 1 to 3 leaf stage of growth with an occasional 4 leaf plant (make application when sprangletop is less than 1/2 inch in height).

Do not use in California. **DO NOT** use this treatment in water seeded rice.

SURFACE WATER ADVISORY

This product may contaminate water through runoff following rainfall events and by seepage through levees. This product has a high potential for runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Levees should be constructed with adequate time prior to chemical application so that they are compacted to reduce seepage and to hold a 3-6 inch flood (2001 Mississippi Rice Growers Guide). Other guidance is located at <http://agronomy.ucdavis.edu/uccerice/water/seep.htm> and from the document "Closed Rice Water Management Systems" from the National Resource Conservation Service of USDA. The University of Arkansas Rice Production Book (<http://www.uaex.edu/other-areas/publications/html>) also provides information concerning levee production.

GROUNDWATER ADVISORY

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical prior to flooding may result in some shallow ground water contamination due to cracks in the subsoil of the rice paddy.

SPRAYING INSTRUCTIONS

AERIAL APPLICATION

For aerial application, apply the recommended rate in 10 to 12 gallons of water per acre to insure adequate coverage. Avoid drift or accidental application from turning aircraft on cotton, soybeans, corn, safflower, seedling legumes, vegetables, orchards, vineyards, gardens, shrubs, and ornamentals. To minimize drift, **DO NOT** apply during periods of gusty winds or winds in excess of 5 mph. Uniform spray distribution may not be achieved when wind velocity is greater than 5 mph. It is recommended that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory information. This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE:

Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

GROUND APPLICATIONS

For ground equipment, apply the recommended rate in 15 to 25 gallons of water per acre to ensure adequate coverage. Use a properly calibrated low-pressure (20 to 40 psi) sprayer equipped with 8002 or larger size Tee-Jet or comparable nozzles to achieve uniform spray distribution and minimize foaming. Nozzle screens must be no finer than 50 mesh. DO NOT apply during periods of gusty winds or when wind velocity is greater than 10 mph.

EARLY POSTEMERGENCE APPLICATIONS

The seedbed should be firm and free of clods and trash. The seedbed must be prepared to allow for good seed coverage. Previous crop residues should be thoroughly mixed into the soil to a depth of 4 to 6 inches by plowing or disking before planting rice.

Uniformly apply recommended treatment by aircraft or ground equipment after rice emergence, according to spraying instructions above, when barnyardgrass is in the 1 to 3 leaf growth stage with an occasional 4 leaf plant (make application when sprangletop is less than 1/2 inch). THE GROWER SHOULD INSPECT FIELDS FREQUENTLY TO CHECK GROWTH OF BARNYARDGRASS AND/OR SPRANGLETOP TO DETERMINE PROPER APPLICATION TIMING. Timing of applications should be based on the growth stage of barnyardgrass and/or sprangletop and not on the growth stage of rice. If rice is too small to maintain a flood on the field, the treatment can still be applied since flooding can be delayed because of the residual activity of pendimethalin.

While the residual activity of pendimethalin allows flooding to be delayed, proper water management practices must be followed for normal rice growth. Flooding should not be delayed if weeds begin to develop after application.

Since soil and weeds must be completely exposed to spray coverage, no flood water should be on field at time of application. If necessary, fields may be flushed prior to treatment to produce vigorous rice and weed growth.

Since the residual activity of pendimethalin is activated by moisture, pendimethalin is most effective in controlling emerging weeds when adequate rainfall or irrigation (flush) is received within 7 days after application.

For maximum direct contact activity delay application if there is a chance of rain within 6 to 8 hours.

WEED SPECIES CONTROLLED

PROWL® HERBICIDE + PROPANIL provides direct contact control of the weeds listed below:

- Barnyardgrass (*Brachiaris* spp.)
- Crabgrass, large (*Digitaria sanguinalis*)
- Croton, woolly (*Croton capitatus*)
- Foxtail (*Setaria* spp.)
- Goosegrass (*Eleusine indica*)
- Gulf Cockspur (*Echinochloa crus-pavonis*)
- Hoorahgrass (*Fimbristylis miliaceae*)
- Mexican weed (*Caperonia palustris*)
- Paragrass (*Panicum purpurascens*)
- Pigweed, redroot (*Amaranthus retroflexus*)
- Redweed (*Melochia corchorifolia*)
- Sour Dock (*Rumex crispus*)
- Spearhead (*Rhynchospora comiculata*)
- Tall Indigo or Coffee Bean (*Sesbania exalta*)
- Texas Millet (*Panicum texanum*)
- Wiregrass (*Eleocharis* spp.)

The following grass species are controlled by the residual activity of **PROWL® HERBICIDE + PROPANIL** at the rates recommended for each soil texture listed below:

- Barnyardgrass (watergrass) (*Echinochloa crus-galli* or *Echinochloa colonum*)
- Crabgrass (*Digitaria* spp.)
- Signalgrass (*Brachiaris platyphylla*)
- Sprangletop (*Leptochloa* spp.)

BROADCAST RATE PER ACRE OF PROWL® HERBICIDE + PROPANIL

Soil Texture	PROWL® HERBICIDE + PROPANIL
COARSE	
sandy loams	3 qts.
MEDIUM	
sandy clay loams, *sandy clays, loams, silts, silt loams	4 qts.
FINE	
silty clay loams, *clay loams, silty clays, clays	4 qts.

1) The addition of 0.75 lbs. of propanil is necessary to obtain maximum contact kill.

*Sometimes considered transitional soils and may be classified as either medium or fine textured soils.

Water drained from treated rice fields must not be used to irrigate other crops or released within 1/2 mile upstream of a potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of a potable water intake in a standing body of water such as a lake, pond or reservoir.

Do not apply to fields where commercial catfish farming is practiced, and do not drain water from treated fields into areas where catfish farming is practiced.

Do not apply to fields where commercial crayfish farming is practiced and do not drain water from treated fields into area where crayfish farming is practiced.

DO NOT apply this mixture within 14 days before or after insecticide applications because serious damage to rice may occur.

- DO NOT** apply in liquid fertilizer.
- DO NOT** make more than one application per season.
- DO NOT** bale or use rice straw for feed or bedding.
- DO NOT** discharge water from treated areas for a period of 30-days after treatment.
- DO NOT** rotate crops other than rice for 30-days following application.
- DO NOT** harvest rice grown within 60-days following treatment.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES
Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

6/7

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2005.

PROWL® is a registered trademark of American Cyanamid Company.



HELENA CHEMICAL COMPANY

225 Schilling Blvd., Suite 300
Collierville, Tennessee 38017
Telephone: 901/761-0050

March 1, 2005

U.S. Environmental Protection Agency
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
1801 South Bell Street
Crystal Mall #2, Room 266A
Arlington, VA 22202-4501

RE: Notification to Add Other Revisions

To Whom It May Concern,

Enclosed, you will find Helena Chemical Company's Notification for numerous products. Helena has updated the copyright company name, all trademarks, and registered trademarks from Helena Chemical Company to Helena Holding Company. This has been done on the following products:

- Helena Wettable Sulphur 5905-289
- Fyfanon 25WP 5905-293
- Sol Oil 97 5905-294
- Sol Oil Plus 5905-302
- Helena Citrusperse Sulfur 5905-350
- Omni Supreme Spray 5905-368
- Helena 90% Sulfur Dust 5905-422
- Helena Liquid Sulfur Six 5905-437
- 6lb. Sodium Chlorate Defoliant Desiccant 5905-461
- Atrazine 4L 5905-470
- Helena Bravo S 5905-472
- Helena 4.8lb. Diuron Flowable Herbicide 5905-482
- Helena 3.5lb. Liquid Copper Concentrate 5905-483
- Copper Z 4/4 Algicide/Herbicide 5905-486
- Dimethoate 4EC 5905-493
- Setre Fluometuron 80WP Herbicide 5905-494
- Setre Prowl Herbicide + Propanil 5905-495
- Setre 1.5lb. Benfluralin EC 5905-496
- 5lb. Dimethoate Systemic Insecticide 5905-497
- Brush Rhap Low Volatile 4-D Herbicide 5905-498

This should also be noted on all alternate brand names of the master labels.

In support of these Notifications, you will find the following:

- EPA Form 8570-1
- 1 copy of the revised label highlighted

If you have any questions, do not hesitate to call me at (901) 752-4420 or by fax at (901) 758-1694. Thank you for your assistance in this matter.

Sincerely,

Mandy K. Styles
Product Registration Supervisor

