PM23 5905-502



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

FEB 2 4 1994

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

BEVERLY NEALE
HELENA CHEMICAL COMPANY
6075 POPLAR AVENUE SUITE 500
MEMPHIS, TN 38119

Subject:

Label Amendment Submission of 9/16/93 Response to PR Notice 93-7

EPA Reg. No. 5905-502

WEED-RHAP A4-MCPA HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.

Page 2

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Sim Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division

Beverly Neale HELENA CHEMICAL CO 6075 POPLAR AVE SUITE 500 MEMPHIS TN 38119

Comment for: EPA Reg Nr.5905-502 WEED-RHAP A4-MCPA HERBICIDE

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

Remove the statement "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application." from its current position within the Agricultural Use Box and place it above the Agricultural Use Box.

The circled statement(s) in the Non-Agricultural Use Requirements box contains errors. Use each statement exactly as it appears on your original label.

Correct the typographical errors circled on your proposed label.

WEED RHAP A4-MCPA

MCP AMINE HERBICIDE

For the Selective Control of Many Broadleaf Weeds in Flax, Small Grains, Rice, Peas, Clover, Alfalfa and Grass Pastures;

For Whitebrush Control; And For Weed Control in Non-Crop Areas

ACTIVE INGREDIENTS:

2-Methyl-4-Chlorophenoxyacetic Acid, dimethylamine salt	52.2%
INERT INGREDIENTS:	<u>47.8%</u>
Total	100.0%

Equivalent to 42.6% or 4 pounds per gallon of MCPA.

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle, (If you do not understand the label, find someone to explain it to you in detail.

STATEMENT OF PRACTICAL TREATMENT

<u>IF SWALLOWED:</u> Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF IN EYES: Flush with plenty of water for at least thirty (30) minutes. Get medical attention. IF ON SKIN: Wash contaminated areas thoroughly with soap and water. Get medical attention if irritation persists.

ACCEPTED
WILL CUMMENTS

In EPA Letter Dates

FEB 24 1994

EPA REG. NO.: 5905-502

EPA EST. NO.:

Under the Federal Insecticide. Fundicide, and Redesticide Act an amended, for the posticide reministrational RPA Reg. No.

NET CONTENTS:

MANUFACTURED BY HELENA CHEMICAL COMPANY MEMPHIS, TN 38119

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful if swallowed. May cause skin irritation. Avoid breathing spray mist.

Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTION EQUIPMENT

Applicators and other handlers must wear:

Long-sleeved shirt and long pants Waterproof gloves Shoes plus socks Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Usenshould: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Drift or run-off may adversely affect nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters or rinsate. Do not contaminate water intended for irrigation or domestic purposes. Do not apply when weather conditions favor drift from target sight.

GROUND WATER CONTAMINATION STATEMENT

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

CLEANING OF EQUIPMENT

When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away

from wells and other water sources.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls

Waterproof gloves

Shoes plus socks

Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets out of treated area until sprays have dried:

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Do not store this product near fertilizers, seeds, insecticides, or fungicides. Protect product from freezing. If subjected to freezing conditions, warm to 70°F and agitate. This will return product to original condition. Reclose all partially used containers by thoroughly tightening the screw cap. Damaged or leaking containers which cannot be used immediately, should be transferred to suitable sound containers and properly marked. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal."

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities. To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

7411

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

PESTICIDE DISPOSAL: Pesticide waste are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to the spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL PRODUCT INFORMATION

USE PRECAUTIONS: MCP Amine Herbicide is injurious to most broadleaf pants. Salts are the least volatile forms of MCP and do not release enough vapors from treated areas to reduce yield of adjacent susceptible crops. This product is a salt of MCP. Susceptible plants may, however, be injured by physical drift from the application of this product. Do not apply directly to or otherwise permit even minute amounts (such as spray drift) to contact fruit trees, vegetables, flowers, ornamentals, or other desirable plants susceptible to MCP. DO NOT APPLY IN THE VICINITY OF COTTON, GRAPES, TO-BACCO OR TOMATOES. Do not use in or near a greenhouse. Do not spray when wind is blowing towards susceptible crops or ornamental plants.

Do not apply this product through any type of irrigation system.

Use coarse sprays to minimize spray drift since, under certain weather conditions, fine spray droplets may drift a mile or more. A spray thickening agent, such as Nalco-Trol® may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on the product label.

(®, Nalco Chemical Company)

With ground equipment spray drift can be lessened by:

- 1. Keeping the spray boom as low as possible.
- 2. Applying 20 gallons or more of spray volume per acre.
- 3. Using large-droplet producing nozzles.
- 4. Using no more than 20 psi spray pressure at the nozzles.
- 5. Spraying only when wind velocity is low.
- 6. Stopping all spraying when wind exceeds 8 MPH.

Do not use hollow cone-type nozzles or other nozzles that produce large amounts of line spray droplets.

With aerial equipment, drift can be minimized by:

- 1. Using no more than 20 psi spray pressure at nozzles.
- 2. Using nozzle types and positions which do not produce fine spray droplets.
- 3. Using a spray boom no longer than 3/4 the wing or rotor span of the aircraft.
- 4. Spraying only when wind is less than 6 MPH.

Determine air movement and direction before foliar application. Use a smoke generator or other means at or near the application site for the detection of air movement, air stability, or temperature inversions. Such a condition exists when there is little or no wind and air temperature is lower near the ground than at higher levels. Use appropriate drift control measures or avoid application when smoke is moving toward nearby desirable susceptible plants or sensitive areas.

Excessive amounts of this herbicide in the soil may temporarily inhibit seed generation or plant growth. Violent windstorms may move soil particles. If MCP is on soil particles and they are blown onto susceptible plants, visible symptoms may appear. Serious injury is unlikely. The hazards of movement of MCP on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.

To avoid injury to desirable plants, do not handle or apply other agricultural chemicals with the same equipment used for MCP unless it has been thoroughly cleaned with a suitable chemical cleaner.

Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station or Extension Service Weed Specialists for advice in selecting treatments from this label to best fit local conditions. Be sure that the use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

WHERE TO APPLY: MCP Amine Herbicide is used to control broadleaf weeds in flax, peas, small grains, established alfalfa and red clover, grass pastures, rangeland, and many non-crop areas. This product will kill or control the following weeds in addition to many other noxious plants susceptible to MCP.

Most Susceptible Weeds (most areas):

Arrowhead	Beggartick	Mustard	Pennycress (field)
Galinsoga	Goatsbeard	Ragweed	Shepherdspurse
Marshelder	Mint (dragonhead)	Water Plantain	Witchweed
Puncturevine	Redstem	and many	Croton (goatweed)
Sneezeweed	Stinkweed	other species	Lambsquarters
Yellow Rocket	Burdock	Cocklebur	Pepperweed (field)
(winter cress)	Jimsonweed	Kochia	Sicklepod

Less Susceptible Weeds:

Burcucumber	Hoary Cress	Honeysuckle	Sunflower
Hempnettle	Sow Thistle	Stinging Nettle	Dandelion
Purslane	Wild Radish	Catsear	Pigweed
White Top	Canada Thistle	Plantain	Vetch
Buttercup			

WHEN TO APPLY: Apply the product during warm weather when weeds are young and actively growing. Uniform spray coverage is necessary for best results. Generally the low dosages listed will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is difficult, the higher dosages will be needed.

HOW TO APPLY:

PREPARING THE SPRAY: Mix the MCP Amine Herbicide with water unless otherwise indicated

on the label. Fill the mixing tank or spray tank with approximately 1/2 the amount of water to be used for spraying. Add the required amount of product with agitation. NOTE: adding oil, wetting agents, or other adjuvants to the spray may reduce selectivity to crops, possible causing crop damage.

SPRAYING CONDITIONS: Use a moderate spray pressure of approximately 20 psi. Enough spray volume should be used to give good coverage of the weeds being sprayed. This is usually 5 to 30 gallons by ground application or 3 to 10 gallons by air. Higher spray volumes may be useful in some situations to provide better spray coverage when weeds are especially dense and/or to reduce spray drift.

WEED CONTROL APPLICATIONS

FLAX: For control of susceptible weeds, use 1/2 pint of MCP Amine Herbicide per acre. Treat flax when 3 to 6 inches tall before buds begin to form. Rates of 3/4 to 1 pint per acre may be needed to control some weeds such as wild buckwheat, smartweed and thistles but flax may be injured at these rates. If seeding grasses such as foxtail are a problem, DOWPON® M Grass Killer may be used alone or in combination with MCP Amine Herbicide. Flax varieties may vary in tolerance to DOWPON® M and MCP. Observe the most restrictive limitations, precautions, and directions for use on the product's labeling. Consult Local Extension Service or University Specialists for advise on herbicide use on flax. Do not forage or graze meat animals on treated areas within seven (7) days of slaughter.

WEED CONTROL IN SMALL GRAINS: NOTE: Do not permit dairy animals, or meat animals being finished for slaughter to forage or graze treated grain fields within two (2) weeks after treatment. Do not forage or graze meat animals on treated areas within seven (7) days of slaughter.

Wheat, Barley, Oats and Rye not underseeded to Legumes: Use 1/2 to 1 pint MCP Amine Herbicide per acre for the most susceptible weeds when grains are in the 3 to 5 leaf stage or up to early boot stage. Apply when weeds are very small for best results. Use up to 3 pints MCP Amine Herbicide per acre for the less susceptible weeds after crop has tillered but not later than early boot stage. Do not apply during the boot to dough stage of the grain.

Wheat, Barley, Oats and Rye underseeded to Alfalfa, Birdsfoot Trefoil, Lespedeza, Red White, or Ladino Clover: For emergency control of susceptible weeds such as mustard and lambsquarters, use 1/3 to 1/2 pint of MCP Amine Herbicide per acre as soon as possible after weeds emerge, but after seedling grain is at least 3 inches tall. The grain and weeds should form a protective canopy over the clover seedlings. Treatment may be made up to the full tiller stage, but not in the boot to dough stages of the grain. To control certain weed species, 1 pint per acre may be needed. Do not use where small seeded legumes, especially vetch, sweet clover, trefoil or alfalfa, are seeded unless injury to them can be tolerated.

RICE: For control of certain broadleaf weeds and sedges, apply 2 to 3 pints of MCP Amine Herbicide in 10 to 30 gallons of water per acre. Treat when rice is fully tillered and 6 to 8 inches above water, usually 35 to 65 days after planting. Do not treat when rice is in the early seedling, boot, or early heading stages or when air temperature is over 90°F. Some rice varieties may be less tolerant to MCP than others. Consult your Agricultural Experiment Station or Extension Service Weed Specialists for appropriate rates and timing of MCP spray and on the tolerance of rice varieties. Do not grow crayfish or catfish in treated rice fields.

RICE IN CALIFORNIA: For early season broadleaf weed and sedge control in rice, apply 1 to 1½ pints of MCP Amine Herbicide per acre by air in 8 to 10 gallons of total spray volume, when rice is in the 3 to 4 leaf stage, and with well established root systems, usually 18 to 24 days after planting. Applications made during hot weather or at the higher dosage rate may cause temporary stunting of the rice. For maximum effectiveness, the weeds should be exposed at time of treatment. If retreatment is necessary, usually 35 to 65 days after planting, the total amount of MCP Amine Herbicide per acre, applied in the two applications, must not exceed 3 pints per season. Do not grow crayfish or catfish in treated rice fields.

PEAS: Use on peas to control susceptible weeds only where recommended by local Agricultural Extension Service or experiment station specialists. Use 1/4 to 1/3 pint of MCP Amine Herbicide in at least 15 gallons of water per acre when peas are 3 to 7 inches tall, before first flowering, and weeds are small. Higher rates of 1/2 to 3/4 pint per acre may be used to improve control of difficult weeds, but crop injury is more likely to occur. Do not apply if peas are taller than 7 inches or when they are stressed for lack of soil moisture. Do not apply when air temperature is over 90°F. Do not feed treated vines to livestock. Do not forage or graze treated pea fields. NOTE: MCP can cause injury and delayed maturity in the pea crop.

ESTABLISHED ALFALFA OR RED CLOVER: For control of yellow rocket and other susceptible annual weeds such as pennycress or tanweed, use 1 pint of MCP Amine Herbicide per acre. Apply in late fall after frost has killed legume top-growth and legumes are dormant. The temperature at time of spraying should be above 40°F.

GRASS PASTURES (INCLUDING GRASS GROWN FOR SEED): Use 2 to 4 pints of MCP Amine Herbicide per acre. Apply when weeds are small and actively growing for best results. Use the higher rates for whitetop., Canada thistle, buttercup and other more difficult to control weeds. Spray perennials in early bud to full bloom stage and regrowth in fall. Legumes may be injured or killed. Do not use this treatment where legumes, especially alfalfa, are present and desirable. In some areas bent, buffalo, carpet and St. Augustine grasses may be susceptible to injury. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage where grass seed production is desired.

CONTROL OF WHITEBUSH ON RANGELAND: Use 3 pints of MCP Amine Herbicide with 1 gallon of diesel oil and enough water for good spray coverage, usually 6 to 10 gallons per acre by aircraft. Apply in spring or fall when plant foliage is well developed and actively growing. Spraying during bloom is recommended but not immediately after shedding of blossoms. Retreatments may be needed in succeeding years. NOTE: First mix the MCP Amine Herbicide in the water, then add the oil with agitation. Spray must be kept agitated during spraying to avoid separation in the tank.

NON-CROP AREAS: For control of susceptible weeds in fencerows, fallowlands, rights-of-way, roadsides, and similar areas, use 6 pints of product per acre in enough water to give sufficient coverage. NOTE: on fallow land, do not plant any crop for 3 months after treatment or until chemical has disappeared from soil.

SPOT TREATMENT: For weed control in pastures, rangelands and in non-crop areas such as fencerows, farmyards, shelter-belts and roadsides: Use ¼ pint of MCP Amine Herbicide in 3 gallons of water to control weeds such as Canada thistle, whitetop, meadow buttercup and Texas blueweed. Apply when weeds are growing vigorously, usually prior to bloom, and spray fall regrowth if necessary.

118f11

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man, or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damage and in no event shall damages or any other recovery of any kind against the Company exceed the price of the product which causes the alleged loss, damage, injury, or other claim. The Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income.

The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability, and remedies.