5905-538

6311/2065 Form Approved.

OMB	No.	2070-0		el exp	ires	<u> 2-28</u>
	_		 			

_	Registration
	Amendment
	Other

OPP Identifier Number

Environmental Protection Agenc Washington, DC 20460			ncy		~	Amend Other					`	
		Applicati	on for F	Pesticio	de - Sect	tion	1			<u> </u>		-
1. Company/Product Num	nber		•	2. EPA f	roduct Man	ager			3. Pro	posed C	lassificat	tion
5905-538  4. Company/Product (Name) Pro-Mate Dimension 0.22% Plus Fertilizer									~	None	R	lestricted
				e Eva	adited Day					CIEDA C		3/-1/21
5. Name and Address of Applicant (Include ZIP Code) Helena Chemical Company 225 Schilling Boulevard, Suite 300 Collierville, Tennessee 38017					6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No.							
Check if t	his is a new address			Produ	ct Name _							
			Sect	ion - I	1							
Amendment - Exp	ain below.	datad			Final printed Agency lett "Me Too" A	er dat	ed	se to			CATIC	
Notification - Expl					Other - Expi	•			•	null T	1 200	JS
152.46, and no other char U.S.C. Sec. 1001 to willful	sions per PR Notice 98-10, nges have been made to the ly make any false statement product may be in violation	e labeling or t nt to EPA. I fu	he confiden urther under I may be su	tial statem stand that ubject to e	nent of formul if this notifican forcement a	a of th	is product. s not consist	I unde lent wit	rstand th the t	that it is erms of I	a violation PR Notice	n of 18 98-10
			Sect	ion - II	<u> </u>							
1. Material This Product \	1		Water	Soluble Pa			2. Type o	f Cont	niner	<u> </u>		
Child-Resistant Packaging  Yes  No  No  Volume  No			Yes Mota V No Glass			etal estic						
* Certification must be submitted If "Yes" No. per container			If "Yes" Packag	es" No. per Paper Other (Specify)								
3. Location of Net Conten	ts Information  Container	4. Size(s) Re	tail Contair	ier		5. Lo	On Lab		irection	าร		
6. Manner in Which Label is Affixed to Product Self Adhesive Lithor Paper Stend			graph Other glued iiled									
			Secti	on - I\								<u></u>
1. Contact Point (Comple	te items directly below fo	r identification	on of individ	dual to be	contacted,	if nec	essery, to p	roces.	s this .	applicati	on.)	<del> </del>
Name Mandy K. Styles			Title Product I	Registrat	ion Supervis	sor		1	phone 1) 752-		iude Are	a Code)
	itements I have made on any knowlingly false or n le law.		i all attachr						1	Recei	Applications  tampe	
2. Signature			3. Title Product Registration Supervisor									
4. Typed Name  Mandy K. Styles			5. Date	3/1/	2005							

A/g NOTIFICATION

X-X-X	
GUARANTEED FERTILIZER ANALYSIS	
TOTAL NITROGEN (N)*X.XX	%
X.XX % Ammoniacal Nitrogen	
AVAILABLE PHOSPHATE (P2O5)XX	%
SOLUBLE POTASH (K2O)X.XX	%
Derived from: xxxxxxx, xxxxx, xxxxx.	
Chlorine (CI) MaxX.XX	%
*x.xx% Slowly Available Nitrogen from polycoated sulful	
coated urea.	
F224	
*For approved fertilizer variations by state, contact	

## Pro-Mate Dimension 0.22 %Plus

### **Fertilizer**

Representative.

#### **Weed Control with Premium Lawn Food**

your local Helena Chemical Company

Pro-Mate® Dimension® 0.22% Plus Fertilizer is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns, ornamental landscape areas, and ornamental turfs, including golf course fairways, roughs, and tee boxes. Not for use on turf being grown for sale or other commerical use, or for commercial seed production, or for research purposes.

In the state of New York, this product may <u>only</u> be used by Professional Turf Applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

#### **ACTIVE INGREDIENT:**

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA Reg. No 5905-538

EPA Est. No.

**Net Contents:** 

KEEP OUT OF REACH OF CHILDREN CAUTION

MANUFACTURED BY: MAR 1 1 2005
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TN 38017

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION! CAUSES EYE IRRITATION. HARMFUL IF
INHALED. Avoid contact with eyes or clothing. Avoid
breathing dust. Wash thoroughly with soap and water
after handling. Remove contaminated clothing and wash
before reuse. Prolonged or frequently repeated skin
contact while handling this material may cause allergic
reaction in some individuals.

#### **FIRST AID**

#### IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

#### **HOT LINE NUMBER**

For additional information, in case of emergency, call 800-424-9300.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not

contaminate water when disposing of equipment washwaters.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in any manner inconsistent with its labeling. In New York State, this product may <u>only</u> be used by Professional Turf Applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

#### STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**STORAGE:** Store this product only in its original container in a dry, cool, secured storage area.

PRODUCT DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state or local procedures.

EMPTY CONTAINER DISPOSAL: Do not reuse bag. Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**SPILL:** In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

#### **GENERAL INFORMATION**

This product is a selective herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

#### **USE PRECAUTIONS**

**Turfgrass Safety** - This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns

and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

#### Cool-Season Grasses

Common Name	Scientific Name
Bentgrass, creeping*	Agrostis palustris
Bluegrass, Kentucky	Poa pratensis
Fescue, fine**	Festuca rubra
Fescue, tall	Fescue arundinacea
Ryegrass, perennial	Lolium perenne

#### Warm-Season Grasses

Common Name	Scientific Name			
Bahiagrass	Paspalum notatum			
Bermudagrass	Cynodon dactylon			
Buffalograss***	Buchloe dactyloides			
Carpetgrass	Axonopus affinis			
Centipedegrass	Eremochloa			
, -	ophiuroides			
Kikuyugrass	Pennisetum			
	clandestinum			
St. Augustinegrass	Stenotaphrum			
	secundatum			
Zoysiagrass	Zoysia japonica			


\*Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside' and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

\*\*Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Biljart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and 'Waldina'. Not all varieties of fine fescue have been tested.

\*\*\*DO NOT use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to Pro-mate® Dimension® 0.22%with Fertilizer. It is suggested that the user evaluate the response of turfgrass varieties to this product, on a limited area, at recommended use rates, prior to initiating large-scale use.



#### Reseeding, Overseeding or Sprigging

Reseeding, overseeding or sprigging of treated areas should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent the establishment of desirable turfgrasses.

When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed.

For best results, use mechanical or power seeding equiptment (slit seeders) designed to give good seed to soil contact.

#### **GOLF COURSE PUTTING GREENS**

DO NOT apply this product to golf course putting greens.

#### Other Use Precautions

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification and verticutting should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

- Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.

# APPLICATION DIRECTIONS Control of Crabgrass

#### **Pre-emergence and Early Postemergence Control**

- This product provides "preemergence" control of crabgrass (including the large, smooth and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.
- It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground; however, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

### **Application Frequency and Timing**

- This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer or fall.
- DO NOT apply more than 5.22 pounds of this product per 1,000 sq ft per application, and no more than 15.66 pounds of this product per 1,000 sq ft per year.

5/8

• In the state of New York, this product may only be used by professional applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

#### Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

- Use Program #1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- Use Program #2 for preemergence control at sites
  where (a) turfgrass is cut low (e.g., golf fairways),
  and (b) turfgrass maintenance or weed control has
  not been conducted during the previous year. This
  program provides 4-6 months of preemergence
  crabgrass control. This program may also be used
  for early postemergence control of crabgrass.

Subsequent, sequential pre-and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be divided between two applications made 6-10 weeks apart and prior to crabgrass emergence.

## Table 1: Recommended Single Application Use Rates\*

(Pounds of this product per 1,000 sq ft)

PROGRAM USE (turigrass cut) REGION	#1 Preemergence (high-cut turf)	#2 Preemergence (low-cut turf) Postemergence (high-cut turf)	#3 Preemergence (low-cut turf) Postemergence (low-cut turf)
North	1.30 lb/	1.91 lb/	2.61 lb/
	1,000 sq. ft.	1,000 sq. ft.	1,000 sq. ft.
Transition	1.91 lb/1,000 sq.ft.	2.61 lb/ 1,000 sq. ft.	3.82 lb/ 1,000 sq. ft.
South	2.61 lb/	3.82 lb/	5.22 lb/
	1,000 sq. ft.	1,000 sq. ft.	1,000 sq. ft.
Coastal South	3.82 lb/	5.22 lb/	5.22 lb/
	1,000 sq. ft.	1,000 sq. ft.	1,000 sq. ft.**

Note: 2.61 lb per 1,000 square feet is equal to 0.25 lb. active ingredient per acre.

\*DO NOT apply more than 5.22 lb of this product per 1,000 sq ft per application, and no more than 15.66 lb of this product per 1,000 sq ft per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. See Map of United States.

\*\* May require split or sequential applications for full control.

In the state of New York, this product may <u>only</u> be used by Professional Turf Applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

North: all areas not designated below. Transition: DE, KS, KY, MD, MO, NJ, VA,

southeastern PA, southern areas of IL, IN, OH, &

coastal areas of CT, NY & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NC, NM,

NV, OK, SC, TN & TX.

Coastal South: HI, FL & southern coastal areas of AL,

GA, LA, MS, NC, SC & TX.

#### **Fall Applications**

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application may be followed by an appropriate spring application to provide season-long control.

#### **Tips for Improved Control**

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few

6/8

days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

# Control of Other Grasses and Broadleaf Weeds Spring Application

Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

Barnyardgrass

Ryegrass (annual & perennial)

Bluegrass (annual)

**Smutgrass** 

Crowfootgrass

Lespedeza (common)

Foxtail (yellow &

Oxalis (buttercup, creeping, &

green)\* Goosegrass yellow woodsorrel) Purslane (common)

Kikuyugrass Speedwell (corn)

Spurge (prostrate & spotted)

#### **Fall Applications**

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

Bluegrass (annual)

Geranium (Carolina)

Bittercress Chickweed Henbit Parsley-piert
Pineappleweed
Shepherdspurse

# APPLICATION EQUIPMENT AND INSTRUCTIONS Apply PRO-MATE DIMENSION 0.22%Plus Fertilizer

with drop or rotary spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to

ensure equipment is functioning properly and applying uniform distribution of granules.

CONDITIONS OF SALE - LIMITED WARRANTY
AND LIMITATIONS OF LIABILITY AND REMEDIES
Read the Conditions of Sale - Warranty and
Limitations of Liability and Remedies before using
this product. If the terms are not acceptable, return
the product, unopened, and the full purchase price
will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product. including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

- Refund of the purchase price paid by buyer or user for product bought, or
- 2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2005.

<sup>\*</sup>also controlled at the "Program 2" rates.

PRO-MATE® is a registered trademark of Helena Holding Company.

DIMENSION® is a registered trademark of Rohm & Haas Company.



#### HELENA CHEMICAL COMPANY

225 Schilling Blvd., Suite 300 Collierville, Tennessee 38017 Telephone: 901/761-0050

March 1, 2005

U.S. Environmental Protection Agency Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) 1801 South Bell Street Crystal Mall #2, Room 266A Arlington, VA 22202-4501

RE:

Notification to Add Other Revisions

To Whom It May Concern,

Enclosed, you will find Helena Chemical Company's Notification for numerous products. Helena has updated the copyright company name, all trademarks, and registered trademarks from Helena Chemical Company to Helena Holding Company. This has been done on the following products:

Pro-Mate Barricade 0.435% Plus Fertilizer	5905-536
Pro-Mate Dimension 0.10% Plus Fertilizer	5905-537
Pro-Mate Dimension 0.22% Plus Fertilizer	5905-538
Pro-Mate Dimension 0.26% Plus Fertilizer	5905-539
Asset PGR	5905-540
Armicarb 100	5905-541
HM-2010	5905-542
Pro-Mate Barricade 0.22% Plus Fertilizer	5905-543
Extra Value Dusting Sulfur	5905-544
Sungro Combo Sulfur Dusting & Spraying	5905-545
Sungro 455 Soluble Oil	5905-546
435 Soluble Oil	5905-547
Empower 2 Granular Insecticide	5905-548
HardBall	5905-549
WildCard Xtra	5905-550
Pro-Mate Ronstar 1% Plus Fertilizer	5905-551
Double Up B+D	5905-552
Helena XL2G	5905-556
This should also be noted on all alternate brand name	es of the master labels.

In support of these Notifications, you will find the following:

EPA Form 8570-1

1 copy of the revised label highlighted

If you have any questions, do not hesitate to call me at (901) 752-4420 or by fax at (901) 758-1694. Thank you for your assistance in this matter.

Sincerely,

Mandy Tr. St.

Mandy K. Styles Product Registration Supervisor