

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

AUG 2 0 2010

Helena Chemical Company c/o Cheryl Wagner Wagner Regulatory Associates, Inc. P.O. Box 640 Hockessin, DE 19707

Subject: Notification: Revised Container Disposal Instructions per PR Notices 2001-6, 2007-1,

and 2007-4

Pro-Mate Dimension 0.22% Plus Fertilizer

EPA Reg. No. 5905-538

Your Re-Submission, Dated August 20, 2010

Dear Ms. Wagner:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notices (PRN) 2007-4, 2007-1 and 2001-6 for the subject product.

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4, 2007-1 and 2001-6 and finds that the label changes requested fall within the scope of the PR Notices. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at (703) 305-1243 or Susan Stanton of my staff at (703) 305-5218.

Sincerely,

Kathryn Montague, Product Manager 23

I Starton for

Herbicide Branch

Registration Division (7505P)
Office of Pesticide Program

, ,						2006	
Please read instructions on reverse	before com		Form	Approv	/ JMB No. 20	70-0060. Approval expires 05-31-98	
O PDA	United S Vironmental Pro Washington, I	otection A		X	Registration Amendme Other	İ	
	Applic	ation for	Pesticide - Se	ction	<u> </u>	· · · · · · · · · · · · · · · · · · ·	
1. Company/Product Number			2. EPA Product N			3. Proposed Classification	
5905-538			K. Montague				
4. Company/Product (Name)			PM#			☑ None Booksisted	
Pro-Mate Dimension 0.22% Plus Fertilizer			None Restricted				
5. Name and Address of Appli Helena Chemical Company c/o Wagner Regulatory Ass P.O. Box 640 Hockessin, DE 19707	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name AUG 2 0 2010						
Li Check if	this is a new address						
Amandment Cyntein bal		Sec	tion - II	al labale	o in connent to		
Amendment - Explain bel Resubmission in respons Notification - Explain belo	Agency let "Me Too" A	ter date Applicati	ion.				
2001-6 and EPA Regulations at 40 confidential statement of formula of	Notices 2007-4, 2007-1, CFR 156.10, 152.46, 15 of this product. I understant cation is not consistent w	and 2001-6. The state of the st	his notification is cons i, 156.146, and 156.19 clation of 18 U.S.C. S i PR Notices 2007-4, 2	56. No d Sec. 100	other changes ha of to willfully mak	as of PR Notices 2007-4, 20071- and ave been made to the labeling or the se any false statement to EPA. I product may be in violation of FIFRA	
1. Material This Product Will	Be Packaged In:						
Child-Resistant Packaging Yes* X No * Certification must	Unit Packaging Yes X No If "Yes" Unit Packaging wgt.	No. per If		aging . per ntainer	X	Metal Plastic Glass Paper	
be submitted						Other (Specify) HDPE lined bags	
3. Location of Net Contents Information X Label Container 50 lb. bag			etall Container	5	X On Lat	abel Directions pel peling accompanying product	
6. Manner in Which Label is A	affixed to Product	X Pap	ograph er glued nciled	<u> </u>	Otheradhe	sive backed label	
			tion - IV				
1. Contact Point (Complete ite		dentification of	of individual to be c	ontacte		والمراب أحاده والمستون والمشارك والمستون والمستون والمستون والمستون والمستون والمستون والمستون والمستون والمستون	
Name Cheryl Wagner	Title	Agent for He	elena Chemical Co	mpany		ne No. (Include Area Code) 4 8551	
I certify that the statements I hav I acknowledge that any knowingly both under applicable law.		ation all attachments	thereto are true, acc	urate ar	nd complete.	6. Date Application Received (Stamped)	
2. Signature 3. Title						• •	
Church Wagner Agent for H			elena Chemical Co	mpany	<u>, </u>		
4 Typed Name		5 Date					

Cheryl Wagner

August 20, 2010

Wagner Regulatory Associates, Inc.

P.O. Box 640 7460 Lancaster Pike, Suite 9 Hockessin, Delaware 19707

August 20, 2010

Document Processing Desk (NOTIF) ATTN: Ms. Kathryn Montague, PM 23 Registration Division (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, Virginia 22202-4501

Dear Ms. Montague:

Pro-Mate Dimension 0.22% Plus Fertilizer Re:

EPA Registration Number 5905-538

Notice of Revised Storage & Disposal Label Language

Wagner Regulatory Associates, Inc., on behalf of Helena Chemical Company, hereby notifies the Agency that the storage and disposal section of the subject label as been revised in accordance with PR Notice 2007-4, 2007-1, and 2001-6. Enclosed for the Agency's file is:

Letter from Helena Chemical Company authorizing Wagner Regulatory to serve as Agent

EPA Notification form (EPA Form 8570-1)

One copy of revised labeling

Please feel free to contact me at (302) 234-8551 if you have any questions or require additional information.

Respectfully submitted,

Cheryl Wagner

Agent for Helena Chemical Company

in Wagney

SPECIMEN LABEL

Pro-Mate® Dimension® 0.22 % **Plus Fertilizer**

Weed Control with Premium Lawn Food

Pro-Mate® Dimension® 0.22% Plus Fertilizer is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns, ornamental landscape areas, and ornamental turfs, including golf course fairways, roughs, and tee

Not for use on turf being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

In the state of New York, this product may only be used by Professional Turf Applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.l./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

ACTIVE INGREDIENT:

Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-

6-(trifluoromethyl)-S,S-dimethyl ester..... 0.22%

INERT INGREDIENTS:....99.78%

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA Reg. No 5905-538

EPA Est. 5905-NM-003, 5905-NC-018, 5905-FL-013, 5905-NE-009

Manufactured For: **Helena Chemical Company** 225 Schilling Blvd. Collerville, TN 38017

SN: 0206/0607

Net Contents: 50 lbs.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTIONI CAUSES EYE IRRITATION. HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction In some individuals.

FIRST AID

IF IN EYES:

- · Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- · Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

For additional information, in case of emergency, call 800-424-9300.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in any manner inconsistent with its labeling. In New York State, this product may only be used by Professional Turf Applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store this product only in its original container in a dry, cool, secured storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state or local procedures.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container, If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or

SPILL: In case of spill, sweep up material and dispose of material according to "product disposal* directions listed above.

d

GENERAL INFORMATION

This product is a selective herbicide that provides control of crabgress and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

USE PRECAUTIONS

Turfgrass Safety - This product may be used on seeded, sodded, or sprigged lawns and omamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanicalrelated stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses

Common Name Bentgrass, creeping* Bluegrass, Kentucky Fescue, fine** Fescue, tail Ryegrass, perennial

Scientific Name Agrostis palustris Poa pratensis Festuca rubra Fescue arundinacea Lolium perenne

NOTIFICATION AUG 2 0 2010

Warm-Season Grasses

Common Name Bahlagrass Bermudagrass Buffalograss*** Carpetgrass Centipedegrass Kikuyugrass St. Augustinegrass Zoysiagrass

Scientific Name Paspalum notatum Cynodon dactylon **Buchloe dectyloides** Axonopus affinis Eremochioa ophiuroides Pennisetum clandestinum Stenotephrum secundatum Zoysia japonica

*Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside' and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties.

Disclaimer: Always refer to the label on the product before using Helena or any other product. 16June2010 revised storage & disposal


SPECIMEN LABEL

APPLICATION DIRECTIONS

"Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Attanta', 'Banner', 'Beauty', 'Biljart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf' and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turigrass varieties for tolerance to Pro-mate® Dimension® 0,22% with Fertilizer. It is suggested that the user evaluate the response of turigrass varieties to this product, on a limited area, at recommended use rates, prior to initiating large-scale use.



Reseeding, Overseeding or Sprigging

Reseeding, overseeding or sprigging of treated areas should be delayed until 12 weeks from the time of application. Reseeding, overseeding or sprigging before 12 weeks after application may prevent the establishment of desirable turigrasses.

When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed.

For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

GOLF COURSE PUTTING GREENS

DO NOT apply this product to golf course putting greens.

Other Use Precautions

The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydroaerification and verticutting should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

- Apply this product directly to established lawns or ornamental turfs only. DO NOT apply to flowers, vegetables, shrubs, or trees.
- DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.
- DO NOT apply this product to pastures. Keep people and pets off treated areas until dust has settled.

Control of Crabgrass

Pre-emergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass (including the large, smooth and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs.
 - It can also provide "early postemergence" control of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground; however, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental turfs. Post-emergence crabgrass control will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So the practical benefit of this product's additional, early postemergence activity is that (compared to strictly preemergence crabgrass products) it can give the user a 2-4 week longer period of time to make applications and still control crabgrass.

Application Frequency and Timing

- This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer or fall.
- DO NOT apply more than 5.22 pounds of this product per 1,000 sq ft per application, and no more than 15.66 pounds of this product per 1,000 sq ft per year.
- In the state of New York, this product may <u>only</u> be used by professional applicators at no more than 5.22 lb./1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

Spring Applications

For single applications made in the spring or early summer, this product should be applied at the appropriate rate corresponding to one of the three control programs listed in Table 1 below, depending on the user's location, the turigrass mowing height, and whether the use is considered to be preemergence or early

postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

- Use Program #1 for preemergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program should not be used for early postemergence crabgrass control.
- Use Program #2 for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass.

Subsequent, sequential pre-and/or postemergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be divided between two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates*
(Pounds of this product per 1 000 so ft)

PROGRAM	#1	#2	#3
USE (turfgrass	Preemergence	Preemergence	Preemergence
cut)	(high-cut turf)	(low-cut turf)	(low-cut turf)
REGION		Postemergence (high-cut turf)	Postemergence (low-cut turf)
North	1.30 (b/ 1,000 sq. ft.	1.91 lb/ 1,000 sq. ft.	2.61 lb/ 1,000 sq. ft.
Transition	1.91 lb/1,000 sq.ft.	2.61 ib/ 1,000 sq. ft.	3.82 lb/ 1,000 sq. ft.
South	2.61 lb/	3.82 lb/	5.22 lb/
	1,000 sq. ft.	1,000 sq. ft.	1,000 sq. ft.

SPECIMEN LABEL

Coastal South	3.82 lb/	5.22 lb/	5.22 lb/	
	1,000 sq. ft.	1,000 sq. ft.	1,000 sq. ft.**	

Note: 2.61 lb per 1,000 square feet is equal to 0.25 lb. active ingredient per acre.

- *DO NOT apply more than 5.22 lb of this product per 1,000 sq ft per application, and no more than 15.66 lb of this product per 1,000 sq ft per year. Postemergence control is limited; see "Crabgrass Control" and
- "Precautions" sections above. Regions include areas listed below. See Map of United States.
- May require split or sequential applications for full control.

In the state of New York, this product may only be used by Professional Turf Applicators at no more than 5.22 lb/1000 sq. ft. per year (0.5 lbs. a.i./acre/year); and is prohibited from use in Nassau and Suffolk Counties.

North:

all areas not designated below.

Transition:

DE, KS, KY, MD, MO, NJ, VA.

Southeastern

PA, southern areas of IL, IN, OH, & coastal areas of CT, NY &

RI.

South:

AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN & TX.

Coastal South:

HI, FL & southern coastal areas of AL, GA, LA, MS, NC, SC &

Fall Applications

This product can also be applied in the late summer or early fall. (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application may be followed by an appropriate spring application to provide season-long control.

Tips for improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed

Control of Other Grasses and Broadleaf Weeds

Spring Application

Used as directed for crabgrass control in the spring, this product will also control (at the "Program 3" rates) or suppress (at the "Program 2" rates) the following weeds when applied prior to their emergence:

Barnyardgrass

Ryegrass (annual & perennial)

Bluegrass (annual)

Smutgrass

Crowfootgrass Foxtail (yellow & Lespedeza (common) Oxalis (buttercup, creeping, &

green)*

yellow woodscrrel)

Goosegrass

Pursiane (common)

Kikuyugrass

Speedwell (corn)

Spurge (prostrate & spotted)

"also controlled at the "Program 2" rates.

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) or suppress (at the 'Program 2' rates) the following weeds when applied prior to their emergence:

Bluegrass (annual)

Geranium (Carolina)

Bittercress

Parsley-piert

Chickweed

Pineappleweed

Henbit

Shepherdspurse

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply PRO-MATE® DIMENSION® 0.22%Plus Fertilizer with drop or rotary spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended

Disclaimer: Always refer to the label on the product before using Helena or any other product. 16June2010 revised storage & disposal

above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY **AND REMEDIES**

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular . purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

- 1. Refund of the purchase price paid by buyer or user for product bought, or
- Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2010. PRO-MATE® is a registered trademark of Helena Holding Company. DIMENSION® is a registered trademark of Dow AgroSciences.