

5905-550

1/31/2008

10918



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number: **5905-550**  
Date of Issuance: **1-31-08**

NOTICE OF PESTICIDE:  
\_ Registration  
x Reregistration  
(under FIFRA, as amended)

Term of Issuance:  
Name of Pesticide Product:  
**Wildcard XTRA**

Name and Address of Registrant (include ZIP Code):

**Helena Chemical Company**  
**225 Schilling Boulevard, Suite 300**  
**Collierville, TN 38017**

**Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.**

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

1. Include the statement "contains petroleum distillate" near the ingredients statement.
2. Per the acute toxicity review, the signal word must be changed to "CAUTION".
3. Per the acute toxicity review, it is recommended that the "If in Eyes" statement appear immediately after the "If Swallowed" statement.
4. Per the acute toxicity review, add "Note to Physician: May pose an aspiration pneumonia hazard. Contains petroleum distillate."
5. Per the acute toxicity review, add to the Hazards to Humans and Domestic Animals section "Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.
6. To the PPE change "A NIOSH ...or HE filter" to "NIOSH-approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-84A. It is recommended that you require the respirator wearer be fit tested, and trained in the use maintenance, and limitations of the respirator."

Signature of Approving Official:  
Erik Kraft  
Acting Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date:  
**1-31-08**

7. The language "This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants," is not needed per the MCPA RED.
8. On the third page of the label, the first sentence must read "This chemical has properties and **characteristics** associated..."
9. Per the Bromoxynil RED add this language to the PPE "Chemical-resistant gloves for cleaning, equipment, mixing and loading, Chemical-resistant apron for cleaning equipment, mixing and loading."
10. Per the Bromoxynil RED, for REI and early-entry PPE requirements add "The restricted-entry interval (REI) for cotton is 4 days and includes scouts and crop advisors. The exemption in the Worker Protection Standard for certified crop advisors does not apply to bromoxynil. Scouts and crop advisors are prohibited from entering the treated area during the entire 4-day REI for bromoxynil. Applicators and other users must inform crop advisors and scouts of this requirement."
11. To the label add a Non-agricultural use box, which includes the statement "Do not enter or allow others to enter until sprays have dried."
12. To the Warranty section add to the extent consistent with applicable law in front of "the company makes no other", "the exclusive remedy" and change "to the extent allowed by law" to "to the extent consistent with applicable law".
13. Revise the application restriction for flax, grasses grown for seed, and the wheat, barley, oats and rye group to include the maximum amount of product/acre/year. Also, the Pasture/Rangeland application restrictions are required for CRP areas.

**Wheat**

"Do not apply more than 0.75 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

**Barley**


"Do not apply more than 0.75 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

**Oats**

"Do not apply more than 0.75 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

**Rye**

"Do not apply more than 0.75 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."



Handwritten signatures and initials at the bottom of the page, including a large signature on the left, a circular mark in the middle, and the number '1-31-00' on the right.

3 of 18

**Flax**

"Do not apply more than 0.25 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

**Sod farms**

"Do not apply more than 1.5 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

**Golf Courses**

"Do not apply more than 1.5 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

**Grass Grown for Seed**

"Do not apply more than 1.5 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

**Pasture/ Rangeland**

"Do not apply more than 1.5 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."


**Noncropland Rights-of-Way (Broadcast Treatment)**

"Do not apply more than 1.5 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

**Noncropland Rights-of-Way (Spot Treatment)**

"Do not apply more than 3.0 lb of MCPA ae/acre per year (registrant state this in amount of formulation per acre per year)."

Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.



1.31.08

Erik Kraft  
Acting Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure

**SPECIMEN LABEL**



in EPA Letter Dated

1.31.08

40918

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

5905-550

**FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN SMALL GRAINS (WHEAT, BARLEY, OATS AND RYE), CONSERVATION RESERVE PROGRAM (CRP) AREAS, GRASSES GROWN FOR SEED PRODUCTION AND FLAX**

**ACTIVE INGREDIENT:**

Octanoic acid of ester of bromoxynil (3,5-dibromo-4-Hydroxybenzoxynil).....31.7%\*  
 Isooctyl ester of 2-methyl-chlorophenoxyacetic acid\*\*.....34.0%\*\*

**INERT INGREDIENTS:**.....34.3%  
 100.0%

\*Bromoxynil octanoate equivalent to 21.8% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

\*\* Equivalent to 21.8% 2-methyl-chlorophenoxyacetic acid or not less than 2.0 pounds MCPA acid per gallon.

EPA Reg. No. 5905-550

EPA Est. No. 228-IL-1

SN 1004/0305

Patent No. 6,232,272

**KEEP OUT OF REACH OF CHILDREN**

**WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.  
 For Medical Emergencies Only, Call 877-325-1840. Product of United Kingdom

**FIRST AID**

- |                                |   |
|--------------------------------|---|
| <b>IF SWALLOWED:</b>           | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul> |
| <b>IF ON SKIN OR CLOTHING:</b> | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |
| <b>IF IN EYES:</b>             | <ul style="list-style-type: none"> <li>• Hold eyelid open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>  |
| <b>IF INHALED:</b>             | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>   |

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

**MANUFACTURED FOR  
 HELENA CHEMICAL COMPANY  
 225 SCHILLING BOULEVARD, SUITE 300  
 COLLIERVILLE TENNESSEE 38017**

Disclaimer: Always refer to the label on the product before using Helena or any other product.

SPECIMEN LABEL

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

**PERSONAL PROTECTIVE EQUIPMENT:**

Some materials that are chemical-resistant to this product are as follows:

- Barrier laminate,
- Butyl rubber,
- Nitrile rubber, or
- Viton gloves

If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

**Personal Protective Equipment (PPE)**

Mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Chemical-resistant gloves when mixing, loading or using any hand-held equipment.

**ADDITIONAL PPE REQUIREMENTS** for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland. The mixers/loaders also must wear:

- a chemical-resistant apron, and
- a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N<sup>2</sup>, R, P, or HE filter

**See engineering controls for additional requirements.**

**USER SAFETY REQUIREMENTS:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS FOR AERIAL APPLICATIONS:**

**Enclosed Cockpits**

**Engineering Controls:** Pilots must use and enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d) (6)]

**User Safety Recommendations:**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards:**

- This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**SPECIMEN LABEL**

- **This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.**

**ENVIRONMENTAL HAZARDS STATEMENTS:**

**Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination System ( NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.**

**APPLICATION BY CHEMIGATION** must be done by fixed pipe, overhead sprinkler systems or hand moved pipe. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

**DURING AERIAL APPLICATIONS**, human flaggers are prohibited unless in enclosed vehicles. Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, hospitals, shopping areas, etc.)

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not cut or weld container.

**WARNING**

**WILDCARD® XTRA** Herbicide contains low volatile isooctyl ester of MCPA. At high air or ground surface temperatures, vapors from this product may cause injury to susceptible plants. This fact should be considered when applying **WILDCARD® XTRA**.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR parts 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Disclaimer: Always refer to the label on the product before using Helena or any other product.

**SPECIMEN LABEL**

- ❖ Coveralls
- ❖ Shoes plus socks,
- ❖ Chemical-resistant gloves made of any waterproof material, and

**STORAGE AND DISPOSAL**

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Store at temperatures above 3° F. If allowed to freeze, remix before using.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**RETURNABLE – REFILLABLE CONTAINERS:** This material may be repackaged in 15 or 30 gallon returnable-refillable containers by Helena or a registered establishment under contract to Helena. After use, return the container to the point of purchase or designated locations. This container must only be refilled with **WILDCARD® XTRA**. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. If the container is not being refilled, return it to the point of purchase.

**GENERAL INFORMATION**

**WILDCARD® XTRA** is formulated as an emulsifiable concentrate containing the equivalent of 2 lbs. per gallon of bromoxynil and 2 pounds per gallon of MCPA.

**WILDCARD® XTRA** is a selective postemergence herbicide for control of important broadleaf weeds infesting small grains (wheat, barley, oats, rye), conservation reserve program areas, and grass grown for seed. Optimum weed control is obtained when **WILDCARD® XTRA** is applied to actively growing weed seedlings. **WILDCARD® XTRA** is primarily a contact herbicide; therefore thorough coverage of the weed seedlings is essential for optimum control.

**WILDCARD® XTRA** has little residual activity. Therefore subsequent flushes of weeds will not be controlled by the initial treatment.

Generally crops that form a good canopy will help shade subsequent weed flushes. However, certain crops or short-straw varieties, for example Yaccora Rojo wheat, may not develop the crop canopy fast enough to shade the subsequent flushes of weeds.

Occasional transitory leaf burn may occur. The temporary leaf burn is similar to that seen with liquid fertilizer. Because the activity of **WILDCARD® XTRA** is mainly contact, recovery of the crop is generally rapid with no lasting effect. Frequency and amount of leaf burn may be greater when crops are stressed by abrasive winds, cool to cold evening temperatures or mechanical injury, such as that caused by hail, sleet or insect feeding. To reduce the potential for temporary leaf burn, applications should be made to dry foliage in the recommended spray volumes per acre when weather conditions are not extreme.

**MIXING, LOADING AND HANDLING INSTRUCTIONS**

**2.5 Gallon Containers**

It is strongly recommended that special care be taken in mixing and loading this product. Hands should be placed on the container in such a way as to avoid possible drip or splash.

Disclaimer: Always refer to the label on the product before using Helena or any other product.

**SPECIMEN LABEL**

**30 Gallon and Bulk Containers**

If you will handle a total of 60 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

**WILDCARD® XTRA ALONE:** Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of **WILDCARD® XTRA**. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

**TANK MIXTURES: WILDCARD® XTRA** can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply **WILDCARD® XTRA** in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water add the recommended amount of **WILDCARD® XTRA** and add water to the spray tank to the desired level. If tank mixing with other product types, add the **WILDCARD® XTRA** first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application. If tank mixing with products other than those listed within each crop section, a compatibility test is recommended to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture with **WILDCARD® XTRA**.

**SPRAYABLE LIQUID FERTILIZERS AND SPRAY ADDITIVES**

**WILDCARD® XTRA** can be applied in combination with sprayable liquid fertilizer or spray additives such as surfactants or crop oil concentrate. When tank mixing with liquid fertilizer always add the fertilizer to the spray tank first and agitate thoroughly before adding **WILDCARD® XTRA**. Always predetermine the compatibility with liquid fertilizer by mixing small proportional quantities in advance. Agitation must be maintained during filling and application operations to ensure that **WILDCARD® XTRA** is evenly mixed with the fertilizer. Leaf burn may occur when **WILDCARD® XTRA** is applied with liquid fertilizer, but new leaves are not adversely affected.

**PRECAUTION:** Fertilizers and spray additives can increase foliage leaf burn when applied with **WILDCARD® XTRA**. Do not apply fertilizers or spray additives with **WILDCARD® XTRA** if leaf burn is a major concern due to environmental conditions, crop or variety sensitivity to **WILDCARD® XTRA**.

**APPLICATION PROCEDURES**

**WILDCARD® XTRA** can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment.

Disclaimer: Always refer to the label on the product before using Helena or any other product.



**SPECIMEN LABEL**

**GROUND APPLICATION**

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage use of flat fan nozzles (maximum tip size 8008) with a spray pressure of 40-60 psi are recommended. Other nozzle types and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. Raindrop® nozzles and flood nozzles are not recommended as weed control with **WILDCARD® XTRA** may be reduced.

In general, a spray volume of 10 to 20 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi and a maximum ground speed of 10 mph may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application which results in poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas. Applications using less than 10 gallons per acre may result in reduced weed control.

When weed infestations are heavy, use of higher spray volumes and spray pressure will be helpful in obtaining uniform weed coverage. When corn or grain sorghum are large enough to interfere with the spray pattern, drop nozzles should be used to obtain uniform weed coverage. If you are unsure of the infestation level or size of crop, consult your local extension service.

Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement.

**Additional requirements for ground boom application:**

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

**AERIAL APPLICATION**

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended. A minimum spray volume of 3 gallons per acre may be used if crop canopy and weed density allow adequate spray coverage. Aerial applications using less than 5 gallons of spray volume per acre may result in reduced weed control.

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off target spray movement. Off target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.

**SPRAY DRIFT MANAGEMENT:**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray ( ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Disclaimer: Always refer to the label on the product before using Helena or any other product.

## SPECIMEN LABEL

**ADDITIONAL REQUIREMENTS FOR AERIAL REQUIREMENTS:**

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

**SPRINKLER IRRIGATION APPLICATION**

**WILDCARD® XTRA** Herbicide can be applied through sprinkler irrigation systems to small grains and grasses grown for seed.

Apply **WILDCARD® XTRA** Herbicide through sprinkler systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle. Do not apply this product through any other type of irrigation system.

**SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH AUTOMATED SPRINKLER IRRIGATION SYSTEM**

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Agitation is recommended in the pesticide supply tank when applying the **WILDCARD® XTRA** Herbicide.
9. **WILDCARD® XTRA** Herbicide should be applied continuously for the duration of the water application with center pivot and continuous lateral move systems. Application of **WILDCARD® XTRA** Herbicide should be made during the last 30-45 minutes of the irrigation set with other overhead sprinkler systems.
10. For best performance, set the sprinkler system to deliver approximately 0.5 inch or less of water per acre.
11. Remove scale, pesticide residues and other foreign matter from the supply tank and entire injector system. Flush with clean water.
12. If **WILDCARD® XTRA** Herbicide is diluted in the supply tank, fill the tank with half of the water amount desired, add the **WILDCARD® XTRA** and then add remaining water amount with agitation. Always dilute with at least 4 parts water to 1 part **WILDCARD® XTRA**.
13. Start the sprinklers and then inject **WILDCARD® XTRA** Herbicide into the irrigation line. **WILDCARD® XTRA** should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to insure adequate mixing. Refer to the **WILDCARD® XTRA** Herbicide label for detailed information on application rates and timings.

**SPECIMEN LABEL**

**CHEMIGATION USER PRECAUTIONS**

Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils. Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.

Allow sufficient time for pesticide to be flushed through all the lines and nozzles before turning off irrigation water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Do not connect an irrigation system used for pesticide application to a public water system. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**GENERAL WEED LIST**

Postemergence application of **WILDCARD® XTRA** Herbicide will control the following weeds when sprayed in the seedling stage.

Maximum weed stage of growth is listed under **WILDCARD® XTRA** RECOMMENDATIONS.

**MOST SUSCEPTIBLE BROADLEAF WEED SPECIES**

- Annual sowthistle (*Sonchus oleraceus*)
- Black mustard (*Brassica nigra*)
- Black nightshade (*Solanum nigrum*)
- Common cocklebur (*Xanthium strumarium*)
- Common lambsquarters (*Chenopodium album*)
- Common tarweed (*Hemizonia congesta*)
- Cow cockle (*Saponaria vaccaria*)
- Cutleaf nightshade (*Solanum triflorum*)
- Eastern black nightshade (*Solanum ptycanthum*)
- Coast fiddleneck (*Amsinckia intermedia*)
- Field pennycress (*Thlaspi arvense*)
- Green smartweed (*Polygonum scabrum*)
- Hairy nightshade (*Solanum sarachoides*)
- Horned Poppy (*Glaucium corniculatum*)
- Jimsonweed (*Datura stramonium*)
- Ladysthumb (*Polygonum persicaria*)
- Lanceleaf sage (*Salvia reflexa*)
- London rocket (*Sisymbrium irio*)
- Marshelder (*Iva xanthifolia*)
- Pennsylvania smartweed (*Polygonum strumarium*)
- Pepperweed spp. (*Lepidium app.*)
- Redroot pigweed (*Amaranthus retroflexus*)
- Russian thistle (*Salsola kali*)

**SUSCEPTIBLE BROADLEAF WEED SPECIES**

- Blue (purple) mustard (*Chlorispora tenella*)
- Common groundsel (*Senecio vulgaris*)
- Common ragweed (*Ambrosia artemisiifolia*)
- Corn chamomile (*Anthemis arvensis*)
- Corn gromwell (*Lithospermum arvense*)
- Fumitory (*Fumaria officinalis*)
- Giant ragweed (*Ambrosia trifida*)
- Hemp sesbania (*Sesbania exaltata*)
- Henbit (*Lamium amplexicaule*)
- Ivyleaf morningglory (*Ipomoea hederacea*)
- Knawel (*Scleranthus annuus*)
- Kochia (*Kochia scoparia*)
- Mayweed (*Anthemis cotula*)
- Prostrate knotweed (*Polygonum aviculare*)
- Puncture vine (*Tribulus terrestris*)
- Tall morningglory (*Ipomoea purpurea*)
- Tansy mustard (*Descurainia pinnata*)
- Tarweed (*Hemizonia spp.*)
- Velvetleaf (*Abutilon theophrasti*)
- Wild radish (*Raphanus raphanistrum*)

Weeds germinating after spraying will not be controlled.

12/09/18

**SPECIMEN LABEL**

- Shepherdspurse (*Capsella bursa-pastoris*)
- Silverleaf nightshade (*Solanum elaeagnifolium*)
- Smooth pigweed (*Amaranthus hybridus*)
- Spiny pigweed (*Amaranthus spinosus*)
- 1 Sunflower (*Helianthus annuus*)
- Tall Waterhemp (*Amaranthus tuberculatus*)
- Tartary buckwheat (*Fagopyrum tataricum*)
- Tumble mustard (*Sisymbrium altissimum*)
- Wild buckwheat (*Polygonum convolvulus*)
- Wild mustard (*Sinapis arvensis*)
- Yellow rocket (*Barbarea vulgaris*)

<sup>1</sup> For control of sunflower, delay application until first sunflower seedlings emerging are 4 inches in height.

**WEED SUPPRESSION**

Canada Thistle (*Cirsium arvense*)

**WILDCARD® XTRA** Herbicide applied at 1-1/2 pints per acre provides burn down of top growth. Regrowth may occur. Make applications when Canada thistle is 8 inches tall to the bud stage.

**WHEAT, BARLEY, OATS AND RYE  
WILDCARD® XTRA RECOMMENDATIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA	1 pint/A	Fall seeded wheat, barley, oats and rye throughout the United States and spring seeded wheat, barley, oats and rye in Idaho, Oregon, Washington, Colorado, Wyoming and Montana. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	<b>MOST SUSCEPTIBLE BROADLEAF WEEDS:</b> Apply to weeds up to the 8 leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter.
	1-1/2 to 2 pints/A		<b>SUSCEPTIBLE BROADLEAF WEEDS:</b> Apply to weeds up to the 4 leaf stage or 2 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.
	2 pints/A		Apply to henbit, knawel and mayweed up to the 4 leaf stage or 2 inches in height, whichever comes first. Apply to kochia and tansy mustard for improved control when these weeds exceed the recommended stage of growth or are growing under cool, dry conditions.
	1-1/2 pints/A	Spring seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	<b>MOST SUSCEPTIBLE AND SUSCEPTIBLE BROADLEAF WEEDS:</b> Apply to weeds that do not exceed the 8 leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter. Apply to kochia up to 2 inches in height.

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.

**SPECIMEN LABEL**

1-1/2 to 2 pints/A	Spring seeded wheat and barley except except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3 leaf stage but before the crop reaches the boot stage.	Apply to kochia that is 2-4 inches height.
Chemigation Only 2 pints/A	Apply to wheat, barley, oats and rye from the 3 leaf stage but before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING LOADING AND HANDLING INSTRUCTIONS section for complete details.	Apply to MOST SUSCEPTIBLE and SUSCEPTIBLE broadleaf weeds up to the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.
Post-harvest 3/4 - 2 pints/A	Make applications following harvest of wheat, barley, oats and rye in the states of North Dakota, South Dakota, Minnesota, and Montana. Do not plant any rotational crop until	Apply 3/4 to 1 pints/A to MOST SUSCEPTIBLE BROADLEAF WEEDS up to the 8 leaf stage or 4 inches in height, whichever comes first. Apply 1-1/2 to 2 pints/A to SUSCEPTIBLE BROADLEAF

**WHEAT, BARLEY, OATS AND RYE  
WILDCARD® XTRA RECOMMENDATIONS (continued)**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
		the following use season.	WEEDS up to the 4 leaf stage or 2 inches in height, whichever comes first. For control of both grasses and broadleaf weeds, tank mix WILDCARD® XTRA with Roundup® or Roundup®+2,4-D such as WEEDONE® or WEEDAR® brand herbicides.

**WHEAT, BARLEY, OATS AND RYE  
WILDCARD® XTRA TANK MIXTURE RECOMMENDATIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA + Rhonox® (MCPA Ester)	3/4 - 2 pints/A + 1/4-1/2 pint/A	Apply to spring seeded wheat, barley, oats and rye from tillering stage, but before boot stage.	For control of MOST SUSCEPTIBLE and SUSCEPTIBLE weeds and improved control of redroot pigweed and kochia. Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia and redroot pigweed up to 2 inches in height or diameter.
WILDCARD® XTRA + Glean® +	3/4 to 1-1/2 pints/A + 1/6-1/3 oz/A +	Apply to wheat and barley from the 3 leaf stage but before the crop reaches the boot stage. Refer to Glean® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.

**SPECIMEN LABEL**

nonionic surfactant	1 qt/100 gal of water		first.
<b>WILDCARD® XTRA</b> + Finesse® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6-1/3 oz/A + 1 qt/100 gal of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Finesse® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
<b>WILDCARD® XTRA</b> + Ally® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/10 oz/A + 1 qt/100 gal of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Ally® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
<b>WILDCARD® XTRA</b> + Banvel®	3/4 to 1-1/2 pints/A + 1/8 - 1/4 pint/A	Fall seeded wheat from the 3 leaf stage but before jointing. Spring seeded wheat from the 3 to 5 leaf stage of growth.	This tank mix improves control of broadleaves such as prostrate knotweed and kochia. Apply to weeds up to the 8 leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to kochia up to 2 inches in height or diameter.

**WHEAT, BARLEY, OATS AND RYE**

**WILDCARD® XTRA TANK MIXTURE RECOMMENDATIONS (continued)**

<b>APPLICATION TIMING AND SPECIFIC COMMENTS</b>			
<b>PRODUCT</b>	<b>RATE</b>	<b>CROP</b>	<b>WEEDS</b>
<b>WILDCARD® XTRA</b> + Harmony® Extra + nonionic surfactant	3/4 to 1-1/2 pints/A + 3/10 - 1/2 oz/A + 1 qt/100 gal of water	Winter wheat. Apply from the 3 leaf stage but before the 3 <sup>rd</sup> node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions. Spring wheat and barley. Apply after the 3 leaf but before the 1 <sup>st</sup> node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed and redroot pigweed. Apply to weeds up to the 8 leaf stage, 4 inches in height or across, whichever comes first.
<b>WILDCARD® XTRA</b> + Amber® + nonionic surfactant	3/4 to 1-1/2 pints/A + 0.28 - 0.56 oz/A + 0.25 - 0.5% v/v	Apply to wheat and barley from the 3 leaf stage, but before the flag leaf is visible. Refer to the Amber® label for crop rotation and other restrictions.	This tank mix improves control of broadleaves such as henbit, tansy mustard, and pigweed. Apply to weeds up to the 4 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
<b>WILDCARD® XTRA</b> + Express® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 - 1/3 oz/A + 1 qt/100 gal of water	Wheat and barley. Apply from the 3 leaf stage but before the flag is visible. Refer to the Express® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed, redroot pigweed and suppression of Canada thistle. Apply to annual weeds up to the 8 leaf stage, 4 inches in height or across, whichever comes first and to Canada thistle 4 to 8 inches tall with 2 to 6 inches of new growth.
<b>WILDCARD® XTRA</b> +	3/4 to 1-1/2 pints/A +	Apply to wheat and barley after the crop begins to tiller up to the 1 <sup>st</sup> node detectable.	This tank mix improves control of kochia, wild buckwheat and suppression of Canada thistle. Apply to annual broadleaf weeds up

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.

**SPECIMEN LABEL**

Curtail® or Curtail® M	2 pints/A		to the 8 leaf stage, 4 inches in height or 2 inches in diameter and to Canada thistle in the rosette to prebud stage.
WILDCARD® XTRA + metribuzin  (Sencor® or Lexone®)	1 pint/A + 1/8 - 3/16 lb ai/A	Winter wheat in Idaho, Oregon and Washington. Apply in spring after growth has started and secondary roots with a minimum of 3 to 4 tillers have been established, but before the forming of joints in the stem. Avoid application when crop has experienced winter kill, frost damage, disease or drought.	This tank mix improves control of broadleaf weeds such as chickweed, filaree, henbit. Apply to weeds up to the 4 leaf stage, 2 inches in height or diameter, whichever comes first. A recognized authority should be consulted concerning the use of this mixture in your area.
WILDCARD® XTRA + Avenge®	1 - 2 pints/A + 2-1/2 to 4 pints/A	Winter wheat. Four leaf to tillering stage. Refer to Avenge® label for varietal and other restrictions. Spring Wheat. Five to 6 leaf stage. Refer to Avenge® label for varietal and other restrictions. Barley. Three to 7 leaf stage.	This tank mix will provide wild oat control in addition to broadleaves. Apply to wild oats in the 3-5 leaf stage and broadleaves that do not exceed the 4 leaf stage or rosettes of 1.5 inches in diameter. Avenge use rates per acre are 2-1/2 pints (1-10 oats per sq. ft.), 3 pints (11-25 oats per sq. ft.) or 4 pints (more than 25 oats per sq. ft.).
WILDCARD® XTRA + Assert®	1 to 1-1/2 pints/A + 1 to 1-1/2 pints/A	Apply to wheat and barley from the 3 leaf stage but before boot stage. Refer to Assert® label for crop rotation and other restrictions.	This tank mix will provide wild oat control in addition to broadleaf weeds. Apply to wild oats at the 1-4 leaf stage and broadleaf weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first. Use Assert at 1-1/2 pints/A west of the Rocky Mountains or if wild oats have initiated tillering. For spray volumes in excess of 10 GPA, add 0.3 fluid oz of nonionic surfactant for each gallon in excess of 10 GPA.

**Restrictions and Precautions: Wheat, Barley, Oats and Rye**

- Do not graze treated field within 45 days after application.
- Do not apply when crops are under moisture stress.
- Do not apply when crop canopy covers the weeds as poor control will result.
- Reduced weed control may occur when weeds are stressed from lack of moisture or cold temperatures.
- Refer to labels of products used in tank mixture for additional restrictions and precautions.
- **Do not apply more than 0.75 lb ae / acre per year.**

**CONSERVATION RESERVE PROGRAM AREAS (CRP)  
WILDCARD® XTRA RECOMMENDATIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA	1 - 2 pints/A	Apply to grasses from the 3 leaf stage.	Apply 1 pint/A to MOST SUSCEPTIBLE and 1-1/2 to 2 pints/A to SUSCEPTIBLE broadleaf weeds up to the 8 leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.

**RESTRICTIONS AND PRECAUTIONS: CRP AREAS**

- Do not allow livestock to graze in treated areas or feed treated grass to livestock.

**SPECIMEN LABEL**

- If legumes are included in CRP area planting, severe injury may occur to legumes treated with **WILDCARD® XTRA**.

**GRASSES GROWN FOR SEED PRODUCTION  
WILDCARD® XTRA Recommendations  
Seedling and Established Grasses**

PRODUCT	RATE	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
	Per ACRE	Per 1000 SQ.FT.	CROP	WEEDS
<b>WILDCARD® XTRA</b>	1 to 3 Pints	0.375 to 1.125 Fl. Oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Established grasses tolerant to <b>WILDCARD® XTRA</b> include bentgrasses, Kentucky Bluegrass, Fescues, Ryegrass, Bermudagrass, St. Augustinegrass and Zoysiagrass. <b>WILDCARD® XTRA</b> may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky Bluegrasses, Pennlawn, Chewings, Illahee or Alta Fescues, Orchard grass, Highland, Seaside or Astoria Bentgrasses, perennial Ryegrasses, Bahiagrass and Zoysiagrass.	Refer to the GENERAL WEED LIST for a listing of susceptible broadleaf weeds. Optimal control will be attained when weeds are treated in the seedling stage (less than 4 leaf stage, 2 inches in height, or 1 inch in diameter).
<b>WILDCARD® XTRA</b>	Chemigation 2 pints/A only	0.75 Fl. Oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Apply through	

**GRASSES GROWN FOR SEED PRODUCTION  
WILDCARD® XTRA Recommendations (continued)  
Seedling and Established Grasses**

PRODUCT	RATE	RATE	APPLICATION TIMING AND SPECIFIC COMMENTS	
	Per ACRE	Per 1000 SQ.FT.	CROP	WEEDS
	1-3pints	0.375 -1.125 Fl Oz.	automated sprinkler irrigation systems with mechanical transfer loading system only. See MIXING, LOADING AND HANDLING INSTRUCTIONS section for complete details. Refer to the list of established grasses that are tolerant to <b>WILDCARD® XTRA</b> .	

**RESTRICTIONS AND PRECAUTIONS:** Grasses grown for seed or sod production

- Do not allow livestock to graze in treated areas or feed treated grasses to livestock.
- Do not apply **WILDCARD® XTRA** to grasses grown for seed production with backpack or hand-held application equipment.

Disclaimer: Always refer to the label on the product before using Helena or any other product.



17 of 18

**SPECIMEN LABEL**

**Note: Do not apply more than 1.5 Lb ae/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.**

**FLAX (*Linum usitatissimum* only)  
WILDCARD® XTRA Recommendations**

<b>APPLICATION TIMING AND SPECIFIC COMMENTS</b>			
<b>PRODUCT</b>	<b>RATE</b>	<b>CROP</b>	<b>WEEDS</b>
<b>WILDCARD® XTRA</b>	0.9 pint- 1 pint/ac	Apply to flax 2 to 8 inches in height. Do not apply <b>WILDCARD® XTRA</b> to flax during or after the bud stage.	Apply to <b>MOST SUSCEPTIBLE</b> weeds that do not exceed the 4 leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.

**RESTRICTIONS AND PRECAUTIONS: Flax (*Linum usitatissimum* only)**

- Do not apply if temperatures are expected to exceed 85° F at or 3 days following application or crop injury may occur.
- Unacceptable crop injury may occur following **WILDCARD® XTRA** application to flax grown on high organic, peat type soils.
- Application under high humidity conditions can injure flax.
- Unless otherwise instructed, do not apply **WILDCARD® XTRA** to flax with crop oil concentrate, surfactants or nitrogen solutions.
- Do not use on ornamental flax.

**Note: Do not apply more than 0.25 lb ae/ acre year.**

**CONDITIONS OF SALE - LIMITED WARRANTY  
AND LIMITATIONS OF LIABILITY AND REMEDIES**

**Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.**

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

**Disclaimer:** Always refer to the label on the product before using Helena or any other product.

180 of 18

**SPECIMEN LABEL**

© Copyright Helena Holding Company, 2005.

**ASSERT®**, **BANVEL®**, and **AVENGE®** are registered trademarks of BASF Corp.

**SENCOR®** is a registered trademark of Bayer AG, Germany.

**RAINDROP®** is a registered trademark of Delavan Corporation.

**CURTAIL®** is a registered trademark of Dow AgroSciences.

**ALLY®**, **EXPRESS®**, **FINESSE®**, **GLEAN®**, **HARMONY®** and **LEXONE®** are registered trademarks of E.I. DuPont DeNemours and Co.

**AMBER®** is a registered trademark of Syngenta Crop Protection.

**RHONOX®**, **WEEDAR®** and **WEEDONE®** are registered trademarks of NuFarm, Inc.

**ROUNDUP®** is a registered trademark of Monsanto Company.

**WILDCARD®** is a registered trademark of Helena Holding Company.