

5905-574

2-27-2008

1/23



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

5905-574

Date of Issuance:

FEB 27 2008

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

OUTLAW

Name and Address of Registrant (include ZIP Code):

Mr. Scott A. Pace
Helena Chemical CO.
225 Schilling Boulevard, Suite 300
Collierville, TN 38017

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following changes to the label:

a. Change the product registration number to "EPA Reg. No. 5905-574".

(continued on page 2)

Signature of Approving Official:

Handwritten signature of Joanne I. Miller, Product Manager (23), Herbicide Branch, Registration Division (7505P)

Date:

FEB 27 2008

EPA Form 8570-6

b. Add the correct EPA Establishment Number.

c. Under "PRECAUTIONARY STATEMENTS" on page 1, make "CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE" all capital letters.

d. On page 2, under "PERSONEL PROTECTIVE EQUIPMENT", change "Applicators and other handlers must wear:" to "All mixers, loaders, flaggers, applicators and other handlers must wear:". After "Chemical-resistant apron when cleaning", add "up spills or equipment, mixing, loading, or otherwise exposed to the concentrate." Under the bullet "Chemical-resistant apron, etc.", skip a line and add "See Engineering Controls for additional requirements and exceptions." In the washing instructins for Personel Protection Equipment, add "exist" after "...washables". Change "Engineering Control Statements" to "Engineering Controls", and replace the statement below "Engineering Controls" with "Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(6)]." Under "ENVIRONMENTAL HAZARDS", change the first sentence to "This product is toxic to fish and aquatic invertebrates." Replace the second sentence with "Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated area." After the first paragraph under "ENVIRONMENTAL HAZARDS", add as a separate paragraph, "The chemicals in this product have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater."

e. On page 3 and before the "AGRICULTURAL USE REQUIREMENTS" box, add "General Application Restrictions", and the sentences "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handles may be in the area during application." In the "AGRICULTURAL USE REQUIREMENTS" box, replace the REI of "48 hours" with "24 hours"; after "Coveralls" add "worn over short-sleeve shirt and short pants"; change "Shoes plus socks" to "Chemical resistant shoes plus socks"; under "Protective Eyewear", add the bullet "Chemical resistant headgear for overhead exposure"; skip a line and add the sentence "Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area." In the "NON-AGRICULTURAL USE REQUIREMENTS" box remove the entire second paragraph, and replace with "Do not allow people or pets in treated area during application. Do not allow people or pets to enter treated area until spray has dried." On a separate line in the "NON-AGRICULTURAL USE REQUIREMENTS" box, add "CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE. Applicator must wear protective eyewear." Remove the last paragraph starting with "TURF USE REQUIREMENTS:". There are no use directions for turf/sod production on product label. In the "STORAGE AND DISPOSAL" box, rewrite the "CONTAINER DISPOSAL" directions to conform to Pesticide Registration (PR) Notice 2007-4 (Labeling Revision Required by the Final Rule "Pesticide Management and Disposal;

f. Replace the three paragraphs under "SPRAY DRIFT MANAGEMENT" with language required for end-use products in the 2,4-D RED, which is:

"A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Apply as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

Temperature Inversion

If applying at wind speeds less than 3 mph, the applicator must determine if:

- a) conditions of temperature inversion exist, or
- b) stable atmospheric conditions exist at or below nozzle height.

Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged, or crops thereof rendered unfit for sale, use, or consumption. Susceptible crop include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding applications of 2,4-D herbicides. Where States have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional Requirements for Aerial Applicators

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way application.

4/
23

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional Requirements for Ground Boom Applications

Do not apply with a nozzle height greater than 4 feet above the crop canopy. This product contains a 2,4-D ester. 2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures."

g. On pages 5 and 6, remove the "AERIAL DRIFT REDUCTION ADVISORY". The Advisory is confusing and unnecessary with the new mandatory spray Drift Management language above.

h. On page 11, under "Restrictions and Limitations", remove the redundant and uncorrected restriction "Restricted entry interval (REI): 48 Hours". Change the sentence "Planting/replanting restrictions for applications of more than 6 pints and up to 8 pints of OUTLAW™ per acre." to "Planting/replanting restrictions for applications of more than 6 pints and up to 8 pints of OUTLAW™ per acre/season:".

i. On page 12, under "Food/Feed Crop Specific Information", remove the sentence "Rates above 4 pints of OUTLAW™ per acre are for spot treatments only." The sentence appears to contradict the 6 pints/A for between crop applications. Remove the same sentence on page 14. After the sentence starting with "Retreatments may be made as needed; etc.", add "There is a minimum of 30 days between applications."

j. On page 13, under "Grazing and Feeding Non-Lactating Animals:": remove the sentence "Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter." This restriction is unnecessary, because the appropriate meat tolerances are established (memo of T. Bloem, HED, 4/20/2005, DP# 313211). Under "Wheat", remove "Non-Food/Feed Use (Land not Harvested, Grazed or Foraged)-Specific Information", as this language is misleading. Instead move "Between Crop Applications" into its place. Under "Rates and Timings", change "...do not exceed a total of 8 pints of OUTLAW™ per treated, etc." to "...do not exceed a total of 8 pints of OUTLAW™ (3.33 pints for wheat) per, etc."

k. Page 14, after the list of tank partners, add the restriction "When tank mixing with products that contain either 2,4-D or dicamba, do not exceed the lowest annual application rate/A for each active ingredient for that crop/use." Under "Farmstead and Fence-row Treatment Application Instructions", fourth paragraph starting with "Do not exceed 40 gallons, etc.", add "(not to exceed 8 pints of OUTLAW™/A)", after "... of spray solution, etc." Remove all references/sentences pertaining to the use of diesel oil. (e.g., 10% diesel oil, etc, "The diesel oil in this tank, etc.", "4). Diesel Oil;

etc.", and "1. Increase diesel oil content to 15%, etc."). Diesel oil is not an approved use with pesticides in the environment. 5/23


l. Page 17, sentence above "CONDITIONS OF SALE AND WARRANTY" starting with the bullet "These crops are considered Food/Feed crops only, etc." This statement is overly broad, potentially misleading, and does not take into consideration release of land by Federal Authorities for grazing/harvest in emergence situations, like this year in most of the Continental U.S.

m. Page 18, under "CONDITIONS OF SALE AND WARRANTY", add to the beginning of the third paragraph starting with "BUYER'S EXCLUSIVE REMEDY, etc." the phrase "TO THE EXTENT ALLOWED BY APPLICABLE LAW,". Add the same phrase to the beginning of the sentences starting with "When Buyer suffers losses, etc." and "IN NO CASE SHALL HELENA OR THE SELLER, etc."

3. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.


for Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure: Label stamped "Accepted with Comments"



OUTLAW™

ACCEPTED
with COMMENTS
In EPA Letter Dated

6/23

FEB 27 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act
General Pesticide
registered under EPA Reg. No. 5905-574

For control of a wide-spectrum of annual, biennial, and perennial broadleaf weeds and brush in Pastures, Rangeland, and Grass (Hay, Silage); Wheat; Conservation Reserve Program lands; General Farmland Areas; Post-Harvest, Fallow, Crop Stubble and Set Aside Acres

ACTIVE INGREDIENT(S):

| | |
|---|--------|
| 3,6-dichloromethoxybenzoic acid..... | 12.18% |
| 2-Ethylhexyl Ester of 2,4-Dichlorophenoxyacetic Acid..... | 24.28% |

INERT INGREDIENTS:

| | |
|-------------|---------|
| TOTAL | 100.00% |
|-------------|---------|

Equivalent to:

12.18% Dicamba Acid, 1.09 lbs./gal

16.10% 2,4-D Acid or 1.45 lbs./gal

Isomer specific by AOAC Method 6.D01-5 (12th Ed.)

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it you in detail.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious or convulsing person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONS AND DIRECTIONS FOR USE

EPA REG. NO. 5905-
EPA EST. NO. 42750-

NET CONTENTS:

Helena Chemical Company
225 Schilling Blvd., Suite 300
Collierville, TN 38017

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

7/23

Applicators and other handlers must wear:

- Long-Sleeved shirt and long pants
- Chemical resistant gloves Category C, such as butyl rubber \geq 14 mils, or nitrile rubber \geq 14 mils, or neoprene rubber \geq 14 mils or viton \geq 14 mils
- Shoes plus socks
- Protective Eyewear
- Chemical-resistant apron when cleaning equipment, mixing or loading

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be re-used until it has been cleaned.

If this container contains over 1 gallon and less than 5 gallons, mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

If this container contains 5 gallons or more in capacity, do not open pour. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Engineering Control Statements

When handlers use enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove PPE immediately after handling this product.
- Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from target area. Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides. Do not contaminate water by cleaning of equipment or disposing of equipment washwaters or rinsate.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Endangered Species Concerns:

The use of any pesticide in a manner that may kill or otherwise harm and endangered species or adversely modify their habitat is a violation of federal law.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

8/23

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **48 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves Category C, such as butyl rubber \geq 14 mils, or nitrile rubber \geq 14 mils, or neoprene rubber \geq 14 mils or viton \geq 14 mils
- Shoes plus socks
- Protective Eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

USE REQUIREMENTS FOR PASTURES, PERENNIAL GRASSLANDS, RANGELAND, FALLOW LAND AND NONCROP AREAS: Do not enter treated areas until spray has dried. For early entry to treated areas, wear eye protection, chemical-resistant gloves Category C, such as butyl rubber \geq 14 mils, or nitrile rubber \geq 14 mils, or neoprene rubber \geq 14 mils or viton \geq 14 mils, long-sleeved shirt, long pants, shoes and socks.

TURF USE REQUIREMENTS: Do not allow persons (other than applicator) or pets on treated area during application. Do not enter treated areas until spray has dried. NOTE: For application to turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes, follow **AGRICULTURAL USE REQUIREMENTS** on this label.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions that might adversely affect the container or its ability to function properly.

Pesticide Storage: Do not store below temperature of 32°F or above 100°F. Store in original container in a well-ventilated area separately from fertilizer, feed, and foodstuffs. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength.

Pesticide Disposal: Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Plastic/Metal Containers: Triple rinse (or equivalent) and add rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable-Refillable Container (Drum/Bulk/Mini-bulk):

When this container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase or to a designated location named at the time of purchase of this product in a bulk container. This container may only be refilled with this herbicide. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact ChemTrec at 1-800-424-9300 or Helena Chemical Co. at 901-761-0050. If not returned to the point of purchase or to the designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

9/23

Steps to be taken in case material is released or spilled:

Dike and contain the spill with inert material (sand, earth, etc) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

I. GENERAL INFORMATION

OUTLAW™ herbicide is a postemergence herbicide for controlling a wide spectrum of annual, biennial, and perennial broadleaf weeds and brush in pastures, rangeland, and grass (hay, silage); wheat; conservation reserve program land; postharvest, fallow, crop stubble, set-aside acres; and general farmstead areas.

Mode of Action

OUTLAW™ contains two active ingredients uniquely formulated to be used alone or tank mixed with other listed products as well as liquid fertilizer solutions. **OUTLAW™** is readily absorbed by plants through shoot and root uptake, translocates throughout the plant's system, and accumulates in areas of active growth. **OUTLAW™** interferes with the plant's growth hormones (auxins) resulting in death of many broadleaf weeds.

For best results, thoroughly clean sprayer equipment (tank, lines and nozzles) immediately after use by flushing system with water and heavy duty detergent or other suitable tank cleaner.

II. APPLICATION INSTRUCTIONS

Apply **OUTLAW™** at the rates and growth stages listed in Tables 1 and 2 as follows unless instructed differently by section on "Food/Feed Crop Specific Information" or "Non-Food/Feed Use (Land not Harvested, Grazed or Foraged)-Specific Information." **OUTLAW™** may be applied using water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence use for all crops listed on this label. Postemergence uses with sprayable fluid fertilizer may be made on pasture, hayland, or wheat crops only. The most effective application rate and timing varies based on the target weed species (refer to Table I). In mixed populations of weeds the correct rate is determined by the weed species requiring the highest rate. Delaying application permits weeds to exceed the maximum size and will prevent adequate control. For certain specified applications liquid fertilizer or oil may replace part or all of the water as diluent. If dry flowable (DF), wettable powder (WP) or flowable (F) tank mix products are to be used, these should generally be added to the spray tank first. Refer to the mixing directions on the labels of the tank mix products.

Irrigation:

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Chemigation Prohibition

Do not apply this product through any type of irrigation system.

Spray Coverage:

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Sensitive Crop Precautions:

OUTLAW™ may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems or foliage. At high temperatures (about 85 degrees or higher), vapors from this product may cause injury to the aforementioned susceptible crops. These plants are most sensitive to **OUTLAW™** during their development or growing stage. Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of **OUTLAW™** with the roots of desirable trees and shrubs.

Drift Reduction Information:

The following information may be helpful in reducing possible spray drift from ground or aerial applications. Avoid making applications when spray particle may be carried by air currents to areas where sensitive crops and plants are growing. Do not spray near sensitive plants if the wind is gusty or in excess of 5 mph and moving in the direction of nearby sensitive crops or if a temperature inversion exists. Always determine the direction and distance of possible spray drift prior to application. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays. Properly maintain and calibrate all spray equipment. The use of agriculturally accepted drift retardants are acceptable and advised. Avoid applications within the vicinity of susceptible plants when at all possible. Do not apply in greenhouses.

AERIAL APPLICATION METHODS AND EQUIPMENT

10/23

Water Volume: Use 3-10 gallons of water per acre. Use the higher spray volume when treating dense or tall vegetation.

Application Equipment: Select nozzles designed to produce minimal amounts of fine spray particles. Make applications at the lowest stage height to reduce the exposure of spray droplets to evaporation and wind. The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in the this labeling as well as applicable state and local regulations and ordinances.

SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

AERIAL DRIFT REDUCTION ADVISORY

[This section is advisory in nature and does not supersede the mandatory label requirements.]

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

11/23

Table 2. Application Rate and Timing - Biennial and Perennial Weeds.

(The addition of liquid fertilizer (28-0-0,32-0-0) at ½ lb/acre)

spray solution has proven to give increase suppression or control of certain species of weeds.)

(certain species of weeds.)

13/23

| Weeds Controlled | Rate Per Acre (according to weed growth stage) | | | | | |
|-------------------------------------|--|-----------|------------------|-----------------|----------------|---------------------------|
| | 1.5 pints | 1.5 pints | 1.5 pints | 1.75 pints | 2.75 pints | 3.5-5.25 pints |
| Bindweed, Field | - | - | - | - | - | actively growing |
| Bittercress | - | 2-3" | - | - | - | - |
| Buckeye species ¹ | - | - | - | - | full leaf | - |
| Bullnettle ² | - | - | - | flower | - | - |
| Chicory | - | - | - | - | early bolting | - |
| Clove, Bur | - | - | Pre-flower | - | - | - |
| Dandelion, Common | - | rosette | - | bolting | - | - |
| Dewberry, Southern ¹ | - | - | - | - | - | spring or fall |
| Dock, Curly | - | - | prior to bolting | - | after bolting | - |
| Elderberry ² | - | - | - | - | - | actively growing |
| Goldenrod, Missouri | - | - | - | 3-15" | flower | - |
| Groundsel, Texas | - | rosette | post-bolting | - | - | - |
| Honeysuckle, Hairy | - | - | - | - | spring or fall | - |
| Horsenettle, Carolina ¹ | - | - | - | - | - | flower or berry |
| Ivy, Poison | - | - | - | after bloom | - | - |
| Knapweed, Black ² | - | - | - | - | - | actively growing |
| , Russian ² | - | - | - | - | - | actively growing |
| , Spotted | - | - | - | - | - | actively growing |
| Marshelder | - | - | - | <12" | 12"/prebloom | - |
| Mesquite ³ | - | - | - | - | - | 45-90 days after budbreak |
| Milkweed, Antelopehorn ² | - | - | - | pre-flower | - | Flower |
| Nightshade, Silverleaf ¹ | - | - | - | full flower | - | - |
| , Black ¹ | - | - | - | full flower | - | actively growing |
| Persimmon, Eastern ³ | - | - | - | - | - | actively growing |
| Prickly, Lettuce | - | - | - | rosette | - | actively growing |
| Rabbitbrush ² | - | - | - | - | - | - |
| Ragwort, Tansy | - | - | - | rosette | - | actively growing |
| Redvine ² | - | - | - | - | - | actively growing |
| Sagebrush, Fringed ² | - | - | - | - | - | actively growing |
| Smartweed | - | - | - | - | - | - |
| Sorrel, Red | - | - | Rosette | bolting | flower | actively growing |
| Sowthistle ² | - | - | - | - | - | actively growing |
| Spurge, Leafy ² | - | - | - | - | - | full leaf |
| Tallow Tree, Chinese ⁴ | - | - | - | - | - | - |
| Thistle, Bull | - | - | Rosette | bolting | - | actively growing |
| , Canada ² | - | - | - | - | - | - |
| , Musk | - | - | - | rosette/bolting | - | - |
| , Plumeless | - | - | Rosette | bolting | - | - |
| Vetch, Hairy | - | 1-4" | 4-8" | 8" full flower | - | - |
| Yankeeeweed | - | - | - | 10-18" | - | Rosette |
| Yellow Starthistle ¹ | - | - | - | - | - | - |

¹ May require repeat applications

² Recommended rate will provide top growth suppression only.

³ For improved root kill or woody species such as mesquite and eastern persimmon spray 3.5 pints of per acre **OUTLAW™** each year for 3 consecutive years.

⁴ Under dense populations, a second application may be needed the following growing season.

For increased control of weeds such as blackberry and dewberry, **OUTLAW™** may be tank mixed with Ally® herbicide (0.1-0.2 ounces per acre), if labeled for the use site.

Ground Application (Banding)

When applying **OUTLAW™** herbicide by banding, determine the amount of herbicide and water volume needed using the following formula:

14/23

$$\frac{\text{Bandwidth in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Banding herbicide rate per acre}$$

$$\frac{\text{Bandwidth in inches}}{\text{Row width in inches}} \times \text{Broadcast rate volume per acre} = \text{Banding water volume per acre}$$

Ground Application (Broadcast)

Water volume: Use 10-25 gallons of spray solution per broadcast acre for optimal performance. Use the higher spray volume when treating dense or tall vegetation.

Application Equipment: Select nozzle design to produce minimal amounts of fine spray particles. Spray nozzles as close to the weeds as is practical for good weed coverage.

Spot or Small Area Application

OUTLAW™ may be applied to individual clumps or small areas of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems. For knapsack or other small capacity sprayers, prepare a solution of **OUTLAW™** in water according to **Table 3** (assuming that the spot treatment rate equates to 40 gallons pre acre on the broadcast basis.) Adding a surfactant (0.5% by volume) can help improve control.

Do not make spot treatments in addition to broadcast or band treatments.

Application equipment: Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

Table 3. – Knapsack Sprayer Dilution Instructions

| Sprayer Capacity (gallons of water) | Amount of OUTLAW™ to add to the spray tank |
|--|--|
| 1 gallon | 1 fluid ounce* |
| 3 gallons | 3 fluid ounces |
| 5 gallons | 5 fluid ounces |

* 1 fluid ounce = 2 tablespoons

III. Additives

To improve burndown of emerged weeds, surfactants and/or low use rates of liquid fertilizers (28-0-0; 32-0-0), or crop oil concentrate may be used with **OUTLAW™** herbicide or **OUTLAW™** tank mixes applied after the weeds have emerged. Crop oil concentrate is for non-food/feed crop uses only. Do not apply tank mixes that include Ammonium Sulfate or Crop Oil Concentrate to any food/feed crop use listed on this label. For food/feed crop use, do not use liquid fertilizers that contain Ammonium Sulfate (AMS) as a source of nitrogen as tolerances in commodities derived from the crop may contain residues that exceed established tolerances.

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic
- contain only EPA-exempt ingredients
- provide good mixing quality in the jar test, and
- be successful in local experience

The exact composition of suitable products will vary; however, vegetable oil and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, see **Compatibility Test for Mix Components**.

Adjuvants containing crop oil concentrate may be used for preplant, pre-emergence and between cropping applications. Do not use crop oil concentrate for postemergence applications in food/feed crops (i.e. grass (silage), pastures, rangeland, and wheat.)

15/23

Nitrogen Source

- **Sprayable liquid fertilizers:** Use ½ GPA of sprayable liquid fertilizers (28-0-0; 32-0-0) per acre. Do not use brass or aluminum nozzles when spraying fertilizers.

Non-ionic Surfactant

The standard label recommendation is 2-4 pints of an 80% active non-ionic spray surfactant per 100 gallons of water. (Rate will vary with the size and condition of weeds to be controlled. Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.)

Table 4. Additive Rate Per Acre.

| Additive ¹ | Rate OUTLAW™ Per Acre |
|---|--|
| Non-ionic Surfactant | 2-4 pints per 100 gallons ² |
| Sprayable Liquid Fertilizers (28-0-0; 32-0-0) | ½ GPA of spray solution |
| Crop Oil Concentrate | 1 quart |

¹ See manufacturer's label for specific rate recommendations.

² Use lowest rate per 100 gallons when weeds are small and actively growing. As weeds increase in size and or become hardened off, the rate of non-ionic surfactant will have to be increased to give optimum coverage and control.

IV. General Tank Mixing Information

Tank Mix Partners/Components

The following products may be tank mixed with OUTLAW™ according to the specific tank mixing instructions in this label and respective product labels.

- Aim™ (carfentrazone-ethyl)
- Ally® (metsulfuron-methyl)
- Amber® (triasulfuron)
- Asulox® (asulam)
- Atrazine
- Banvel® (dicamba)
- Clarity® (dicamba)
- Curtail™ (clopyralid + 2,4-D)
- Cyclone® (paraquat)
- Dicamba DMA (dicamba)
- Distinct® (diflufenzopyr + dicamba)
- Evik® (ametryn)
- Fallowmaster® (glyphosate + dicamba)
- Fallow Star™ (glyphosate + dicamba)
- Finesse® (chlorsulfuron + metsulfuron-methyl)
- Gly Star™ Plus (glyphosate)
- Gramoxone® Extra (paraquat)
- Grazon™ P+D (picloram + 2,4-D)
- Kerb™ (pronamide)
- Landmaster® (glyphosate + 2,4-D)
- MCPA
- Paramount® (quinclorac)
- Rave™ (dicamba + triasulfuron)
- Roundup® Ultra (glyphosate)
- Sencor® (metribuzin)

16/23

- Sinbar® (terbacil)
- Tordon™ (picloram)
- Touchdown® (glyphosate)
- 2,4-D

See "VI. Food/Feed Crop Specific Information" section for more information for more details. Read and follow the applicable **Restrictions and Limitations and Directions for Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes. Physical incompatibility, reduced weed control, or crop injury may result from mixing **OUTLAW™** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes adjust accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order

If an inductor is used, rinse it thoroughly after each component has been added. Maintain constant agitation during application.

1. **Water** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
2. **Agitation.** Maintain constant agitation throughout mixing and application.
3. **Products in PVA bags.** Place any product contained in water-soluble bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, and suspo-emulsions)
5. **Water-soluble products (such as OUTLAW™).**
6. **Emulsifiable concentrates** (such as oil concentrate, when applicable).
7. **Water-soluble additives** (such as liquid fertilizers (28-0-0; 32-0-0), when applicable).*
8. **Remaining quantity of water.**

* If sprayable fluid fertilizer is used as the carrier.

Always perform the **Compatibility Test** before mixing into the spray tank. Also, when using a sprayable fluid fertilizer as the carrier, any product contained in PVA bags must first be completely dissolved in water before the contents can be added to the fertilizer mix.

V. Restrictions and Limitations

- **Maximum seasonal use rate:** Refer to Table 5.
- **Preharvest Interval (PHI):** Refer to "Food/Feed Crop Specific Information"
- **Restricted entry Interval (REI):** 48 Hours
- **Crop Rotational Restrictions:**
- The interval between application and planting rotational crop is given below. Always exclude counting days when the ground is frozen. Planting at intervals less than specified below may result in crop injury. Moisture is essential for the degradation of this herbicide in soil.

Planting/replanting restrictions for OUTLAW™ applications of 6 pints per acre or less: No rotational cropping restrictions apply at 120 days or more following application. Additionally, for annual crop uses in this label including sorghum, follow the preplant use directions under "VI. Food/Feed Crop Specific Information." For barley, oat, wheat, and other grass seedlings, the interval between application and planting is 10 days per pint per acre.

Planting/replanting restrictions for applications of more than 6 pints and up to 8 pints of OUTLAW™ per acre: Corn, sorghum, cotton (east of the Rocky Mountains) and all other crops grown in areas with 30" or more of annual rain fall may be planted 120 days or more after application. Barley, oat, wheat and other grass seedlings may be planted, if the interval from

application to planting is 10 days per pint per acre east of the Mississippi River and 15 days per pint per acre west of the Mississippi River. For all other areas with less than 30" of annual rainfall, the interval between application and planting is 180 days or more.

17/23

- **Arid (dry) conditions:** it is extremely important that the addition of a suitable Nonionic Surfactant, Oil, or sprayable fertilizer be used when applying **OUTLAW™**. Higher rates of **OUTLAW™** may be needed to control susceptible weeds in this environment.
- **Rainfast Period:** Rainfall or irrigation occurring within 4 hours after postemergence applications may reduce effectiveness of **OUTLAW™**.
- **Stress:** Do not apply to crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result.
- Do not apply to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Do not apply this product through any type of **irrigation** equipment. Do not contaminate irrigation ditches or water used for domestic purposes.
- This product cannot be used to **formulate** or reformulate another pesticide product.

Table 5. Crop Specific Restrictions and Limitations.

| Crop | Maximum Rate Per Acre Per Application | Maximum Rate Per Acre Per Season | Livestock Grazing or Feeding ¹ | Aircraft Application |
|---------------------------|---------------------------------------|----------------------------------|---|----------------------|
| Between Crop Applications | 6 pints | 8 pints | Yes | Yes |
| Pasture, Hay, Silage | 4 pints | 8 pints | Yes | Yes |
| Wheat | 2 pints | 3.33 pints | Yes | Yes |

¹ Refer to "Food/Feed Crop Specific Information" for grazing and feeding restrictions.

VI. Food/Feed Crop Specific Information

Pastures, Rangeland and Grass (Hay, Silage)

OUTLAW™ is recommended for use for pasture (including pasture grown for hay), rangeland, grass grown for hay or silage, between crop applications/fallow systems, Conservation Reserve Programs, and general farmstead (non-cropland only).

Refer to **Tables 1 and 2** for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

Rates above 4 pints of **OUTLAW™** per acre are for spot treatments only.

Retreatments may be made as needed; however, do not exceed a total of 8 pints of **OUTLAW™** per treated acre during a growing season.

Uses described in this section also pertain to small grains (such as barley, corn, forage sorghum, oats, rye, sudangrass, or wheat) grown for pasture, hay, and silage only. Newly seeded areas including small grains grown for pasture or hay, may be injured if rates of **OUTLAW™** are greater than 2 pints per acre are applied.

In newly established hybrid Bermudagrass, Pangolagrass, and stargrasses (*Cynodon* spp.) use 1.75 to 3.5 pints of **OUTLAW™** per acre to control or suppress weeds after planting vegetative propagules (stolens) of hybrid bermudagrasses. In addition to the weeds listed in **Tables 1 and 2**, this rate of **OUTLAW™** will control or suppress annual sedges, broadleaf signalgrass, crabgrass, and goosegrass. Best results will be obtained if **OUTLAW™** is applied at the germinating stage of weeds. Under favorable conditions, this is usually 7-10 days after planting these grasses. Reduced control can be expected if weeds are allowed to reach 1" in height before application or if germination of weeds occurs 10 days after application.

Do not use on bentgrass, susceptible grass pastures (such as carpetgrass, buffalograss, or St. Augustine grass), lezpedeza, wild winter peas, vetch, clover, and alfalfa pastures as injury will occur.

When perennial weeds are reaching maturity, mowing and allowing some regrowth will enhance control. Difficult to control weeds may require a repeat application.

For pasture renovations, wait 3 weeks per quart (2 pints) of **OUTLAW™** used per acre before interseeding or injury may occur.

If grasses are grown for seed or for seed () purposes, do not apply after grass reaches joint ()

18/23

Grazing and Feeding Non-Lactating Animals: There is no waiting period between treatment and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

Grazing and Feeding Lactating Animals: Do not graze lactating dairy animals within 7 days of treatment.

Dry hay and Silage: Treated grasses may be harvested for dry hay or silage but do not harvest within 37 days of treatment.

Pasture and Rangeland Tank Mixes

OUTLAW™ may be applied in tank mixes with one or more of the following herbicides:

| | | | |
|--------|----------|-------------|-------|
| Ally® | Barivel® | Dicamba DMA | 2,4-D |
| Amber® | Clarity® | Rave™ | |

Wheat

PREHARVEST APPLICATIONS:

OUTLAW™ can be used to control weeds that may interfere with harvest of wheat. Apply up to 2 pints of **OUTLAW™** per acre as a broadcast or spot treatment to annual broadleaf weeds when wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing but before weeds canopy. A waiting interval of 7 days is required before harvest. Do not use preharvest-treated wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better. For control of additional broadleaf weeds or grasses, **OUTLAW™** may be tank mixed with other herbicides such as **Ally** or **Gly Star™ Plus** that are registered for preharvest use in wheat. Preharvest use of **OUTLAW™** is not registered for use in California.

Between Crop Applications/Fallow Systems, Conservation Reserve Programs, and General Farmstead

These uses are considered Food/Feed Crops when harvested, grazed or foraged. Consult section on "General Tank Mixing Information" for adjuvant restrictions and section on "Additives" for specific use directions.

VII. Non-Food/Feed Use (Land not Harvested, Grazed or Foraged)-Specific Information.

Between Crop Applications

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL

OUTLAW™ can be applied postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply to weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

See "V. Restrictions and Limitations" for the recommended interval between application and planting to prevent crop injury.

Rates and Timings:

Apply 1.5-6 pints of **OUTLAW™** per acre. Refer to Table 1 to determine use rates for specific targeted weed species. Retreatments may be made as needed; however, do not exceed a total of 8 pints of **OUTLAW™** per treated acre during a growing season. For best performance, apply **OUTLAW™** when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if **OUTLAW™** is applied when the majority of weeds have at least 4-6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage. The addition of liquid fertilizers (28-0-0,32-0-0) at 1/2 GPA has shown to increase efficacy.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for **OUTLAW™**. For seedling control, a follow-up program or other cultural practices could be instituted.

Between Crop Tank Mixes:

In tank mixes with one or more of the following herbicides, apply 1.5-1.75 pints of **OUTLAW™** per acre for control of annual weeds, or 1.75-7.25 pints of **OUTLAW™** per acre for control of biennial and perennial weeds

- Aim™
- Ally®
- Amber®
- Atrazine
- Curtail™
- Cyclone®

19/
23

- Distinct®
- Fallowmaster®
- Fallow Star™
- Finesse®
- Glyphosate (Gly Star™ Plus)
- Gramoxone® Extra
- Kerb™
- Landmaster® BW
- Paramount®
- Sencor®
- Tordon™ 22K
- Touchdown®
- 2,4-D

Conservation Reserve Programs and General Farmstead

OUTLAW™ is recommended for use for Conservation Reserve Programs, general farmstead (non-cropland only), weed and brush control, or use in State Recognized Noxious Weed areas (non-cropland areas).

Refer to **Tables 1 and 2** for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

Rates above 4 pints of **OUTLAW™** per acre are for spot treatments only.

Retreatments may be made as needed; however, do not exceed a total of 8 pints of **OUTLAW™** per treated acre during a growing season.

Farmstead and Fence-row Treatment Application Instructions

OUTLAW™ may be applied using water or oil and water emulsions in spot application to control undesirable vegetation using handgun or similar types of application equipment. In addition to weed species listed in **Tables 1 and 2**, these treatments may be used to control or suppress woody plant species listed in **Table 7**.

To prepare soil and water emulsions, mix in the order and proportions indicated below.

The solution should remain milky colored without an oily layer on top when under agitation. If an oily layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

Do not exceed 40 gallons of spray solution per treated acre per application. One gallon of **OUTLAW™** in forty gallons of spray solution contains 1.09 pounds acid equivalent of dicamba and 1.45 pounds acid equivalent of 2,4-D. Spray plants to wet. Do not allow this spray mix to contact desirable vegetation.

To control brush, briars, and weeds along fence-rows surrounding pasture and ranch lands, and fallow fields, use a tank mix of 2.5% **OUTLAW™**, 87.5% water, 10% diesel oil, and sufficient emulsifier (to mix the diesel and emulsifier). The diesel oil in this tank mix will damage or kill desirable grasses and should not be used in pastures or where damage to desirable species cannot be tolerated.

- 1). Water: Begin by agitating a thoroughly clean sprayer tank with the desired quantity of clean water. Maintain constant agitation during complete mixing procedure.
- 2). Emulsifier: Add 0.5% volume to volume of water.
- 3). **OUTLAW™**: add 2.5 gallons per 100 gallons of total intended solution.
- 4). Diesel Oil: Add 10 gallons per 100 gallons of total intended solution.

Maintain constant agitation during application. Under good agitation, the spray solution should be milky white with no oil layer on top. If oil layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

FOR SPRAYING FOLIAR APPLICATIONS:

1. Spray when leaves have reached full size but have not hardened due to drought or maturity.
2. Spray individual plants to wet with handgun.
3. For larger stems (up to 3" in diameter) and hard to control species, direct spray stream to base of stems to wet the stem at soil surface in addition to wetting the foliage.
4. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FOR DORMANT BASAL APPLICATIONS:

1. Increase diesel oil content to 15% or 15 gallons of diesel oil per 100 gallons of total solution.
2. Spray in late winter and early spring before plants break dormancy.

20/23

3. Spray the bottom 24" of the target stem to wet on all sides.
4. For larger stems (up to 3" in diameter) direct the spray solution to () of target stems to wet the soil at the stem/soil junction in addition to wetting the stem.
5. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

FOR CUT SURFACE TREATMENTS:

Apply **OUTLAW™** in an undiluted state as a cut surface treatment to control unwanted trees and prevent sprouts of cut trees.

- **Frill or Girdle Treatments:** Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint the cut surface with **OUTLAW™**.
- **Stump Treatments:** Spray or paint freshly cut surface with **OUTLAW™**. The cambium layer (the area adjacent to the bark) should be thoroughly wet. Treat stumps within 6 hours after cutting.

Table 7. The following list of trees and vines can be controlled on farmsteads and fencerows as foliar, basal, or cut surface treatments:

| | |
|-----------------------|------------------------|
| Alder | Kudzu |
| Ash | Locust, Black |
| Aspen | Maple |
| Basswood | Mesquite |
| Beech | Oak |
| Blackberry | Oak, Poison |
| Blackgum | Olive, Russian |
| Cedar | Persimmon, Eastern |
| Cherry | Pine |
| Chinquapin | Plum, Sand (Wild Plum) |
| Cottonwood | Poplar |
| Creosotebush | Rabbitbrush |
| Dewberry | Redcedar, Eastern |
| Dogwood | Rose, McCartney |
| Elm | Rose, Multiflora |
| Grape | Sagebrush, Fringe |
| Greenbriar | Sassafras |
| Hawthorn (Thornapple) | Spruce |
| Hemlock | Sumac |
| Hickory | Sweetgum |
| Honeylocust | Sycamore |
| Honeysuckle | Tarbrush |
| Hornbeam | Willow |
| Huckleberry | Witchhazel |
| Huisache | Yaupon |
| Ivy, Poison | Yucca |

Weeds listed in this label:

| Common Name | Scientific Name |
|--------------------------|-----------------------------------|
| ANNUALS | |
| Beebalm, Spotted | <i>Monarda punctata</i> |
| Broomweed, Common | <i>Gutierrezia dracunculoides</i> |
| Buckwheat, Wild | <i>Polygonum convulvulus</i> |
| Buffalobur | <i>Solanum rostratum</i> |
| Burdock | <i>Arctium spp.</i> |
| Buttercup, Corn | <i>Ranunculus arvensis</i> |
| Chickweed, Common | <i>Stellaria media</i> |
| Cockle, Corn | <i>Agrostemma githago</i> |
| Cocklebur, Common | <i>Xanthium strumarium</i> |
| Coreopsis, Plains | <i>Coreopsis tinctoria</i> |
| Croton, Woolly | <i>Croton capitatus</i> |
| Devilsclaw, | <i>proboscidea luisianica</i> |
| Dogfennel (Cypressweed) | <i>Eupatorium capillifolium</i> |
| Eveningprimrose, Cutleaf | <i>Oenothera lacinata</i> |
| Flax | <i>Linum catharticum</i> |
| Fleabane, Annual | <i>Erigeron annuus</i> |
| Flixweed | <i>Descurainia sophia</i> |
| Henbit | <i>Lamium amplexicaule</i> |

| Common Name | Scientific Name |
|--------------------------|---------------------------------|
| Knotweed, Prostrate | <i>Polygonum aviculare</i> |
| Kochia | <i>Kochia scoparia</i> |
| Lambsquarters, Common | <i>Chenopodium album</i> |
| Lettuce, Prickly | <i>Lactuca serriola</i> |
| Mallow, Common | <i>Malva neglecta</i> |
| Mornigglory, Ivyleaf | <i>Ipomea hederacea</i> |
| Tall | <i>Ipomea purpurea</i> |
| Mustard, Annual | <i>Brassica spp.</i> |
| Tansy | <i>Descurainia pinnata</i> |
| Pennycress, Field | <i>Thlaspi arvense</i> |
| Pepperweed, Virginia | <i>Lepidium virginicum</i> |
| Pigweed, Prostrate, | <i>Amaranthus blitoides</i> |
| Redroot, | <i>Amaranthus retroflexus</i> |
| Smooth, | <i>Amaranthus hybridus</i> |
| Tumble | <i>Amaranthus albus</i> |
| Poorjoe | <i>Diodia teres</i> |
| Purslane, Common | <i>Portulaca oleracea</i> |
| Ragweed, Common, | <i>Ambrosia artemisiifolia</i> |
| Lance-leaf, | <i>Ambrosia bidentata</i> |
| Western | <i>Ambrosia psilostachya</i> |
| Sedge | <i>Cyperus compressus</i> |
| Shepherdspurse | <i>Capsella bursa-pastoris</i> |
| Smartweed, Pennsylvania | <i>Polygonum pennsylvanicum</i> |
| Sneezeweed, Bitter | <i>Helenium amurum</i> |
| Sunflower, Common (wild) | <i>Helianthus annuus</i> |
| Thistle, Russian | <i>Salsola iberica</i> |

24
/23

24
/23

| Common Name | Scientific Name |
|---------------------------------|-----------------------------------|
| BIENNIALS AND PERENNIALS | |
| Bindweed, field | <i>Convolvulus arvensis</i> |
| Bittercress | <i>Cardamine spp.</i> |
| Buckeye | <i>Aesculus spp.</i> |
| Bullnettle | <i>Cnidoscopus stimulosus</i> |
| Chicory | <i>Cichorium intybus</i> |
| Clover, Hop | <i>Trifolium aureum</i> |
| Dandelion | <i>Taraxacum officinale</i> |
| Dock, Curly | <i>Rumex crispus</i> |
| Elderberry | <i>Sambucus canadensis</i> |
| Goldenrod, Missouri | <i>Solidago missouriensis</i> |
| Goldenweed, Common | <i>Isocyp, a cprpmopifolia</i> |
| Groundset | <i>Senecio vulgaris</i> |
| Honeysuckle, Hairy | <i>Lonicera</i> |
| Horsenettle | <i>Solanum carolinense</i> |
| Ivy, Poison | <i>Rhus radicans</i> |
| Knapweed, Black | <i>Centaurea nigra</i> |
| Russian | <i>Centaurea repens</i> |
| Spotted | <i>Centaurea maculosus</i> |
| Marshelder | <i>Ina annua</i> |
| Mesquite | <i>Prosopis juliflora</i> |
| Milkweed, Antelopehorn | <i>Asclepius</i> |
| Nightshade, Silverleaf | <i>Solanum elaeagnifolium</i> |
| Black | <i>Solanum nigrum</i> |
| Persimmon, Eastern | <i>Diospyros virginiana</i> |
| Rabbitbrush | <i>Chrysanthemum pulchellus</i> |
| Ragwort, Tansy | <i>Senecio jacobia</i> |
| Redvine | <i>Brunnichia ovata</i> |
| Sagebrush, Fringed | <i>Artemisia frigida</i> |
| Smartweed, Swamp | <i>Polygonum coccineum</i> |
| Sorrel, Red (Sheep Sorrel) | <i>Rumex acetosella</i> |
| Sowthistle, Perennial | <i>Sonchus arvensis</i> |
| Spurge, Leafy | <i>Euphorbia esula</i> |
| Starthistle, Yellow | <i>Centaurea solstitialis</i> |
| Tallow Tree, Chinese | <i>Sapium sebiferum</i> |
| Thistle, Bull | <i>Cirsium vulgare</i> |
| Canada | <i>Cirsium arvense</i> |
| Musk | <i>Carduus nutans</i> |
| Plumeless | <i>Carduus acanthoides</i> |
| Vetch | <i>Vicia spp.</i> |
| Yankeeeweed | <i>Eupatorium compositifolium</i> |

Food/Feed Crop Uses

This product can be used on the following:

- Conservation Reserve Program Land
- Fallow Systems (Between Crop Application)
- General Farmstead

Grass (Hay or Silage)

Pastures

Rangeland

Wheat

Look inside for complete **Restrictions and Limitations** and **Application Instructions**

- These crops are considered Food/Feed crops only when harvested, grazed, or foraged. Otherwise, they are considered non-Food/Feed uses.

CONDITIONS OF SALE AND WARRANTY

The **DIRECTIONS FOR USE** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions or presence of other materials. All such risks shall be assumed by the Buyer.

HELENA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **HELENA MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.**

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **IN NO CASE SHALL HELENA OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** HELENA and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of HELENA. No employee or agent of HELENA or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

Ally®, Finesse®, and Sinbar® are registered trademarks of E. I. du Pont de Nemours & Company.

Banvel®, Clarity®, Distinct®, Fallowmaster®, and Paramount® are registered trademarks of BASF Corporation.

Asulox® is a registered trademark of Bayer CropScience.

Amber®, Evik®, Cyclone®, Sencor®, Gramoxone®, Rave™ and Touchdown® are trademarks and/or registered trademarks of Syngenta Crop Protection.

Landmaster® and Roundup® Ultra are registered trademarks of Monsanto Company.

Curtail™, Grazon™, Kerb™ and Tordon™ are trademarks of Dow AgroSciences LLC.

Aim™ is a trademark of FMC Corporation.

Fallow Star™ and Gly Star™ Plus are trademarks of Albaugh, Inc.

The contents of this package and/or use of the contents are covered by U.S. Patent Nos. 5,877,112; 6,232,272; 5,580,567 and 5,741,502.

23/
23