

5905-578

6/2/2014

1/45



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Bill Washburn  
Helena Chemical Company  
225 Schilling Boulevard, Suite 300  
Collierville, TN 38017

JUN -2 2014

Subject: Notification per PR Notice 98-10 – Addition of state restriction  
Product Name: TAPOUT  
EPA Registration Number: 5905-578  
Application Dated: May 23, 2013  
Decision Number: 479037

Dear Mr. Washburn:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product. Our review finds that the action falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and placed in our records.

If you have any questions, please contact Sarah Meadows at (703)347-0505 or at meadows.sarah@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "K. Montague".

Kathryn V. Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

2/45

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 05-31-98

|   |  |                               |
|---|--|-------------------------------|
| <b style="font-size: 1.2em;">EPA</b><br>United States<br><b>Environmental Protection Agency</b><br>Washington, DC 20460 | <input type="checkbox"/> <b>Registration</b><br><input type="checkbox"/> <b>Amendment</b><br><input checked="" type="checkbox"/> <b>Other:</b> | OPP Identifier Number<br><br> |
|---|--|-------------------------------|

**Application for Pesticide - Section I**

|  |   |  |
|--|---|--|
| 1. Company/Product Number<br><b>5905-578</b>   | 2. EPA Product Manager<br><b>Kathryn Montague</b> | 3. Proposed Classification<br><input type="checkbox"/> None <input type="checkbox"/> Restricted  |
| 4. Company/Product (Name)<br><b>TAPOUT</b>   | PM#<br><b>23</b>                                  |  |
| 5. Name and Address of Applicant (Include ZIP Code)<br><br><b>Helena Chemical Company</b><br><b>225 Schilling Boulevard, Suite 300</b><br><b>Collierville, TN 38017</b><br><br><input type="checkbox"/> Check if this is a new address |   | 6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to:<br><p style="text-align: right; font-weight: bold; font-size: 1.2em;">NOTIFICATION</p> EPA Reg. No. _____<br><br>Product Name <span style="float: right; font-weight: bold; font-size: 1.2em;">JUN - 2 2014</span> |

**Section - II**

|  |  |
|--|--|
| <input type="checkbox"/> Amendment - Explain below.                            | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application  |
| <input checked="" type="checkbox"/> Notification - Explain below.              | <input type="checkbox"/> Other - Explain below   |

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)  
**Notification of addition of California crop restriction per PR Notice 98-10.** This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

1. Material This Product Will Be Packaged In:

|   |   |   |  |
|---|---|---|--|
| Child-Resistant Packaging<br><input type="checkbox"/> Yes*<br><input type="checkbox"/> No | Unit Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No<br>If "Yes"<br>Unit Packaging wgt.    No. per container | Water Soluble Packaging<br><input type="checkbox"/> Yes<br><input type="checkbox"/> No<br>If "Yes"<br>Package wgt.    No. per container | 2. Type of Container<br><input type="checkbox"/> Metal<br><input type="checkbox"/> Plastic<br><input type="checkbox"/> Glass<br><input type="checkbox"/> Paper<br><input type="checkbox"/> Other (Specify) |
|---|---|---|--|

**\*Certification must be submitted**

|  |                             |   |
|--|-----------------------------|---|
| 3. Location of Net Contents Information<br><input type="checkbox"/> Label <input type="checkbox"/> Container | 4. Size(s) Retail Container | 5. Location of Label Directions<br><input type="checkbox"/> On Label<br><input type="checkbox"/> On labeling accompanying product |
|--|-----------------------------|---|

6. Manner in Which Label is Affixed to Product     Lithograph     Other \_\_\_\_\_  
 Paper glued  
 Stenciled

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)

|                              |   |  |
|------------------------------|---|--|
| Name<br><b>Bill Washburn</b> | Title<br><b>Registration Specialist</b> | Telephone No. (Include Area Code)<br><b>901-752-4420</b> |
|------------------------------|---|--|

**Certification**

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

|                                       |  |  |
|---------------------------------------|--|--|
| 2. Signature<br>                      | 3. Title<br><b>Registration Specialist</b> | 6. Date Application Received (Stamped) |
| 4. Typed Name<br><b>Bill Washburn</b> | 5. Date<br><b>May 23, 2013</b>             |  |

31/45



**HELENA CHEMICAL COMPANY**

225 Schilling Blvd., Suite 300  
Collierville, Tennessee 38017  
Telephone: 901-761-0050

May 23, 2013

Ms. Kathryn Montague, PM 23  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504P)  
U.S. Environmental Protection Agency  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501

**Subject: TAPOUT  
EPA Reg. No. 5905-578**

**Notification of the Addition of California Crop Restrictions  
per PR Notice 98-10**

Dear Ms. Montague:

It was recently brought to our attention that while the restriction of using Tapout in California is mentioned in two different places for both Canola and Flax, the restriction is not present in the tank-mix section of the label. Therefore, we are submitting this notification to add "Not for Use in California" to the Canola and Flax tank-mix instructions on the subject label. The following documents are submitted in support of this notification:

- EPA Form 8570-1 (Application for Pesticide: Other)  
One copy of EPA Form 8570-1 dated May 23, 2013 addressing the need for a notification.
- 2 Copies of Revised Labeling  
One copy with changes tracked identified by "STK" in the filename at the bottom of the label and on clean copy identified by the "CLN" in the filename at the bottom of the label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Should you have any questions or comments, please do not hesitate to contact me at 901-752-4420 or by e-mail at [washburnb@helenachemical.com](mailto:washburnb@helenachemical.com)

Sincerely,

Bill Washburn  
Registration Specialist

JUN -2 2014

7/  
45**TAPOUT®**

| Active Ingredient      | By Wt. |
|------------------------|--------|
| *Clethodim .....       | 12.6%  |
| Other Ingredients..... | 87.4%  |
| Total.....             | 100.0% |

Contains Petroleum Distillates

\*(E).2..[1 -[[[(3-chloro-2-propenyl)oxyimino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1 -one

Contains 0.97 lbs. Clethodim per gal.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### FIRST AID

|                                |   |
|--------------------------------|---|
| <b>IF SWALLOWED:</b>           | <ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
| <b>IF IN EYES:</b>             | <ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor immediately for advice</li> </ul>                                      |
| <b>IF ON SKIN OR CLOTHING:</b> | <ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>   |
| <b>IF INHALED:</b>             | <ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>                                  |

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (800) 424-9300, collect day or night, for emergency medical treatment information.

#### NOTE TO PHYSICIAN

Ingestion of this product or subsequent vomiting can result in aspiration of light hydrocarbon liquid, which can cause pneumonitis. If ingested, probable mucosal damage may contraindicate the use of gastric lavage. May pose an aspiration pneumonia hazard. Contains petroleum distillate.

EPA Reg. No. 5905-578  
EPA Est. No.  
AD 021811  
NET CONTENTS

Manufactured For  
Helena Chemical Company  
225 Schilling Boulevard, Suite 300  
Collierville, TN 38017

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate or viton ≥14 mils, shoes plus socks and protective eyewear.

**Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.** Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwater or rinsate.

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

~~Solano Grass: Solano County, California: the vernal lakes area bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south and Travis Air Force Base to the west.~~

Wild Rice: Hays County, Texas.

**PHYSICAL OR CHEMICAL HAZARDS:**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval. (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls over short-sleeved shirt and short pants, chemical-resistant gloves such as barrier laminate or viton ≥14 mils and shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep all unprotected persons out of operating areas, or vicinity where there may be drift. Do not enter treated areas without protective clothing until sprays have dried.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance in proper disposal methods.

**CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon):** Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Non-Refillable (<5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Non-Refillable (>5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container (250 gallon & bulk):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**TANK MIXES**

**NOTICE:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

**TABLE OF CONTENTS**

**PRODUCT INFORMATION**

Control Symptoms

**APPLICATION INFORMATION**

Timing of Applications

Ground Application

Air Application

Spot Treatment

Chemigation — Garlic and Onion (Dry Bulb and Green) Sprinkler Irrigation Application

Use Precautions

**RESTRICTIONS AND LIMITATIONS**

- All Uses
- Spray Drift Management
- Resistance Management

**Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT**

- Directions for Annual Grasses
- Directions for Annual & Perennial Grass Control in Established Alfalfa and Mint
- Directions For Reduced Rate Use To Control Small Annual Grasses Table
- Directions for Perennial Grasses
- Directions for Annual Bluegrass Control

**TANK MIXES - PRODUCT INFORMATION**

Tank Mix Application of TAPOUT and Broadleaf Herbicides for Control of Grasses and Broadleaf Weeds.

- Mixing Instructions
- Information on Antagonism

**Alfalfa**

Table 2. TAPOUT Tank Mixes with Broadleaf Herbicides for Alfalfa.....

**Canola**

Table 3. Reduced Rate TAPOUT Tank Mixes with Broadleaf Herbicides for Canola.....

**Cotton**

Table 4: TAPOUT Tank Mixed with Cobra and MSMA applied Post Directed to Cotton.....

Table S. TAPOUT Tank Mixed with Glyphosate to Control Emerged Grasses in Cotton as a Broadcast Application.....

**Dry and Succulent Shelled Beans**

Table 6. TAPOUT Tank Mixes with Broadleaf Herbicides for Dry and Succulent Shelled Beans.....

**Flax**

Table 7. Reduced Rate TAPOUT Tank Mixes with Broadleaf Herbicides for Flax.....

**Peanut**

Table 8. TAPOUT Tank Mixes with Broadleaf Herbicides For Peanut.....

Directions for Grass Suppression for Harvest Efficiency in Peanut with TAPOUT .....

**Soybean**

Table 9. TAPOUT Tank Mix with Broadleaf Herbicides for the Control of Volunteer Corn (Including Roundup Ready) in Soybean.....

Table 10. TAPOUT Tank Mixes with Broadleaf Herbicides for Soybean.....

**Sugar Beet**

Table 11. TAPOUT Tank Mixed with Broadleaf Sugar Beet Herbicides.....

Table 12. TAPOUT Plus Betanex or Betamix Tank Mix for Three Sequential Applications for Annual Grass Control (Micro-Rate Application).....

Directions for Use for Micro-Rate Applications to Sugar Beet.....

Table 13. Tank Mix Application of TAPOUT and Fungicides for Control of Grass Weeds and Diseases in Sugar Beet.....

Table 14. Tank Mix Application of TAPOUT and Insecticides for Control of Grass Weeds and Insects in Alfalfa, Cotton, Mint, Peanut (including perennial), Soybean and Sunflower.....

**Fallow Land**

Directions for Use.....

Product Information.....

Table 15. TAPOUT in Tank Mixes to Control Annual and Perennial Grasses in Fallow

Land.....

Directions for Grass Suppression in Non-Crop Areas with TAPOUT.....

Table 16. TAPOUT for the Control and/or Suppression of Tall Fescue in Native Prairie Warm-Season Grass

**Restoration.....**

Projects

Special Application Instructions/Precautions.....

Table 17. TAPOUT for the Suppression of Tall Fescue Seed-Heads in Non-Producing Agricultural Areas.....

Special Application Instructions/Precautions.....

**Directions For Use in Ornamentals.....**

Ornamental Trees.....

Ground Covers.....

Garden Flowers and Plants.....

Shrubs.....

Directions for Annual Grasses in Ornamentals.....

Directions for Annual Bluegrass Control with TAPOUT in Ornamentals.....

Directions for Perennial Grasses in Ornamentals.....

Non-Bearing Food Crops.....

Conifer Trees.....

Non-Crop or Non-Planted Areas.....

**THE FOLLOWING STATEMENT ON CHEMIGATION WILL BE USED ONLY IF A SUPPLEMENTAL LABEL IS CREATED.**

**CHEMIGATION**

[Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed. Refer to supplemental labeling entitled, "Application of TAPOUT Onions (dry bulb and green) and Garlic by Chemigation", for use directions for chemigation.]

May be applied to onions and garlic by sprinkler irrigation systems. Do not apply by chemigation to any other crop, or to this crop using any other type of irrigation system.

**PRODUCT INFORMATION**

**FOR USE ON:** Alfalfa, Asparagus, Bean (dry) and Pea (shelled)<sup>1</sup>, Bean and Pea (succulent shelled)<sup>2</sup>, Broccoli, Cabbage, Canola\*, Carrot, Cauliflower (and other Head and Stem Brassica Vegetables)<sup>3</sup>, Celery, Clover (grown in Idaho, Oregon and Washington only), Conifers, Cotton, Cranberry, Cucumber, Eggplant (and other Fruiting Vegetables)<sup>4</sup>, Fallow Land (and other non-producing agricultural areas), Flax\*, Garden Beet, Garlic, Herbs<sup>5</sup>, Hops, Horseradish (and other Root Vegetables)<sup>6</sup>, Legume Vegetables (edible podded)<sup>7</sup>, Lettuce, Head and Leaf (and other Leafy Greens)<sup>8</sup>, Melons (including Cantaloupe and Watermelon)<sup>9</sup>, Mint, Mustard Greens (and other Leafy Brassica Greens)<sup>10</sup>, Mustard Seed\*, Non-Bearing Food Crops, Non-Crop or Non-Planted Areas, Onions (dry bulb and green), Ornamentals, Peanut (including perennial), Peppers (bell and non-bell), Potato, Radish, Rhubarb (and other Leaf Petioles)<sup>11</sup>, Safflower, Sesame, Shallot (dry bulb), Soybean, Squash (including Pumpkin)<sup>9</sup>, Strawberry, Sugar Beet, Sunflower, Sweet Potato, Turnip Greens, Tomato and Yam (and other Tuberous and Corm Vegetables)<sup>12</sup>.

**\*Not for use in California**

<sup>1</sup> Other Bean (dry) and Pea (shelled) crops approved for use with TAPOUT include: Bean (*Lupinus* spp.), grain lupin, sweet lupin, white lupin and white sweet lupin; Bean (*Phaseolus* spp.), field, kidney, lima (dry), navy, pinto and tepary; Bean (*Vigna* spp.), adzuki bean, black-eyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, broad (dry), chickpea (garbanzo), guar, lablab bean and lentil; Pea (*Pisum* spp.), field and pigeon.

<sup>2</sup> Other Bean & Pea (succulent shelled) crops approved for use with TAPOUT include: Bean (*Phaseolus* spp.), broad bean (succulent), lima bean (green); Bean (*Vigna* spp.), black-eyed pea, cowpea, Southern pea; Pea (*Pisum* spp.), English pea, garden pea, green pea and pigeon pea.

<sup>3</sup> Other head and stem brassica vegetables approved for use with TAPOUT include: Chinese broccoli, Brussels sprouts, Chinese (napa) cabbage, Chinese mustard, cavalo broccolo and kohlrabi.

<sup>4</sup> Other Fruiting Vegetables (except tomato) approved for use with TAPOUT include: eggplant, groundcherry, pepino, peppers (all) and tomatillo.

<sup>5</sup> Other Herb crops approved for use with TAPOUT include: angelica, balm, basil, borage, burnet, camomile, catnip, chervil (dried), chive, Chinese chive, clary, coriander (leaf), costmary, culantro (leaf), curry (leaf), dill (dillweed), horehound, hyssop, lavender, lovage (leaf), marigold, maqoram (*Origanum* spp.), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage and savory, summer and winter.

<sup>6</sup> Other root vegetables approved for use with TAPOUT include: burdock, edible; celeriac; chervil; turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip.

<sup>7</sup> Other Edible Podded Legume Vegetable crops approved for use with TAPOUT include: Bean (*Phaseolus* spp.), runner, snap and wax; Bean (*Vigna* spp.), asparagus, chinese longbean, moth, yardlong, jackbean; Pea (*Pisum* spp.), dwarf, edible-pod, snow, sugar snap, pigeon and sword bean.

<sup>8</sup> Other Leafy Greens crops approved for use with TAPOUT include: amaranth (Chinese spinach, leafy amaranth and tampala), arugula (roquette), chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden, yellow rock and winter), dandelion, dock (sorrel), endive (escarole), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio (red chicory), spinach, spinach (New Zealand and Vine (Indian and Malabar)).

<sup>9</sup> Other cucurbit crops approved for use with TAPOUT include: Chayote (fruit), Chinese Wax Gourd, Citron Melon, Edible Gourd, Ghorkin and Muskmelons (all) including Honeydew Melon.

<sup>10</sup> Other leafy brassica greens approved for use with TAPOUT include: broccoli raab, Chinese (bok choy) cabbage, collards, kale, mizuna, mustard greens, mustard spinach, rape greens.

<sup>11</sup> Other leaf petiole crops approved for use with TAPOUT include: cardoon, celtuce, Chinese celery, Florence fennel, and Swiss chard.

<sup>12</sup> Other tuber and corm vegetables approved for use with TAPOUT include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, bitter and sweet cassava, chayote (root), chufa, dasheen (taro), ginger, leren, tanager, turmeric; yam, bean and



9/145

yam, true.

**TAPOUT** is a selective postemergence herbicide for control of annual and perennial grasses. **TAPOUT** does not control sedges or broadleaf weeds.

**Control Symptoms**

Treated grass weeds show a reduction in vigor and growth. Early chlorosis/necrosis of younger plant tissue is followed by a progressive collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days after application, depending on grass species treated and environmental conditions.

**APPLICATION INFORMATION**

**Timing of Applications**

Apply **TAPOUT** postemergence to actively growing grasses according to rate table recommendations. Applications made to grass plants stressed by insufficient moisture, hot or cold temperatures, or to grass plants exceeding specified growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, **TAPOUT** should be applied, as soon as possible, after irrigation (within 7 days). In arid regions, a second application of **TAPOUT** will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of **TAPOUT** may reduce weed control.

**Ground Application**

Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 5 gals. and a maximum of 40 gals of spray solution per acre. Under the following conditions a minimum of 10 gals. per acre is required: ultra narrow row cotton, narrow row soybeans, broadleaf herbicide tank mixes, perennial grasses, volunteer corn, drought or stress conditions, heavy grass pressure or when grasses are at or near maximum height. Failure to use a minimum of 10 gals per acre under these conditions can result in poor coverage and reduced grass control requiring repeat applications. Spray pressures should reflect a minimum of 30 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Applications to garlic, onions (dry bulb and green) or shallots (dry bulb and green) should be made in a minimum of 20 gals of spray solution per acre.

**Air Application**

Use a minimum of 3 gals of spray solution per acre unless otherwise directed in this label. Increase spray volumes up to 10 gals as grass or crop foliage becomes dense. For garlic, onions (dry bulbs and green) or shallots (dry bulb and green): When applying by air do not exceed 16 fl oz/A in a single application. In California, air applications to garlic, onion or shallot should be made in a minimum of 20 gals of spray solution per acre. In states other than California, air application to garlic, onion or shallot should be made in a minimum of 10 gals of spray solution.

**NOTE: Crop injury may occur when TAPOUT is applied to garlic, onion or shallot with aerial equipment.**

**Spot Treatment**

When using hand sprayers or high volume sprayers utilizing hand guns, mix 1/3 to 2/3% (0.44 oz. to 0.85 oz. per gal.) **TAPOUT** and treat to wet vegetation, while not allowing runoff of spray solution. For uses requiring crop oil concentrate; include crop oil concentrate at 1% (1.3 oz. per gal.) by volume. For uses requiring nonionic surfactant, include non-ionic surfactant at 1/4% (0.33 oz. per gal.) by volume.

**NOTE: If TAPOUT is applied as a spot treatment care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.**

**CHEMIGATION — GARLIC AND ONIONS (Dry Bulb and Green)  
SPRINKLER IRRIGATION APPLICATION**

**Do not apply TAPOUT by chemigation in the states of Idaho, Montana, Oregon and Washington.**

Apply **TAPOUT** at the high rate specified for annual grasses (32 fl. oz./A) when the grass height is at the high end of the range (application to larger grasses may not provide adequate control). Add a crop oil concentrate containing at least 15% emulsifier at 1 quart per acre or non-ionic surfactant with at least 80% active ingredient at 0.25% v/v of total spray solution.

Apply **TAPOUT** in 0.1 to 0.2 acre inch of water either at the end of a regular irrigation set or as a separate application not associated with a regular irrigation using the least amount of water that provides proper distribution and coverage. Application of more than label specified quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness. Use a metering device to inject the **TAPOUT** into the irrigation water at a constant flow. Constant agitation must be maintained in the chemical supply tank during the entire period of herbicide application. Inject the product with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period.

It is not recommended that **TAPOUT** be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

**Use Precautions**

1. Apply this product only through irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set or hand move. Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
3. If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.
5. A person knowledgeable of chemigation system and responsible for its operation or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

**RESTRICTIONS AND LIMITATIONS**

**ALL USES**

- Do not apply if rain is expected within 1 hour of application, as control may be unsatisfactory.
- Do not plant rotational crops until 30 days after application of **TAPOUT** unless crop is listed on **TAPOUT** label.
- Do not apply a postemergence broadleaf herbicide within one day following application of **TAPOUT** or reduced grass control may result.
- **TAPOUT** is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.

11/45

- Do not apply under conditions of stress. Applying **TAPOUT** under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate **TAPOUT** effectively, and will be less susceptible to herbicide activity.
- Application on Long Island, New York, is restricted to no more than 32 fl oz of **TAPOUT** (0.25 lb a.i) per acre per season.

Optimal perennial grass control can be obtained if rhizomes or stolons are cut up by preplant tillage practices (disking, plowing, etc.) to stimulate maximum emergence of grass shoots. Cultural practices, such as continuous no-tillage in which the perennial grass rhizomes or stolons are not cut up, results in a very staggered, non-uniform weed emergence. Due to this non-uniform weed emergence, no fewer than two **TAPOUT** applications per season per year are recommended at the appropriate weed-growth stage rate under continuous no-till conditions.

Grass crops such as corn, rice, small grains, sorghum or turf, etc. are highly sensitive to **TAPOUT**.

While all the vegetable crops on this label have been tested and are tolerant to **TAPOUT**, not all specialty varieties of these crops have been tested. It is advised that, before applying **TAPOUT** to specialty varieties of vegetable crops on this label, crop tolerance be investigated first using a small section of the field. It is possible that injury symptoms can occur. Symptoms may appear as leaf speckling or stunting.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotation and other crop restrictions.

Tank mixes of **TAPOUT** and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs, an additional application of **TAPOUT** may be necessary.

**SPRAY DRIFT MANAGEMENT**

- Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:
- Do not apply when conditions are favorable for drift (high temperatures, drought and low relative humidity), especially when sensitive plants are located nearby.
- Do not spray if wind speed is 10 mph or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions.
- Do not spray if winds are gusty.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Do not allow **TAPOUT** to come in contact with desirable grass crops such as corn, rice, small grains, sorghum, or turf, as these and other grass crops will be injured or killed.

Further reductions in drift can be obtained by:

1. Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
2. Orienting nozzles straight back with the windstream, using straight stream orifices for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
3. Increasing the volume of spray mixture (for example a minimum of 10 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
4. Applying as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

**Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

**RESISTANCE MANAGEMENT**

**TAPOUT** is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to **TAPOUT** and other Group 1 herbicides. Weed species with acquired resistance to Group 1 may eventually dominate the weed population if Group 1 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **TAPOUT** or other Group 1 herbicides. Repeated use of **TAPOUT** (or similar postemergence grass herbicide with the same mode of action) may lead to the selection of naturally occurring biotypes that are resistant to these products in some grass species.

If poor performance occurs and cannot be attributed to adverse weather or application conditions, a resistant biotype may be present. This is most likely to occur in fields where other control strategies such as crop rotation, mechanical removal and other classes of herbicides are not used from year to year.

To delay herbicide resistance consider:

- Avoiding the consecutive use of **TAPOUT** or other target site of action Group I herbicides that have similar target site of action, on the same weed species.
- Using tank mixtures or premixes with herbicides from different target site of action groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

**Table 1. CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT**

| Crop <sup>(1)</sup>   | Minimum Time From Application to Harvest (PHI)                            | Annual Grass Use Rate Per Acre <sup>(2)</sup> | Perennial Grass Use Rate Per Acre <sup>(2)</sup> | Adjuvant Recommendation <sup>(3)</sup><br><small>(Use a CPDA Certified adjuvant when possible)</small>   | Ammonium Sulfate Recommendation <sup>(4)</sup> | Special Use Instructions And Restrictions   |
|---|---|---|--|--|--|---|
| <b>Alfalfa, Seedling</b>  | 15 days before grazing, feeding or harvesting (cutting) for forage or hay | 9 to 16 fl oz                                 | 12 to 32 fl oz                                   | NIS at 0.25% v/v or COC/MSO at 1 qt/A<br>Or<br>1% v/v<br><br>See tank mix label for specific adjuvant recommendations,   | 2.5 to 4 lbs/A                                 | Do not apply more than 32 fl oz/A per application.<br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br>For repeat applications make on a minimum of a 14 day interval.         |
| <b>Alfalfa, Established</b> including: Sainfoin, Holy clover, Birdsfoot trefoil | 15 days before grazing, feeding or harvesting (cutting) for forage or hay | 12 to 16 fl oz                                | 12 to 32 fl oz                                   | NIS at 0.25% v/v or COC/MSO at 1 qt/A<br>1% v/v<br><br>See tank mix label for specific adjuvant recommendations.<br><br><small>(Use a CPDA Certified adjuvant when possible)</small> | 2.5 to 4 lbs/A                                 | Do not apply more than 32 fl oz/A per application.<br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br>For repeat applications make on a minimum of a 14 day interval.         |
| <b>Asparagus</b>  | 1 day   | 9 to 16 fl oz                                 | 12 to 16 fl oz                                   | NIS at 0.25% v/v<br><small>(Use a CPDA Certified adjuvant when possible)</small>   | None   | Do not apply more than 16 fl oz/A in a single application,<br>For repeat applications make on a minimum of a 14 day interval.<br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season. |

|  |                |                                    |                       |  |             |  |
|--|----------------|------------------------------------|-----------------------|--|-------------|--|
| <p><b>Bean, Dry including:</b><br/>         Bean (<i>Lupinus</i> spp.)<br/>         Grain Sweet White<br/>         White Sweet Bean (<i>Phaseolus</i> spp.)<br/>         Field Kidney Lima (dry) Navy Pinto Tepary<br/>         Bean (<i>Vigna</i> spp.)<br/>         Adzuki Bean Black-eyed Pea Catjang Cowpea Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea</p> | <p>30 days</p> | <p>9 to 16 fl oz<sup>(5)</sup></p> | <p>12 to 32 fl oz</p> | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p> | <p>None</p> | <p>Do not apply more than 32 fl oz/A per application.<br/><br/>         For repeat applications make on a minimum of a 14 day interval.<br/><br/>         Do not apply more than 64 fl. oz/A (0.5 lb al/A) per season.</p>         |
| <p>Urd Bean Broad (dry) Chickpea (garbanzo) Guar Lablab Bean Lentil</p>  |                |                                    |                       |  |             |  |
| <p><b>Bean, Succulent Shelled including:</b><br/>         Bean (<i>Phaseolus</i> spp.)<br/>         Broad Bean (succulent) Lima Bean (green) Bean (<i>Vigna</i> spp.)<br/>         Black-eyed Pea Cowpea Southern Pea</p>  | <p>21 days</p> | <p>9 to 16 fl oz<sup>(5)</sup></p> | <p>12 to 16 fl oz</p> | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p> | <p>None</p> | <p>Do not apply more than one (1) application per acre per season.</p>   |
| <p>Beet, Garden</p>  | <p>30 days</p> | <p>9 to 16 fl oz</p>               | <p>12 to 16 fl oz</p> | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p> | <p>None</p> | <p>Do not apply more than 16 fl. oz/A in a single application.<br/><br/>         Do not apply more than 64 fl oz/A (0.5 lb al/A) per season.<br/><br/>         For repeat applications make on a minimum of a 14 day interval.</p> |

|  |  |                              |                |  |                |  |
|--|--|------------------------------|----------------|--|----------------|--|
| <b>Brassica Vegetables, Head and Stem</b><br>(6)<br>including:<br>Broccoli<br>Cabbage<br>Cauliflower<br>Brussels sprouts | 30 days  | 9 to 16 fl oz                | 12 to 16 fl oz | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)  | None           | Do not apply more than 16 fl oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval   |
| <b>Canola</b><br>(Not for use in California)   | 70 days  | 9 to 12 fl oz <sup>(5)</sup> | 12 ft. oz.     | NIS at 0.25% v/v<br>Use a CPDA Certified adjuvant when possible)   | None           | Do not apply after crop has begun bolting. Crop injury may occur when <b>TAPOUT</b> is applied during the bloom period.<br><br>Do not apply more than 12 fl oz/A in a single application.<br><br>Do not apply more than 12 fl oz/A per season.                                       |
| <b>Carrot</b>  | 30 days  | 9 to 16 fl oz                | 12 to 16 fl oz | NIS at 0.25% v/v<br>Use a CPDA Certified adjuvant when possible)   | None           | Do not apply more than 16 fl oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.  |
| <b>Clover</b>  | 15 days<br>before grazing, feeding or harvesting (cutting) for forage or hay | 9 to 16 fl oz                | 12 to 32 fl oz | NIS at 0.25% v/v<br>Use a CPDA Certified adjuvant when possible)   | 2.5 to 4 lbs/A | For use on clover grown in the states of Idaho, Oregon and Washington only.<br><br>Do not more than 32 fl oz in a season.<br><br>For repeat applications make on a minimum of a 14 day interval.   |
| <b>Cotton</b>  | 60 days  | 9 to 16 fl oz                | 12 to 32 fl oz | NIS at 0.25% v/v or COC/MSO at 1 qt/A or 1% v/v<br>See tank mix label for specific adjuvant recommendations<br>(Use a CPDA Certified adjuvant when possible) | 2.5 to 4 lbs/A | Do not graze treated fields or feed treated forage or hay to livestock.<br><br>Do not apply more than 32 fl.oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval. |
| <b>Cranberry</b>   | 30 days  | 9 to 16 fl oz                | 12 to 16 fl oz | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)  | None           | Do not apply more than 16 fl oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a.i/A) per season.<br><br>Do not apply between the "hook" stage and full fruit set.<br><br>For repeat applications make on a minimum of a 14 day interval.               |

|  |         |                                |                  |   |                 |   |
|--|---------|--------------------------------|------------------|---|-----------------|---|
| <b>Cucurbits</b><br>including:<br>Cantaloupes (all)<br>Cucumber<br>Gherkin<br>Honeydew<br>Melon<br>Muskmelons (all)<br>Pumpkin<br>Squash (all)<br>Watermelon | 14 days | 9 to 16 fl oz                  | 12 to 16 fl oz   | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Do not apply more than 16 fl oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb. a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.  |
| <b>Fallow Land</b><br><b>Conifer Trees</b><br>(and other non-producing agricultural areas)<br><br><b>Non-Crop or Non-Planted Areas</b>                       | N/A     | 9 to 16 fl oz.                 | 12 to 32 fl oz   | NIS at 0.25% v/v<br>or<br>COC/MSO at 1 qt./A<br>or<br>1% v/v<br>(Use a CPDA Certified adjuvant when possible) | 2.5 to 4 lbs./A | Do not plant any crop for 30 days after application unless Clethodim is registered for use in that crop.<br><br>Do not apply more than 32 fl oz/A per application.<br><br>Do not apply more than 64 fl. oz/A (0.5 lb. a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval. |
| <b>Flax</b><br>(Not for use in California)   | 60 days | 9 to 16 fl. oz. <sup>(5)</sup> | 12 to 16 fl oz.  | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Apply prior to bloom. Crop injury may occur when TAPOUT is applied during the bloom period.<br><br>Do not apply more than 16 fl oz./A per application.<br><br>Do not apply more than 32 fl. oz. in a season.<br><br>For repeat applications make on a minimum of a 14 day interval.                             |
| <b>Fruiting Vegetable</b><br>(except Tomato)<br>including:<br>Eggplant<br>Groundcherry<br>Pepino<br>Peppers (all)<br>Tomatillo                               | 20 days | 9 to 16 fl. oz.                | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Do not apply more than 16 fl. oz/A in a single application.<br><br>Do not apply more than 64 fl oz./A (0.5 lb. a.i/A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.  |

16/45

|  |                |                                    |   |   |             |  |
|--|----------------|------------------------------------|---|---|-------------|--|
| <p><b>Herbs</b><br/>including:<br/>Angelica<br/>Balm<br/>Basil<br/>Borage<br/>Burnet<br/>Camomile<br/>Catnip<br/>Chervil (dried)<br/>Chive<br/>Chive, Chinese<br/>Clary<br/>Coriander (leaf)<br/>Costmary<br/>Culantro (leaf)<br/>Curry (leaf)<br/>Dill (dillweed)<br/>Horehound<br/>Hyssop<br/>Lavender<br/>Lovage (leaf)<br/>Marigold<br/>Marjoram<br/>(<i>Origanum</i> spp.)<br/>Nasturtium<br/>Parsley (dried)<br/>Pennyroyal<br/>Rosemary<br/>Rue<br/>Sage<br/>Savory, Summer<br/>and Winter<br/>Sweet Bay<br/>Tansy<br/>Tarragon<br/>Thyme<br/>Wintergreen<br/>Woodruff<br/>Wormwood</p> | <p>14 days</p> | <p>9 to<br/>16<br/>fl<br/>oz.</p>  | <p>12<br/>to<br/>16<br/>fl<br/>oz.</p>  | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p> | <p>None</p> | <p><b>TAPOUT</b> has not been tested on all herbs, and herb varieties. It is the responsibility of the user to test <b>TAPOUT</b> on a small portion of the crop to be treated before treating the entire field.</p> <p>Crop tolerance should be verified to <b>TAPOUT</b> on a small area of the herb crop, at the desired <b>TAPOUT</b> rate and with the same crop oil concentrate that will be used on the herb field. If no crop response is evident seven (7) days after treatment, <b>TAPOUT</b> may be used on the entire field at the rate tested and with the same crop oil used in the tolerance test.</p> <p>Do not apply more than 16 fl. oz/A in a single application.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Do not apply more than 64 fl. oz./A (0.5 lb. a.i/A) per season.</p> |
| <p><b>Hops</b></p>   | <p>21 days</p> | <p>9 to<br/>16<br/>fl<br/>oz.</p>  | <p>12<br/>to<br/>16<br/>fl<br/>oz.</p>  | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p> | <p>None</p> | <p>Do not apply more than 16 fl. oz/A in a single application.</p> <p>For repeat application make on a minimum of a 14 day interval.</p> <p>Do not apply more than 64 fl. oz/A (0.5 lb. a.i/A) per season.</p>   |
| <p><b>Leaf Lettuce</b></p>   | <p>14 days</p> | <p>9 to<br/>16<br/>fl.<br/>oz.</p> | <p>12<br/>to<br/>16<br/>fl.<br/>oz.</p> | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p> | <p>None</p> | <p>Do not apply more than 16 fl. oz/A in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.5 lb. al/A) per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p>  |



|  |         |                 |                  |   |      |   |
|--|---------|-----------------|------------------|---|------|---|
| <b>Leafy Brassica Greens,</b><br>including:<br>Broccoli Raab<br>Cabbage,<br>Chinese (bok choy)<br>Collards<br>Kale.<br>Mizuna<br>Mustard Greens<br>Mustard Spinach<br>Rape Greens  | 14 days | 9 to 16 fl oz   | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible) | None | Do not apply more than 16 fl oz./A in a single application.<br><br>Do not apply more than 64 fl. oz./A (0.5 lb. a.i/A) per season.<br><br>For repeat application make on a minimum of a 14 day interval.  |
| <b>Leaf Petioles</b><br>including:<br>Cardoon<br>Celery<br>Celtuce<br>Chinese Celery<br>Fennel, Florence (finocchio)<br>Rhubarb<br>Swiss Chard   | 30 days | 9 to 16 fl. oz. | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible) | None | Do not apply more than 16 fl. oz/A in a single application.<br><br>For repeat applications make on a minimum of a 14 day interval.<br><br>Do not apply more than 64 fl. oz./A (0.5 lb. a.i/A) per season. |
| <b>Leafy Greens</b><br>including:<br>Amaranth<br>Chinese<br>Spinach  | 14 days | 9 to 16 fl oz.  | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible) | None | Do not apply more than 16 fl. oz./A in a single application.<br><br>For repeat applications make on a minimum of a 14 day interval.   |
| Leafy Amaranth<br>Tampala<br>Arugula (roquette)<br>Chervil<br>Chrysanthemum, Edible-leaved<br>Chrysanthemum, Garland<br>Corn Salad<br>Cress<br>Garden<br>Upland (yellow rock and winter)<br>Dandelion<br>Dock (sorrel)<br>Endive (escarole)<br>Lettuce, Head and Leaf<br>Orach<br>Parsley<br>Purslane<br>Garden<br>Winter<br>Radicchio (red chicory)<br>Spinach<br>Spinach<br>New Zealand<br>Vine (Indian and Malabar) |         |                 |                  |   |      | Do not apply more than 64 fl. oz./A (0.5 lb. a.i/A) per season.   |

|  |                |                                      |                         |   |                        |   |
|--|----------------|--------------------------------------|-------------------------|---|------------------------|---|
| <p><b>Legume Vegetables, Edible Podded</b> including:<br/>         Bean (<i>Phaseolus</i> spp.)<br/>         Runner<br/>         Snap<br/>         Wax<br/>         Bean (<i>Vigna</i> spp.)<br/>         Asparagus<br/>         Chinese<br/>         Longbean<br/>         Moth<br/>         Yardlong<br/>         Jackbean<br/>         Pea (<i>Pisum</i> spp.)<br/>         Dwarf<br/>         Edible-pod<br/>         Snow<br/>         Sugar Snap<br/>         Pigeon<br/>         Sword Bean</p> | <p>21 days</p> | <p>9 to 16 fl oz<sup>(5)</sup></p>   | <p>12 to 16 fl oz.</p>  | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p>                                | <p>None</p>            | <p>Do not apply more than 16 fl oz/A in a single application.</p> <p>Do not apply more than one (1) application per acre per season.</p> <p>For peas apply before bloom, but no later than 21 days before harvest.</p>  |
| <p><b>Mint</b></p>   | <p>21 days</p> | <p>9 to 16 fl. oz.</p>               | <p>12 to 32 fl. oz.</p> | <p>NIS at 0.25% v/v or COC/MSO at 1 qt./A or 1%v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p> | <p>2.5 to 4 lbs./A</p> | <p>Do not apply more than 32 fl. oz./A in a single application.</p> <p>Do not apply more than 64 fl. oz./A (0.5 lb. al/A) per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p>  |
| <p><b>Mustard Seed</b><br/>         (Not for use in California)</p>  | <p>75 days</p> | <p>9 to 12 fl. oz.<sup>(5)</sup></p> | <p>12 fl. oz.</p>       | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p>                                | <p>None</p>            | <p>Do not apply after crop has begun bolting. Crop injury may occur when TAPOUT is applied during the bloom period.</p> <p>Do not apply more than 12 fl. oz./A per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p>   |
| <p><b>Onions (Dry Bulbs Only)<sup>(7,8)</sup></b><br/> <b>Garlic<sup>(7,8)</sup></b><br/> <b>Shallots (Dry Bulbs Only)<sup>(7,8)</sup></b></p>   | <p>45 days</p> | <p>9 to 16 fl oz.</p>                | <p>12 to 32 fl. oz.</p> | <p>NIS at 0.25% v/v<br/>         (Use a CPDA Certified adjuvant when possible)</p>                                | <p>None</p>            | <p>Do not apply more than 32 fl. oz/A per application.</p> <p>Do not apply more than 64 fl. oz./A (0.5 lb. a.i/A) per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Minimum of 20 gals/A spray volume by ground in entire U.S.</p> <p>Minimum of 20 gals./A spray volume by air in California.</p> <p>In states other than California, air applications to onions, garlic or shallots should be made in a minimum of 10 gals/A.</p> |

|  |                       |   |   |  |                           |   |
|--|-----------------------|---|---|--|---------------------------|---|
| <p><b>Onions, Green</b><br/>(7,8)<br/>including:<br/>Leeks<br/>Scallions or<br/>Spring Onions<br/>Japanese<br/>Bunching<br/>Onions<br/>Green Shallots<br/>Green Eschalots</p> <p>(Not for use in<br/>California)</p> | <p>14 days</p>        | <p>9 to<br/>16<br/>fl oz</p>                                  | <p>12 to<br/>16<br/>fl oz</p>                           | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p>  | <p>None</p>               | <p>Do not apply more than 16 fl. oz./A in a single application.</p> <p>Do not apply more than 64 fl oz./A (0.5 lb. a.i/A) per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Air applications to onions, garlic or shallots should be made in a minimum of 10 gals/A. (7)</p>            |
| <p><b>Ornamentals</b></p> <p><b>Non-Bearing Food Crops</b><sup>(7)</sup></p>   | <p>N/A</p> <p>N/A</p> | <p>9 to<br/>16<br/>fl. oz.</p> <p>9 to<br/>16<br/>fl. oz.</p> | <p>12 to<br/>32 fl. oz.</p> <p>12 to<br/>16 fl. oz.</p> | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p>  | <p>None</p>               | <p>Do not apply more than 64 fl. oz./A (0.5 lb. al/A) per season.</p> <p>For repeat applications make on a minimum of a 14 day interval.</p> <p>Sugar maples cannot be tapped for syrup within one year of TAPOUT application.</p> <p>Do not apply more than 16 fl oz./A in a single application to non-bearing food crops.</p> |
| <p><b>Pea, Shelled</b><br/>(<i>Pisum</i> spp.)<br/>Field<br/>Pigeon</p>  | <p>30 days</p>        | <p>9 to<br/>16<br/>fl oz<br/>(5)</p>                          | <p>12 to<br/>16<br/>fl oz.</p>                          | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p>  | <p>None</p>               | <p>Do not apply more than 16 fl. oz./A per application.</p> <p>Do not apply more than one (1) application per acre per season.</p> <p>Apply before bloom but not later than 30 days prior to harvest. (9)</p>   |
| <p><b>Pea, Succulent Shelled</b><br/>(<i>Pisum</i> spp.)<br/>English Pea<br/>Garden Pea<br/>Green Pea<br/>Pigeon Pea</p>   | <p>21 days</p>        | <p>9 to<br/>16<br/>fl oz<sup>(5)</sup></p>                    | <p>12 to<br/>16 fl oz</p>                               | <p>NIS at 0.25% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p>  | <p>None</p>               | <p>Do not apply more than 16 fl oz./A per application.</p> <p>Do not apply more than one (1) application per acre per season.</p> <p>Apply before bloom but not later than 21 days prior to harvest. (9)</p>  |
| <p><b>Peanut</b><br/>(including perennial)</p>   | <p>40days</p>         | <p>9 to<br/>16<br/>fl oz</p>                                  | <p>12 to<br/>32 fl oz</p>                               | <p>NIS at 0.25% v/v<br/>or<br/>COC/MSO at 1 qt/A<br/>or 1% v/v<br/>(Use a CPDA Certified adjuvant<br/>when possible)</p> | <p>2.5 to<br/>4 lbs/A</p> | <p>Do not apply more than 32 V fl oz/A in a single application.</p> <p>Do not apply more than 64 fl oz/A (0.5 lb al/A) per season, except in Florida where the limit is 64 fl oz/A per year (perennial peanuts only).</p> <p>For repeat applications make on a minimum of a 14 day interval.</p>                                |

|   |         |                              |                  |   |                 |   |
|---|---------|------------------------------|------------------|---|-----------------|---|
| <b>Potato</b>   | 30 days | 9 to 16 fl. oz.              | 12 to 32 fl. oz. | NIS at 0.25% v/v or COC/MSO at 1 qt./A or 1%v/v<br><br>(Use a CPDA Certified adjuvant when possible)  | 2.5 to 4 lbs./A | Do not apply more than 32 fl. oz./A in a single application.<br><br>Do not apply more than 64 fl. oz./A (0.5 lb. a.i./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.   |
| <b>Radish</b>   | 15 days | 9 to 16 fl. oz.              | 12 to 16 fl. oz. | NIS at 0.25%v/v<br>(Use a CPDA Certified adjuvant when possible)  | None            | Do not apply more than 16 fl. oz./A in a single application.<br><br>Do not apply more than 32 fl. oz. (0.25 lb. a.i) per acre in a season.<br><br>For repeat applications V make on a minimum of a 14 day interval.   |
| <b>Root Vegetables (except radish)<sup>(10)</sup></b> | 30 days | 9 to 16 fl. oz.              | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Do not apply more than 16 fl. oz./A in a single application.<br><br>Do not apply more than 64 fl. oz./A (0.5 lb. aI/A) per V season.<br><br>For repeat applications make on a minimum of a 14 day interval.   |
| <b>Safflower</b>                                      | 70 days | 9 to 16 fl. oz.              | 12-16 fl. oz.    | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Do not apply more than 16 fl oz/A in a single application.<br><br>For repeat applications make on a minimum of a 14 day interval.<br><br>Do not apply more than 64 fl oz/A (0.5 lb aI/A) per season.  |
| <b>Sesame</b>   | 14 days | 9 to 16 fl oz                | 12 to 16 fl oz   | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None            | Do not apply during flowering.<br><br>Do not apply more than 16 fl oz/A in a single application.<br><br>For repeat applications make on a minimum of a 14 day interval.<br><br>Do not apply more than 64 fl oz/A (0.5 lb aI/A) per season.                                    |
| <b>Soybean</b>  | 60 days | 9 to 16 fl oz <sup>(5)</sup> | 12 to 32 fl oz   | NIS at 0.25% v/v or COC/MSO at 1 qt/A or 1% v/v<br><br>See tank mix label for specific adjuvant recommendations.<br>(Use a CPDA Certified adjuvant when possible) | 2.5 to 4 lbs/A  | Do not apply more than 32 fl oz/A per application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a.i./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.<br><br>Do not graze treated fields or feed treated forage or hay to livestock. |

|  |         |                              |                  |   |                |  |
|--|---------|------------------------------|------------------|---|----------------|--|
| <b>Strawberry</b>  | 4 days  | 9 to 16 fl oz                | 12 to 16 fl oz   | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None           | Do not apply more than 16 fl oz/A in a single application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval. |
| <b>Sugar Beet</b>  | 40 days | 9 to 16 fl oz <sup>(5)</sup> | 12 to 32 fl oz   | NIS at 0.25% v/v or<br>COC/MO at 1 qt/A or 1% v/v<br><br>See tank mix label for specific adjuvant recommendations.<br>(Use a CPDA Certified adjuvant when possible) | 2.5 to 4 lbs/A | Do not apply more than 32 fl oz/A per application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.         |
| <b>Sunflower</b>   | 70 days | 9 to 16 fl oz <sup>(5)</sup> | 12 to 32 fl oz   | NIS at 0.25% v/v or<br>COC/MO at 1 qt/A or 1% v/v<br><br>See tank mix label for specific adjuvant recommendations.<br>(Use a CPDA Certified adjuvant when possible) | 2.5 to 4 lbs/A | Do not apply more than 32 fl. oz./A per application.<br><br>Do not apply more than 64 fl oz/A (0.5 lb a./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.       |
| <b>Sweet Potato, Yam and other tuberous and corm vegetables (except potato)<sup>(12)</sup></b> | 30 days | 9 to 16 fl oz                | 12 to 32 fl oz   | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None           | Do not apply more than 32 fl oz/A per application.<br>Do not apply more than 64 fl oz/A (0.5 lb a./A) per season.<br>For repeat applications make on a minimum of a 14 day interval.                 |
| <b>Tomato</b>  | 20 days | 9 to 16 fl oz                | 12 to 32 fl oz   | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None<br>V      | Do not apply more than 32 fl oz/A per application.<br><br>Do not apply more than 64 fl. oz. /A (0.5 lb. a./A) per season.<br><br>For repeat applications make on a minimum of a 14 day interval.     |
| <b>Turnip Greens</b>   | 14 days | 9 to 16 fl oz                | 12 to 16 fl. oz. | NIS at 0.25% v/v<br>(Use a CPDA Certified adjuvant when possible)   | None           | Do not apply more than 16 fl. oz. /A in a single application.<br>Do not apply more than 64 fl. oz. /A (0.5 lb. a./A) per season.<br>For repeat application make on a minimum of a 14 day interval.   |

N/A = Not Applicable

1. **TAPOUT** is not specified for use on vegetable crops being grown for seed production unless specific use directions are provided.
2. See annual and perennial grass control tables for specific use rate recommendations.
3. Non-ionic surfactant (NIS) in this case refers to an adjuvant containing at least 80% non-ionic surfactant. Crop oil concentrate in this case refers to both crop oil concentrate and crop oil concentrate blends. Acceptable crop oil concentrates would be those that contain a minimum of 80% oils and 15% emulsifier. Acceptable crop oil concentrate blends would be those that contain a minimum of 60% oils and 25-40% surfactants and emulsifiers. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following

criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. **Use a CPDA Certified Adjuvant when possible.**

4. If Ammonium Sulfate is going to be used it must be spray grade ammonium sulfate. The use of ammonium sulfate or Ammonium Sulfate Replacement products does not take the place of the required adjuvant.
5. See DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.
6. Other head and stem brassica vegetables approved include: Chinese broccoli; Brussels sprouts; Chinese (napa) cabbage; Chinese mustard; cavalo broccoli; and kohlrabi.
7. If TAPOUT is applied as a spot treatment to garlic, onion, shallot or non-bearing food crops care should be taken to not exceed the maximum rate allowed on a "per acre" basis or crop injury may occur.
8. In California, do not apply TAPOUT to garlic, onions or shallots until crop has at least two full leaves. In California, 14 days spray intervals are recommended between the application of TAPOUT and liquid nitrogen and other herbicide applications. Injury to crop may occur when shorter intervals are observed.
9. Applications of TAPOUT to peas during the bloom period could result in severe crop injury, including loss of yield and delayed maturity.
10. Other root vegetables approved for use with TAPOUT include: burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; parsley, turnip-rooted; parsnip; radish, oriental; rutabaga; salsify; salsify, black; salsify, Spanish; skirret and turnip.
11. See TAPOUT TANK MIX WITH BROADLEAF HERBICIDES FOR THE CONTROL OF VOLUNTEER CORN (INCLUDING ROUNDUP READY®) IN SOYBEAN.
12. Other tuber and corm vegetables approved for use with TAPOUT include: arracacha, arrowroot, Chinese artichoke, Jerusalem artichoke, edible canna, cassava, bitter and sweet chayote (root), chufa, dasheen (taro), ginger, leren, tariier, turmeric and bean yam.

**DIRECTIONS FOR ANNUAL GRASSES  
ALL CROPS**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not exceed the maximum per application rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT

| GRASS SPECIES          | SCIENTIFIC NAME                    | WEED HEIGHT<br>(inches) | APPLICATION RATES         |  |
|------------------------|------------------------------------|-------------------------|---------------------------|--|
|                        |                                    |                         | MINIMUM RATE<br>fl. oz./A | Maximum rate <sup>(1)</sup><br>Fl. oz./A |
| Barnyardgrass          | <i>Echinochloa crus-galli</i>      | 2 to 8                  | 9                         | 16                                       |
| Broadleaf Signalgrass  | <i>Brachiaria platyphylla</i>      | 2 to 6                  | 9                         | 16                                       |
| Brome                  |                                    |                         |                           |  |
| California             | <i>Bromus carinatus</i>            | 2 to 6                  | 9                         | 16                                       |
| Cheat                  | <i>Bromus secalinus</i>            | 2 to 6                  | 9                         | 16                                       |
| Downy                  | <i>Bromus tectorum</i>             | 2 to 6                  | 9                         | 16                                       |
| Ripgut                 | <i>Bromus diandrus</i>             | 2 to 6                  | 9                         | 16                                       |
| Canarygrass            | <i>Phalaris canariensis</i>        | 1 to 4                  | 9                         | 16                                       |
| Crabgrass              |                                    |                         |                           |  |
| Hairy                  | <i>Digitaria adscendens</i>        | 2 to 6**                | 9                         | 16                                       |
| Large                  | <i>Digitaria sanguinalis</i>       | 2 to 6**                | 9                         | 16                                       |
| Smooth                 | <i>Digitaria ischaemum</i>         | 2 to 6**                | 9                         | 16                                       |
| Southern               | <i>Digitaria ciliaris</i>          | 2 to 6**                | 9                         | 16                                       |
| Crowfootgrass          | <i>Dactyloctenium aegyptium</i>    | 2 to 6**                | 9                         | 16                                       |
| Fall Panicum           | <i>Panicum dichotomiflor</i>       | 2 to 8                  | 9                         | 16                                       |
| Field Sandbur          | <i>Cenchrus incertus</i>           | 2 to 6                  | 9                         | 16                                       |
| Foxtail                |                                    |                         |                           |  |
| Giant                  | <i>Setaria faberi</i>              | 2 to 12                 | 9                         | 16                                       |
| Green                  | <i>Setaria viridis</i>             | 2 to 8                  | 9                         | 16                                       |
| Yellow                 | <i>Setaria glauca</i>              | 2 to 8                  | 9                         | 16                                       |
| Goosegrass             | <i>Eleusine indica</i>             | 2 to 6**                | 9                         | 16                                       |
| Itchgrass              | <i>Rottboellia cochinchinensis</i> | 2 to 6                  | 9                         | 16                                       |
| Junglerice             | <i>Echinochloa colona</i>          | 2 to 6                  | 9                         | 16                                       |
| Lovegrass (Stinkgrass) | <i>Eragrostis cilianensis</i>      | 2 to 6                  | 9                         | 16                                       |
| Rabbitsfootgrass       | <i>Polyogon monspeliensis</i>      | 1 to 4                  | 9                         | 16                                       |
| Red Rice               | <i>Oryza sativa</i>                | 1 to 3                  | 9                         | 16                                       |
| Rygrass                |                                    |                         |                           |  |
| Hardy                  | <i>Lolium remotum</i>              | 2 to 6                  | 9                         | 16                                       |
| Italian                | <i>Lolium multiflorum</i>          | 2 to 6                  | 9                         | 16                                       |

|                                  |                                |          |    |    |
|----------------------------------|--------------------------------|----------|----|----|
| Seedling Johnsongrass            | <i>Sorghum halepense</i>       | 4 to 10  | 9  | 16 |
| Shattercane                      | <i>Sorghum bicolor</i>         | 6 to 18  | 9  | 16 |
| Southwestern Cupgrass            | <i>Eriochloa gracilis</i>      | 2 to 6   | 9  | 16 |
| Sprangletop                      |                                |          |    |    |
| Amazon                           | <i>Leptochloa panicoides</i>   | 2 to 6   | 9  | 16 |
| Bearded                          | <i>Leptochloa fascicularis</i> | 2 to 6   | 9  | 16 |
| Mexican                          | <i>Leptochloa uninervia</i>    | 2 to 6   | 9  | 16 |
| Red                              | <i>Leptochloa filiformis</i>   | 2 to 6   | 9  | 16 |
| Texas Panicum                    | <i>Panicum texanum</i>         | 2 to 6   | 9  | 16 |
| Volunteer Cereals <sup>(3)</sup> |                                |          |    |    |
| Barley                           | <i>Hordeum vulgare</i>         | 2 to 6   | 9  | 16 |
| Oats                             | <i>Avena sativa</i>            | 2 to 6   | 9  | 16 |
| Rye                              | <i>Secale cereale</i>          | 2 to 6   | 9  | 16 |
| Wheat <sup>(2)</sup>             | <i>Triticum aestivum</i>       | 2 to 6   | 9  | 16 |
| Volunteer Corn <sup>(2,3)</sup>  | <i>Zea mays</i>                | Up to 12 | 6  | 12 |
| Volunteer Corn <sup>(3)</sup>    | <i>Zea mays</i>                | Up to 24 | 9  | 14 |
| Volunteer Corn <sup>(2,3)</sup>  | <i>Zea mays</i>                | Up to 36 | 12 | 16 |
| Volunteer Grain Sorghum          | <i>Sorghum bicolor</i>         | 8 to 12  | 9  | 16 |
| Wild Oats                        | <i>Avena fatua</i>             | 2 to 6   | 9  | 16 |
| Wild Proso Millet                | <i>Panicum miliaceum</i>       | 2 to 10  | 9  | 16 |
| Witchgrass                       | <i>Panicum capillare</i>       | 2 to 8   | 9  | 16 |
| Woolly Cupgrass                  | <i>Eriochloa villosa</i>       | 2 to 8   | 9  | 16 |

\*Generally occurs between 3-leaf stage and tillering.

\*\*Length of lateral growth.

1. Rates higher than 16 fl. oz./A may be applied in certain geographic areas, cropping situations or environmental conditions, where experience has shown that higher rates are needed for satisfactory control of annual grasses. In these situations, rates from 16 to 32 fl. oz./A may be applied. Do not apply more than 16 fl. oz./A of TAPOUT per application to the following crops: garden beets, broccoli, cabbage, carrot, cauliflower (and other head and stem brassica vegetables), celery, cranberry, cucurbits, flax, fruiting vegetables (except tomato), green onion, leaf lettuce, radish (and other root vegetables), rhubarb (and other leaf petioles), strawberry and non-bearing food crops. Do not apply more than 12 fl. oz./A of TAPOUT per application to canola or mustard seed.
2. When a cereal grain crop (such as wheat) is interseeded for crop establishment or is planted as wind breaks to aid crop establishment, the minimum TAPOUT use rate for control is 12 fl. oz./A.
3. Includes Roundup Read, Liberty Link® and IMI-CORN® volunteer corn; however not Sethoxydim-Resistant volunteer corn.

**DIRECTIONS FOR ANNUAL & PERENNIAL GRASS CONTROL  
IN ESTABLISHED ALFALFA AND MINT WITH TAPOUT**

| GRASS SPECIES                                     | WEED SPECIES AND SIZE | APPLICATION RATES  |
|---|-----------------------|--|
| Annual & Perennial Grasses Listed in Grass Tables | See Tables            | See Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT |

**Mowing:** The best control of annual grasses can be achieved by applying TAPOUT before grass weeds are mowed. Once a grass is mowed it becomes tougher to control, as much of the available leaf surface has been removed. In areas without a killing frost, some annuals can over-winter after having been mowed multiple times. These grasses form large crowns and may contain many viable buds. These grasses, even though they may be an annual grass, may require repeated applications of TAPOUT for partial or complete control.

**Irrigated Alfalfa and Mint:** Irrigation practices can be very critical to the successful use of TAPOUT in established alfalfa and mint and may be necessary to initiate active growth of the weeds prior to application. Generally applications 2 to 4 days after irrigation are most effective. Irrigation made shortly after application (2 days) can be effective, but more consistent grass control occurs when the irrigation is made before the application.

**Aerial Application:** Apply TAPOUT in a minimum of 10 GPA in established alfalfa and mint when applying by air.

**Annual Grass Control:** Apply TAPOUT at the grass sizes indicated in the Recommendation for Annual Grass Table and rates indicated. If a grass has been cut, apply TAPOUT after active growth has resumed and regrowth has reached the minimum height and before it reaches the maximum height indicated. Apply before the alfalfa/mint canopy covers the grasses and interferes with the spray coverage. Some annual grasses are spring- and summer- germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to TAPOUT may vary from region to region. Also some annuals germinate over an

extended period of time, and because control of small grasses is desired, applications after each weed flush may be required. As a general rule spray spring and summer germinating grasses as early in the season as possible, after initial green-up. Spray fall - germinating weeds in the fall soon after they begin growing but before any damage is done due to frost. Late fall applications may be less effective due to environmental conditions, such as frost, slower plant growth or the onset of flowering.

**Perennial Grass Control:** TAPOUT effectively controls perennial grasses such as bermudagrass, Johnsongrass, quackgrass, wirestem muhly, tall fescue, foxtail barley and orchardgrass. Due in part to lack of tillage, perennial grasses are more difficult to control in a perennial crop such as established alfalfa or mint. A program of repeated applications is usually necessary for best results. The best way to control perennial grasses is to do so in the year of stand establishment before rhizomes and stolons become large and difficult to kill.

Use the high rate under heavy grass pressure and/or when grasses are at or near maximum height.

**DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES CANOLA, FLAX, LEGUME VEGETABLES (DRY AND SUCCULENT), MUSTARD SEED, SOYBEAN, SUGARBEET AND SUNFLOWER (REDUCED RATE RECOMMENDATIONS NOT FOR USE IN CALIFORNIA)**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Regrowth by tillering may occur if application is made when plants are stressed by lack of moisture, excessive moisture, low or high temperatures and/or under very low humidity.

| GRASS SPECIES         | SCIENTIFIC NAME                | WEED HEIGHT (inches) | RATE FL. OZ/ACRE <sup>(1)</sup> |
|-----------------------|--------------------------------|----------------------|---------------------------------|
| Barnyardgrass         | <i>Echinochloa crus-galli</i>  | 1 to 4               | 6                               |
| Broadleaf Signalgrass | <i>Brachiaria platyphylla</i>  | 1 to 4               | 8                               |
| Crabgrass             |                                |                      |                                 |
| Large                 | <i>Digitaria san guinalis</i>  | 1 to 3*              | 6                               |
| Large                 | <i>Digitaria sanguinalis</i>   | 1 to 4*              | 8                               |
| Smooth                | <i>Digitaria ischaemum</i>     | 1 to 3*              | 6                               |
| Smooth                | <i>Digitaria ischaemum</i>     | 1 to 4*              | 8                               |
| Southern              | <i>Digitaria char</i>          | 1 to 4*              | 8                               |
| Fall Panicum          | <i>Panicum dichotomiflorum</i> | 1 to 4               | 6                               |
| Foxtail               |                                |                      |                                 |
| Giant                 | <i>Setaria faberi</i>          | 1 to 4               | 6                               |
| Green                 | <i>Setaria viridis</i>         | 1 to 4               | 6                               |
| Millet                | <i>Setaria italica</i>         | 1 to 4               | 8                               |
| Yellow                | <i>Setaria glauca</i>          | 1 to 4               | 6                               |
| Seedling Johnsongrass | <i>Sorghum halepense</i>       | 1 to 6               | 8                               |
| Shattercane           | <i>Sorghum bicolor</i>         | 4 to 10              | 6                               |
| Texas Panicum         | <i>Panicum texanum</i>         | 1 to 4               | 8                               |
| Volunteer Cereals     |                                |                      |                                 |
| Barley                | <i>Hordeum vulgare</i>         | 1 to 4               | 8                               |
| Oats                  | <i>Avena sativa</i>            | 1 to 4               | 8                               |
| Wheat                 | <i>Triticum aestivum</i>       | 1 to 4               | 8                               |
| Volunteer Corn**      | <i>Zea mays</i>                | 4 to 12              | 6                               |
| Wild Proso Millet     | <i>Panicum miliaceum</i>       | 1 to 6               | 6                               |
| Wild Oats             | <i>Avena fatua</i>             | 1 to 4               | 8                               |

\*Length of lateral growth.

\*\*Not S.R. Corn

<sup>(1)</sup>Always add a non-ionic surfactant at 0.25% v/v total spray volume unless crop specific restrictions and limitations advise otherwise.

**DIRECTIONS FOR PERENNIAL GRASSES (ALL CROPS)**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- Do not exceed the maximum per application rate listed in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT.



| GRASS SPECIES   | WEED HEIGHT (inches)    | APPLICATION RATE       |                        |
|---|-------------------------|------------------------|------------------------|
|   |                         | MINIMUM RATE fl. oz./A | MAXIMUM RATE fl. oz./A |
| Bermudagrass ( <i>Cynodon dactylon</i> )  |                         |                        |                        |
| First Application   | 3 (or up to 6" runners) | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 3 (or up to 6" runners) | 12                     | 32                     |
| Fescue, Tall ( <i>Festuca arundinacea</i> )   |                         |                        |                        |
| First Application   | 4 to 8                  | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 4 to 8                  | 12                     | 32                     |
| Foxtail Barley ( <i>Hordeum jubatum</i> )   |                         |                        |                        |
| First Application   | 2 to 6                  | 12                     | 32                     |
| Repeat Application (if regrowth occurs)   | 2 to 6                  | 12                     | 32                     |
| Orchardgrass ( <i>Dactylis glomerata</i> )  |                         |                        |                        |
| First Application   | 4 to 8                  | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 4 to 8                  | 12                     | 32                     |
| Quackgrass ( <i>Elytrigia repens</i> )  |                         |                        |                        |
| First Application   | 4 to 12                 | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 4 to 12                 | 12                     | 32                     |
| Rhizome Johnsongrass ( <i>Sorghum halepense</i> )   |                         |                        |                        |
| First Application   | 12 to 24                | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 6 to 18                 | 9                      | 24                     |
| Wirestem Muhly ( <i>Muhlenbergia frondosa</i> )   |                         |                        |                        |
| First Application   | 4 to 8                  | 12                     | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 4 to 8                  | 12                     | 32                     |
| Perennial Bluegrass<br>[Roughstalk ( <i>Poa trivialis</i> )<br>[Kentucky ( <i>Poa pratensis</i> ) |                         |                        |                        |
| First Application   | 2 to 4                  | 12                     | 32                     |
| Repeat Application(s)   | 2 to 4                  | 12                     | 32                     |
| Bentgrass ( <i>Agrostis</i> spp.)   |                         |                        |                        |
| First Application   | 2 to 4                  | -                      | 32                     |
| Repeat Application(s) (if regrowth occurs)  | 2 to 4                  | -                      | 32                     |

| DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL WITH TAPOUT<br>ALL CROPS   |            |                            |                           |
|--|------------|----------------------------|---------------------------|
| GRASS SPECIES  | WEED STAGE | APPLICATION RATES          |                           |
|  |            | MINIMUM RATE<br>Fl. Oz./ A | MAXIMUM RATE<br>Fl. Oz./A |
| Annual Bluegrass ( <i>Poa annua</i> )  | to 4-leaf  | 12*                        | **                        |
| <p>Apply under favorable soil moisture and humidity, which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</p> <p>Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass. Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</p> <p>See Table 1 for crop specific adjuvant recommendations.</p> <p>*Use a minimum of 17 fl. oz./A to control annual bluegrass in seedling and established alfalfa and mint.</p> <p>**See Special Use Instructions and Restrictions in Table 1, CROP SPECIFIC USE DIRECTIONS, RESTRICTIONS AND LIMITATIONS FOR TAPOUT, for maximum application rates.</p> |            |                            |                           |

### TANK MIXES PRODUCT INFORMATION

The labels for each of the herbicides listed for tank mixing with **TAPOUT** are unique to the characteristics of those products and contain restrictions and limitations that may be more restrictive than the **TAPOUT** label in certain considerations. Those concerns may include, but are not limited to:

1. Geographic restrictions - all products are not registered for use in all areas and rates may vary from one region of labeled use to another;
2. Crop rotation restrictions;
3. Applicator certification requirements;
4. Worker safety rules (e.g. protective clothing, reentry time, posting);
5. Soil type or soil characteristics (e.g. pH, OM);
6. Maximum dosage or number of applications per season;
7. Rain free period required; or
8. Application timing (e.g. pre-harvest interval)
9. Do not exceed the total season rates.

**THE MOST RESTRICTIVE LABELING OF ANY PRODUCT USED IN A TANK MIX MUST BE FOLLOWED.**

#### TANK MIX APPLICATION OF TAPOUT AND BROADLEAF HERBICIDES FOR CONTROL OF GRASSES AND BROADLEAF WEEDS

- Apply only to actively growing grass and broadleaf weeds at specified height or growth stage listed on each label.
- Apply when the first grass or broadleaf weed species in a mixed population reaches the specified height or growth stage for treatment.
- Apply under favorable soil moisture and humidity that exist a few days after rainfall or within seven days after irrigation.
- Always add the appropriate adjuvant to the spray mix at the rate specified for each specific tank mix combination.
- Tank mix applications may sometimes result in reduced grass control and possible increases in crop injury as compared to either product used alone. If regrowth occurs, or an additional flush of new grass emerges, make a second application of **TAPOUT**, as specified in the respective size and rate tables.
- Do not tank mix **TAPOUT** when broadleaf weeds are tall and/or dense enough to prevent proper grass coverage.

#### MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of **TAPOUT**. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing **TAPOUT** with Other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**

Failure to agitate the spray solution may result in improper mixing of the herbicides and unsatisfactory weed control. Mixing and compatibility qualities should be verified by a jar test.

#### INFORMATION ON ANTAGONISM

Tank mixes of **TAPOUT** with postemergence broadleaf herbicides have shown some reduction or failure to control certain grass species which would have otherwise been controlled when **TAPOUT** is applied alone. Activity of the postemergence broadleaf herbicide in the tank mix is not affected.

ALFALFA

**Table 2. TAPOUT I EC TANK MIXES WITH BROADLEAF HERBICIDES FOR ALFALFA**  
(Refer to the recommendation tables above for specific grassed and growth stages.)

| PRODUCT <sup>(1)</sup>  | APPLICATION RATES/ACRE <sup>(2)</sup>                 |   | SPRAY ADDITIVES         |                   |                         |                                    |
|---|---|---|-------------------------|-------------------|-------------------------|------------------------------------|
|   | ANNUAL GRASSES  | PERENNIAL GRASSES                         | GROUND APPLICATION      |                   | AIR APPLICATION         |                                    |
|   |   |   | Adjuvant Recommendation | AMS               | Adjuvant Recommendation | AMS                                |
| TAPOUT + 2,4-DB <sup>(3)</sup>                                      | 12 to 32 fl. oz.<br>Refer to 2,4-DB label             | 16 to 32 fl. oz.<br>Refer to 2,4-DB label | NIS at 0.25% v/v        | AMS at 2.5 lbs./A | NIS at 0.25% v/v        | 17 lbs/100 gals of spray solution  |
| TAPOUT + PURSUIT® DG <sup>(4)</sup> or PURSUIT <sup>(4)</sup>       | 12 to 32 fl. oz. + 1.08 to 2.16 oz. or 3 to 6 fl. oz. |   | NIS at 0.25% v/v        | AMS at 2.5 lbs./A | NIS at 0.25% v/v        | 17 lbs/100 gals. of spray solution |
| TAPOUT + BUCTRIL® 2L <sup>(5)</sup> or BUCTRIL GEL <sup>(5,6)</sup> | 12 to 32 fl. oz. + 1.0 to 1.5 pts. or 0.5 to 0.75 pt. |   | NIS at 0.25% v/v        | AMS at 2.5 lbs./A | NIS at 0.25% v/v        | 17 lbs/100 gals. of spray solution |
| TAPOUT + RAPTOR®  | 12 to 32 fl. oz. + 4 to 6 fl. oz.                     |   | NIS at 0.25% v/v        | AMS at 2.5 lbs./A | NIS at 0.25% v/v        | 17 lbs/100 gals. of spray solution |

- Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
- If grass regrowth occurs or an additional flush of new grass emerges, make a second application of TAPOUT alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
- TAPOUT plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.
- Before using this tank mix, read and understand the PURSUIT or PURSUIT DG labels for geographical restrictions and restrictions regarding alfalfa growth stage and type. Failure to do so can result in crop injury to alfalfa. Do not feed, graze or harvest alfalfa for 30 days following an application of PURSUIT to alfalfa.
- In the states of Washington, Oregon, Idaho, Montana, Wyoming, Colorado, Utah, Nevada and the western halves of North Dakota, South Dakota, Nebraska and Kansas: The TAPOUT plus BUCTRIL or BUCTRIL GEL tank mix must be applied in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 2 trifoliates. Unacceptable crop injury may occur to alfalfa seedlings less than the 2 trifoliolate leaf stage. TAPOUT plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 80°F at and 3 days following application can result in unacceptable crop injury. In the states not listed above, apply in the fall or spring to seedling alfalfa when the majority of the field has a minimum of 4 trifoliolate leaves. When alfalfa stand is uneven and conditions favor leafburn, unacceptable crop injury may occur to alfalfa in the 2 trifoliolate or smaller stage of growth. TAPOUT plus BUCTRIL or BUCTRIL GEL applications made when temperatures are expected to exceed 70°F at and 3 days following application can result in unacceptable crop injury. Crop leaf burn can occur following TAPOUT plus BUCTRIL or BUCTRIL GEL application. Warm, humid conditions may enhance leaf burn. New crop growth will not be affected.
- Do not apply when alfalfa is under moisture, temperature, insect or disease stress or has been stressed by other pesticide carryover or application.
- Always use a CPDA Certified Adjuvant when possible.

**CANOLA (Not for Use in California)**

**Table 3. REDUCED RATE TAPOUT TANK MIXES WITH BROADLEAF HERBICIDES FOR CANOLA (Refer to the recommendation tables above for specific grasses and growth stages.)**

| PRODUCT   | APPLICATION RATES/ACRE        |                  | ADJUVANT RECOMMENDATIONS | AMMONIUM SULFATE |          |
|---|-------------------------------|------------------|--------------------------|------------------|----------|
|   | ANNUAL GRASSES <sup>(1)</sup> |                  |                          | GROUND           | AIR      |
| TAPOUT <sup>(2)</sup><br>+<br>LIBERTY® <sup>(3)</sup> | 8 to 10 fl.<br>+              | 28 to 34 fl. oz. | NIS at 0.25% v/v         | 3 lbs./A         | 3 lbs./A |
| TAPOUT<br>+<br>STINGER® <sup>(4)</sup>                | 8 to 10 fl. oz.<br>+          | 0.33 pts./A      | NIS at 0.25% v/v         | 3 lbs./A         | 3 lbs./A |

1. Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.
2. Do not apply TAPOUT tank mix during or after bolting or flowering or crop injury will occur.
3. For use only on LIBERTY LINK® Canola.
4. See STINGER label for weeds controlled.
5. Always use a CPDA Certified Adjuvant when possible.

**COTTON**

**Table 4. TAPOUT TANK MIXED WITH COBRA AND MSMA APPLIED POST DIRECTED TO COTTON**

| PRODUCT <sup>(1)</sup>  | APPLICATION RATES/ACRE(2)   |                   | CROP OIL CONCENTRATE <sup>(3)</sup> | COMMENTS   |
|---|---|-------------------|-------------------------------------|--|
|   | ANNUAL GRASSES  | PERENNIAL GRASSES |                                     |  |
| TAPOUT<br>+<br>COBRA<br>+<br>MSMA<br>(4 lbs./gal.)<br>OR<br>MSMA<br>(6.6 lbs./gal.) | 12 to 16 fl. oz.  | 16 to 32 fl. oz.  | 1%                                  | Reduce broadcast rate in proportion to the band area actually treated. |
|   | See COBRA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the TAPOUT label for weed height and species controlled. |                   |                                     |  |
|   | See MSMA label for rates to control broadleaf weeds and height limitations for cotton. Refer to the TAPOUT label for weed height and species controlled.  |                   |                                     |  |

1. Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
2. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of TAPOUT alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
3. Always use a crop oil concentrate at the listed rate (but not less than 1 pt/A) in the finished spray volume.
4. If at the time of application, grass height is so tall that post-directed applications cannot get good coverage over the top of the grassy weeds, then poor control may result and a second (non-post directed) application of TAPOUT may be necessary.
5. Always use a CPDA Certified Adjuvant when possible.

**Table 5. TAPOUT TANK MIXED WITH GLYPHOSATE TO CONTROL EMERGED GRASSES IN COTTON AS A BROADCAST APPLICATION**

| PRODUCT                   | APPLICATION RATE/ACRE (1)   |                   | ADJUVANT   |  | COMMENTS  |
|---------------------------|---|-------------------|--|--|---|
|                           | ANNUAL GRASSES  | PERENNIAL GRASSES | Glyphosate formulation with built in adjuvant.   | Glyphosate formulation without built in adjuvant.                                  |   |
| TAPOUT<br>+<br>GLYPHOSATE | 9 to 16 fl. oz.<br>See glyphosate label for rates to control broadleaf weeds and height limitations for cotton. | 12 to 32 fl. oz.  | Ammonium sulfate at 8.5 to 17 lbs. per 100 gals. of carrier plus glyphosate label adjuvant recommendation. | Ammonium sulfate at 8.5 to 17 lbs. per 100 gals. Of carrier plus NIS at 0.25% v/v. | See charts for grasses controlled.<br><br>Use a minimum of 10 gals. Of spray solution per acre. |

1. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of TAPOUT at the specified rate with the appropriate amount of crop oil.
2. Always use a CPDA Certified Adjuvant when possible.

**DRY AND SUCCULENT SHELLED BEANS**

**Table 6. TAPOUT TANK MIXES WITH BROADLEAF HERBICIDES FOR DRY AND SUCCULENT SHELLED BEANS**  
(Refer to the recommendation tables above for specific grasses and growth stages.)

| PRODUCT <sup>(1)</sup>                  | APPLICATION RATES/ACRE <sup>(2)</sup> |                                      | ADJUVANT                                      |  |
|---|---------------------------------------|--------------------------------------|---|--|
|   | ANNUAL GRASSES                        | PERENNIAL GRASSES                    | GROUND  | AIR  |
| TAPOUT<br>+<br>BASAGRAN <sup>®(3)</sup> | 9 to 12 fl. oz.<br>+<br>1 to 2 pts.   | 12 to 24 fl. oz.<br>+<br>1 to 2 pts. | COC at 1% v/v<br>+<br>AMS at 2.5 lbs./A       | COC at 1% v/v<br>+<br>AMS at 17<br>lbs./100 gal. v/v   |
| TAPOUT<br>+<br>RAPTOR <sup>(3)</sup>    | 9 to 12 fl. oz.<br>+<br>4 fl. oz.     | -                                    | NIS at 0.25%<br>v/v<br>+<br>AMS at 2.5 lbs./A | NIS at 0.25%<br>v/v<br>+<br>AMS at 17<br>lbs./100 gal. |

1. Broadleaf weed control may be reduced when grass populations are tall enough or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
3. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of TAPOUT alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
4. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed.
5. Always use a CPDA Certified Adjuvant when possible.

**FLAX (Not for Use in California)**

**Table 7. REDUCED RATE TAPOUT TANK MIXES WITH BROADLEAF HERBICIDES FOR FLAX**  
(Refer to the recommendation tables above for specific grasses and growth stages.)

| PRODUCT   | APPLICATION RATES/ACRE                    | ADJUVANT  |                          |
|---|---|---|--------------------------|
|   | ANNUAL GRASSES <sup>(1)</sup>             | GROUND  | AIR                      |
| TAPOUT<br>+<br>BRONATE ADVANCED <sup>™(2,3)</sup> | 6 to 9 fl. oz.<br>+<br>11.4 fl. oz.       | AMS at 2.4 to 4.0<br>lbs./A<br>+<br>NIS at 0.125% v/v | AMS at 2.5 to 4.0 lbs./A |
| TAPOUT<br>+<br>BRONATE <sup>®(2,3)</sup>          | 6 to 9 fl. oz.<br>+<br>0.9 pt.            | AMS at 2.4 to 4.0 lbs./A<br>+<br>NIS at 0.125% v/v    | AMS at 2.5 to 4.0 lbs./A |
| TAPOUT<br>+<br>BUCTRIL <sup>(2,3)</sup>           | 6 to 9 fl. oz.<br>+<br>0.125 lb. al/A     | AMS at 2.4 to 4.0 lbs./A<br>+<br>NIS at 0.125% v/v    | AMS at 2.5 to 4.0 lbs./A |
| TAPOUT<br>+<br>MCPA23                             | 8 to 10 fl. oz.<br>+<br>0.25 to 0.5 pt.   | AMS at 2.4 to 4.0 lbs./A<br>+<br>NIS at 0.125% v/v    | AMS at 2.5 to 4.0 lbs./A |
| TAPOUT<br>+<br>CURTAIL <sup>® M(2,3)</sup>        | 6 to 9 fl. oz.<br>+<br>1.33 to 1.75 pt./A | AMS at 2.4 to 4.0 lbs./A<br>+<br>NIS at 0.125% v/v    | AMS at 2.5 to 4.0 lbs./A |

1. Annual grasses and sizes controlled with these tank mixtures are those that are identified in the DIRECTIONS FOR REDUCED RATE USE TO CONTROL SMALL ANNUAL GRASSES table.
2. Do not apply TAPOUT tank mix during or after the bud stage or to ornamental flax or crop injury may occur.
3. Do not apply tank mixes if temperatures are expected to exceed 85F at (or 3 days following) application or crop injury may occur.
4. Always use a CPDA Certified Adjuvant when possible.

PEANUT

**Table 8. TAPOUT TANK MIXES WITH BROADLEAF HERBICIDES FOR PEANUT**  
 (Refer to the recommendation tables above for specific grasses and growth stages.)

| PRODUCT <sup>(1)</sup>       | APPLICATION RATES/ACRE <sup>(2)</sup>   | ADJUVANT RECOMMENDATIONS                |  |
|------------------------------|---|---|--|
|                              | ANNUAL GRASSES                          | GROUND                                  | AIR  |
| TAPOUT<br>+<br>BASAGRAN      | 9 to 16 fl. oz.<br>+<br>1 to 2 pts.     | COC at 1% v/v<br>+<br>AMS at 2.5 lbs./A | COC at 1% v/v<br>+<br>AMS at 17 lbs./100 gals. |
| TAPOUT<br>+<br>ULTRA BLAZER® | 9 to 16 fl. oz.<br>+<br>0.5 to 1.5 pts. | COC at 1% v/v<br>+<br>AMS at 2.5 lbs./A | COC at 1% v/v<br>+<br>AMS at 17 lbs./100 gals. |
| TAPOUT<br>+<br>STORM®        | 9 to 16 fl. oz.<br>+<br>1.5 pts.        | COC at 1% v/v<br>+<br>AMS at 2.5 lbs./A | COC at 1% v/v<br>+<br>AMS at 17 lbs./100 gals. |

1. Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
2. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of TAPOUT alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
3. Always use a CPDA Certified Adjuvant when possible.

**DIRECTIONS FOR GRASS SUPPRESSION FOR HARVEST EFFICIENCY  
 IN PEANUT WITH TAPOUT**

| GRASS SPECIES  | WEED STAGE   | APPLICATION RATES         |                           |
|--|--|---------------------------|---------------------------|
|  |  | MINIMUM RATE<br>fl. oz./A | MAXIMUM RATE<br>fl. oz./A |
| Annual and perennial grasses that exceed height claimed for control on height charts "RECOMMENDATIONS FOR ANNUAL GRASSES" and "RECOMMENDATIONS FOR PERENNIAL GREASSES" | Up to and including grasses in the seed head stage | 32                        | 64                        |

- Do not apply as part of a tank mix when applying TAPOUT for grass suppression.
- Add a crop oil concentrate at 1 gal. /A by ground to the finished spray volume.
- Always use a CPDA Certified Adjuvant when possible.

SOYBEAN

**Table 9. TAPOUT TANK MIX WITH BROADLEAF HERBICIDES FOR THE CONTROL OF VOLUNTEER CORN (INCLUDING ROUNDUP READY) IN SOYBEAN**

(Refer to the recommendation tables above for specific volunteer corn sizes and use rates.)

| PRODUCT   | WEED SIZE AND APPLICATION RATES |                   | SPRAY ADDITIVES   |  |   |   |
|---|---------------------------------|-------------------|---|--|---|---|
|   | VOLUNTEER CORN HEIGHT (inches)  | TAPOUT RATES/ACRE | GROUND APPLICATION  |  | AIR APPLICATION   |   |
|   |                                 |                   | NIS   | AMS                                      | NIS   | AMS                                       |
| TAPOUT + gIyphosate <sup>(1,2,3)</sup><br>1 to 3 lbs. al/A<br>(ROUNDUP READY soybeans only) | Up to 12                        | 6fl.oz.           | Adjuvant Loaded Glyphosate: None Required<br><br>Adjuvant Unloaded Glyphosate: NIS at 0.25% v/v | 8.5 to 17 lbs/100 gals of spray solution | Adjuvant Loaded Glyphosate: None Required<br><br>Adjuvant Unloaded Glyphosate: NIS at 0.25% v/v | 8.5 to 17 lbs/100 gals. of spray solution |
|   | Up to 24                        | 9 fl. oz.         |   |  |   |   |
|   | Up to 36                        | 12 fl. oz.        |   |  |   |   |
| TAPOUT FIRSTRATE®<br>0.3 oz./A  | Up to 12                        | 6 fl. oz.         | NIS at 0.25% v/v  | AMS at 2.5 lbs./A                        | NIS at 0.25% v/v  | 17 lbs./100 gals. of spray solution       |
|   | Up to 24                        | 9 fl. oz.         |   |  |   |   |
|   | Up to 36                        | 12 fl. oz.        |   |  |   |   |
| TAPOUT + PURSUIT 70 DG<br>1.44 oz./A  | Up to 12                        | 6fl.oz.           | NIS at 0.25% v/v  | AMS at 2.5 lbs./A                        | NIS at V-0.25% v/v  | 17lbs./100 gals. of spray solution        |
|   | Up to 24                        | 9 fl. oz.         |   |  |   |   |
|   | Up to 36                        | 12 fl. oz.        |   |  |   |   |
| TAPOUT + RAPTOR<br>4 to 5 fl. oz./A   | Up to 12                        | 6fl. oz.          | NIS at 0.25% v/v  | AMS at 2.5 lbs/A                         | NIS at 0.25% v/v  | 17 lbs./100 gals. of spray solutions      |
|   | Up to 24                        | 9 fi. OZ          |   |  |   |   |
|   | Up to 36                        | 12 fl. oz.        |   |  |   |   |

1. This tank mix may be applied postemergence to ROUNDUP Ready soybeans up through the full flowering stage. Do not apply less than 60 days before harvest.
2. Avoid contact with foliage, green stems or fruit crops or any desirable plants and trees, other than soybeans with the ROUNDUP Ready gene as severe plant injury or death will result.
3. Do not allow the TAPOUT plus glyphosate to mist, drip, drift or splash onto desirable vegetation as minute quantities of the tank mix can cause severe damage or destruction to the crops, plants or other areas on which treatment was not intended. The likelihood of injury occurring from drift of this product is greatest when winds are gusty or in excess of 5 miles per hour. Even under lesser wind velocities, avoid conditions that allow spray drift to occur such as combinations of spray pressure and nozzle type that will result in fine particles (mist) that are likely to drift.
4. Always use a CPDA Certified Adjuvant when possible.

**Table 10. TAPOUT TANK MIXES WITH BROADLEAF HERBICIDES FOR SOYBEAN**  
 (Refer to the recommendation tables above for specific grasses and growth stages.)

| PRODUCT                                      | APPLICATION RATES/ACRE <sup>(2)</sup>  | SPRAY ADDITIVE RECOMMENDATIONS   |                      |  |   |
|--|--|--|----------------------|--|---|
|  | ANNUAL GRASSES <sup>(3)</sup>  | GROUND APPLICATION   |                      | AIR APPLICATION  |   |
|  |  | COC/NIS <sup>(4)</sup>   | AMS                  | COC/NIS <sup>(4)</sup>   | AMS                                       |
| TAPOUT<br>+<br>COBRA                         | 9 to 20 fl. oz.<br>+<br>6 to 12 fl. oz.  | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A    | AMS at 2.5<br>lbs/A  | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC 1% v/v (but<br>not less than 1<br>pt/A)    | 17 lbs./100 gals.<br>of spray solution    |
| TAPOUT<br>+<br>FIRSTRATE <sup>(5)</sup>      | 9 to 20 fl. oz.<br>+<br>0.3 oz.  | NIS at 0.25%<br>v/v<br>or<br>COC at 1 pt./A                                      | AMS at 2.5<br>lbs./A | NIS at 0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A)                            | 17 lbs/100 gals.<br>of<br>spray solution  |
| TAPOUT<br>+<br>FLEXSTAR® HL <sup>(5)</sup>   | 9 to 20 fl. oz.<br>+<br>Refer to the<br>FLEXSTAR HL<br>label<br>for specific<br>application rates. | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A    | AMS at 2.5<br>lbs/A  | NIS at 0.25%<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt/A)     | 17 lbs/100 gals.<br>of<br>spray solution  |
| TAPOUT<br>+<br>FRONTROW™ <sup>(5)</sup>      | 9 to 20 fl. oz.<br>+<br>Refer to<br>FRONTROW label<br>for use rates                                | NIS at 0.25%<br>v/v<br>or<br>COC at 1 pt./A                                      | AMS at 2.5<br>lbs./A | NIS at 0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A)                            | 17 lbs./100 gals.<br>of<br>spray solution |
| TAPOUT<br>+<br>Harmony® GT XP <sup>(5)</sup> | 9 to 12 fl. oz.<br>+<br>0.042 to 0.083 oz.   | NIS at 0.125 to 0.25%<br>v/v   | AMS at 2.5<br>lbs/A  | -  | -   |
| TAPOUT<br>+<br>PHOENIX™                      | 9 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.  | NIS at 0.25%<br>v/v plus COC at<br>0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A | AMS at 2.5<br>lbs/A  | NIS at 0.25%<br>plus<br>COC at 0.25%<br>v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |
| TAPOUT<br>+<br>PURSUIT 70 DG <sup>(5)</sup>  | 12 to 20 fl. oz.<br>+<br>1.44 oz.  | NIS at 0.25% v/v<br>or<br>COC at 1 pt./A   | AMS at 2.5<br>lbs/A  | MS at 0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A)                             | 17 lbs./100 gals.<br>of<br>spray solution |
| TAPOUT<br>+<br>RAPTOR(1 AS) <sup>(5)</sup>   | 12 to 20 fl. oz.<br>+<br>4 to 5 fl. oz.  | NIS at 0.25%<br>v/v<br>or<br>COC at 1 pt./A                                      | AMS at 2.5<br>lbs./A | NIS at 0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A)                            | 17 lbs./100 gals.<br>of<br>spray solution |
| TAPOUT<br>+<br>RESOURCE®                     | 9 to 20 fl. oz.<br>+<br>4 to 12 fl. oz.  | NIS at 0.25%<br>v/v plus CCC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A    | AMS at 2.5<br>lbs/A  | -  | -   |



|  |   |  |                     |   |   |
|--|---|--|---------------------|---|---|
| <b>TAPOUT</b><br>+<br><b>COBRA</b><br>+<br><b>FIRSTRATE</b> <sup>(5)</sup>       | 9 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>0.3 oz.  | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A  | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |
| <b>TAPOUT</b><br>+<br>Cobra<br>+<br>Harmony GT XP <sup>(5)</sup>                 | 9 to 12 fl. oz.<br>+<br>6 to 8 fl. oz.<br>+<br>0.042. oz.   | NIS at 0.125 to<br>0.25% v/v plus<br>COC at 0.125%<br>v/v  | AMS at 2.5<br>lbs/A | -   | -   |
| <b>TAPOUT</b><br>+<br><b>COBRA</b><br>+<br><b>PURSUIT 70 DG</b> <sup>(5)</sup>   | 12 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>1.44 oz.   | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A  | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of spray solution    |
| <b>TAPOUT</b><br>+<br><b>COBRA</b><br>+<br><b>RAPTOR (1 AS)</b> <sup>(5)</sup>   | 12 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>4 to 5 fl. oz.   | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A  | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |
| <b>TAPOUT</b><br>+<br><b>COBRA</b><br>+<br><b>RESOURCE</b>                       | 9 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>4 to 6 fl. oz.  | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>COC at 1 to 2<br>pts./A  | AMS at 2.5<br>lbs/A | -   | -   |
| <b>TAPOUT</b><br>+<br><b>FIRSTRATE</b><br>+<br><b>FLEXSTAR HL</b> <sup>(5)</sup> | 9 to 20 fl. oz.<br>+<br>0.3 oz.<br>+<br>Refer to the<br>FLEXSTAR HL<br>label<br>for specific<br>application rates | NIS at 0.25%<br>v/v plus COC at<br>0.25% v/v<br>or<br>Equivalent<br>blended product<br>or<br>COC at 1 to 2<br>pts./A | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |
| <b>TAPOUT</b><br>+<br><b>PHOENIX</b><br>+<br><b>FIRSTRATE</b> <sup>(5)</sup>     | 12 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>0.3 oz.  | NIS at 0.25%<br>v/v plus COC at<br>0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A                                     | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of spray solution    |
| <b>TAPOUT</b><br>+<br><b>PHOENIX</b><br>+<br><b>PURSUIT 70 DG</b> <sup>(5)</sup> | 16 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>1.44 oz.   | NIS at 0.25% v/v plus<br>COC at 0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A  | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |

|   |   |  |                              |  |  |
|---|---|--|------------------------------|--|--|
| <p><b>TAPOUT</b><br/>+<br/><b>PHOENIX</b><br/>+<br/><b>RAPTOR(1 AS)<sup>(5)</sup></b></p>   | <p>12 to 20 fl. oz.<br/>-<br/>6 to 12.5 fl. oz.<br/>+<br/>4 to 5 fl. oz.</p>                  | <p>NIS at 0.25%<br/>v/v plus COC at<br/>0.125 to 0.25%<br/>v/v<br/>or<br/>COC at 1 pt./A</p> | <p>AMS at 2.5<br/>lbs./A</p> | <p>NIS at 0.25% v/v<br/>plus COC at<br/>0.25% v/v<br/>or<br/>COC at % v/v<br/>(but not less than<br/>1 pt./A)</p>  | <p>17 lbs./100 gals.<br/>of spray solution</p>     |
| <p><b>TAPOUT</b><br/>+<br/><b>PHOENIX</b><br/>+<br/><b>RESOURCE</b></p>   | <p>9 to 20 fl. oz.<br/>+<br/>6 to 12.5 fl. oz.<br/>+<br/>4 to 6 fl. oz.</p>                   | <p>NIS at 0.25%<br/>v/v plus COC at<br/>0.125 to 0.25%<br/>v/v<br/>or<br/>COC at 1 pt./A</p> | <p>AMS at 2.5<br/>lbs./A</p> | -  | -  |
| <p><b>TAPOUT</b><br/>+<br/><b>RESOURCE</b><br/>+<br/><b>PURSUIT 70 DG<sup>(5)</sup></b></p>   | <p>12 to 20 fl. oz.<br/>+<br/>4 fl. oz.<br/>+<br/>1.44 oz.</p>                                | <p>NIS at 0.25%<br/>v/v plus COC at<br/>0.25% v/v<br/>or<br/>COC at 1 to 2<br/>pts./A</p>    | <p>AMS at 2.5<br/>lbs./A</p> | -  | -  |
| <p><b>TAPOUT</b><br/>+<br/><b>SYNCHRONY<sup>®</sup></b><br/><b>STS<sup>(5)</sup>&gt;</b><br/>or<br/><b>SYNCHRONY XP</b><br/><b>(mp)<sup>(5)</sup></b></p>                     | <p>12 to 20 fl. oz.<br/>+<br/>0.25 oz.<br/>or<br/>0.375 oz.</p>                               | <p>NIS at 0.25%<br/>v/v<br/>or<br/>COC at 1 pt./A</p>  | <p>AMS at 2.5<br/>lbs./A</p> | <p>NIS at 0.25% v/v<br/>or<br/>COC at 1% v/v<br/>(but not less than<br/>1 pt./A)</p>                               | <p>17 lbs./100 gals.<br/>of<br/>spray solution</p> |
| <p><b>TAPOUT</b><br/>+<br/><b>SYNCHRONY STS<sup>(5)</sup></b><br/>or<br/><b>SYNCHRONY XP</b><br/><b>(mp)<sup>(5)</sup></b><br/>(STS Soybeans Only)</p>                        | <p>12 to 20 fl. oz.<br/>+<br/>0.5 oz.<br/>or<br/>0.75 oz.</p>                                 | <p>NIS at 0.25%<br/>v/v or<br/>COC at 1 pt./A</p>  | <p>AMS at 2.5<br/>lbs./A</p> | <p>NIS at 0.25% v/v<br/>or<br/>COC at 1% v/v<br/>(but not less than<br/>1 pt./A)</p>                               | <p>17 lbs./100 gals.<br/>of<br/>spray solution</p> |
| <p><b>TAPOUT</b><br/>+<br/><b>COBRA</b><br/>+<br/><b>RESOURCE</b><br/>+<br/><b>FIRSTRATE<sup>(5)</sup></b></p>  | <p>9 to 12 fl. oz.<br/>+<br/>6 to 12.5 fl. oz.<br/>+<br/>4 to 6 fl. oz.<br/>+<br/>0.3 oz.</p> | <p>NIS at 0.25%<br/>v/v plus COC at<br/>0.25% v/v<br/>or<br/>COC at 1 to 2<br/>pts./A</p>    | <p>AMS at 2.5<br/>lbs./A</p> | -  | -  |
| <p><b>TAPOUT</b><br/>+<br/><b>COBRA</b><br/>+<br/><b>SYNCHRONY STS<sup>(5)</sup></b><br/>or<br/><b>SYNCHRONY XP</b><br/><b>(mp)<sup>(5)</sup></b></p>                         | <p>12 to 20 fl. oz.<br/>+<br/>6 to 12 fl. oz.<br/>+<br/>0.25 oz.<br/>or<br/>0.375 oz.</p>     | <p>NIS at 0.25%<br/>v/v plus COC at 0.25%<br/>v/v<br/>or<br/>COC at it 1 to 2 pts./A</p>     | <p>AMS at 2.5<br/>lbs./A</p> | <p>NIS at 0.25% v/v<br/>plus COC at<br/>0.25% v/v<br/>or<br/>COC at 1% v/v<br/>(but not less than<br/>1 pt./A)</p> | <p>17 lbs./100 gals.<br/>of spray solution</p>     |
| <p><b>TAPOUT</b><br/>+<br/><b>COBRA</b><br/>+<br/><b>SYNCHRONY STS<sup>(5)</sup></b><br/>or<br/><b>SYNCHRONY XP</b><br/><b>(mp)<sup>(5)</sup></b><br/>(STS Soybeans Only)</p> | <p>12 to 20 fl. oz.<br/>+<br/>6 to 12 fl. oz.<br/>+<br/>0.5 oz.<br/>or<br/>0.75 oz.</p>       | <p>NIS at 0.25%<br/>v/v plus COC at<br/>0.25% v/v<br/>or<br/>COC at 1 to 2<br/>pts./A</p>    | <p>AMS at 2.5<br/>lbs./A</p> | <p>NIS at 0.25% v/v<br/>plus COC at<br/>0.25% v/v<br/>or<br/>COC at 1% v/v<br/>(but not less than<br/>1 pt./A)</p> | <p>17 lbs./100 gals.<br/>of spray solution</p>     |

|  |  |  |                     |   |   |
|--|--|--|---------------------|---|---|
| <b>TAPOUT</b><br>+<br><b>PHOENIX</b><br>+<br><b>RESOURCE</b><br>+<br><b>FIRSTRATE<sup>(5)</sup></b>  | 9 to 20 fl. oz.<br>+<br>6 to 12.5 fl. oz.<br>+<br>4 to 6 fl. oz.<br>+<br>0.3 oz. | NIS at 0.25%<br>v/v plus COC at<br>0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A | AMS at 2.5<br>lbs/A | -   | -   |
| <b>TAPOUT</b><br>+<br><b>PHOENIX</b><br>+<br><b>SYNCHRONY</b><br><b>STS<sup>(5)</sup></b><br>or<br><b>SYNCHRONY XP</b><br>(mp) <sup>(5)</sup>              | 12 to 20 fl. oz.<br>+<br>6 to 12 fl. oz.<br>+<br>0.25 oz.<br>or<br>0.375         | NIS at 0.25%<br>v/v plus COC at<br>0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>Or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs./100 gals.<br>of<br>spray solution |
| <b>TAPOUT</b><br>+<br><b>PHOENIX</b><br>+<br><b>SYNCHRONY STS<sup>(5)</sup></b><br>or<br><b>SYNCHRONY XP</b><br>(mp) <sup>(5)</sup><br>(STS Soybeans Only) | 12 to 20 fl. oz.<br>+<br>6 to 12 fl. oz.<br>+<br>0.5 oz.<br>or<br>0.75 oz.       | NIS at 0.25%<br>v/v plus COC at<br>0.125 to 0.25%<br>v/v<br>or<br>COC at 1 pt./A | AMS at 2.5<br>lbs/A | NIS at 0.25% v/v<br>plus COC at<br>0.25% v/v<br>or<br>COC at 1% v/v<br>(but not less than<br>1 pt./A) | 17 lbs/100 gals.<br>of<br>spray solution  |

1. Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
2. If grass regrowth occurs or an additional flush of new grass emerges, make a second application of **TAPOUT** alone (without a tank mix herbicide), according to the appropriate size and rate recommendations.
3. Annual grasses and sizes controlled with these tank mixtures are those that are identified in the **RECOMMENDATIONS FOR ANNUAL GRASSES** table.
4. Contact local Helena Chemical Company representative for proper COC/NIS adjuvant selection.
5. Refer to **FIRSTRATE**, **FLEXSTAR HL**, **FRONTROW**, **HARMONY GT XP**, **PURSUIT DG**, **RAPTOR**, **SYNCHRONY STS** and **SYNCHRONY XP (mp)** for geographic and rotational restrictions.
6. Always use a CPDA Certified Adjuvant when possible.

**SUGAR BEET**

**Table 11. TAPOUT TANK MIXED WITH BROADLEAF SUGAR BEET HERBICIDES**

| PRODUCTS          | APPLICATION RATES/A             | ADJUVANT INFORMATION |
|-------------------|---------------------------------|----------------------|
| TAPOUT            | 9 to 12 fl. oz.                 |                      |
| +<br>BETAMIX®     | See label for rate information. | None required        |
| or<br>BETANEX®    | See label for rate information. | None required        |
| or<br>PROGRESS®   | See label for rate information. | None required        |
| and/or<br>STINGER | See label for rate information. | See below            |
| and/or<br>UPBEET® | See label for rate information. | See below            |

NIS at 0.25% unless BETAMIX, BETANEX or PROGRESS is in the tank, then use no adjuvant.

**TABLE 12. TAPOUT PLUS BETANEX OR BETAMIX TANK MIX FOR THREE SEQUENTIAL APPLICATIONS FOR ANNUAL GRASS CONTROL (MICRO RATE APPLICATION)**

| PRODUCT        | APPLICATION RATES/ACRE <sup>(1)</sup> | GRASSES CONTROLLED (inches)  | METHYLATED SEED OIL <sup>(2)</sup> |          |
|----------------|---------------------------------------|--|------------------------------------|----------|
|                | ANNUAL GRASSES                        |  | GROUND                             | AIR      |
| TAPOUT         | 3 to 6 fl. oz.                        | Green Foxtail (1 to 2)<br>Yellow Foxtail (1 to 2)<br>Barnyardgrass (1 to 2)<br>Wild Oat (1 to 2)<br>Volunteer Cereals (1 to 2) | 1.5% v/v                           | 1.5% v/v |
| +<br>BETANEX   | Refer to label                        |  |                                    |          |
| or<br>BETAMIX  | Refer to label                        |  |                                    |          |
| or<br>PROGRESS | Refer to label                        |  |                                    |          |
| or<br>STINGER  | Refer to label                        |  |                                    |          |
| or<br>UPBEET   | Refer to label                        |  |                                    |          |

- 1) Broadleaf weed control may be reduced when grass populations are tall or dense enough to intercept the spray pattern and prevent them from receiving complete coverage. Tank mixing is not specified in these situations.
- 2) Always use a methylated seed oil at the listed rate (but not less than 1 pt/A) in the finished spray volume.
- 3) Always use a CPDA Certified Adjuvant when possible.

**Directions for Use for Micro-Rate Applications to Sugar Beet**

Multiple micro-rate applications of TAPOUT in tank mixtures with reduced rates of BETANEX or BETAMIX and methylated seed oils may be applied by air or ground equipment to sugar beet to control early germinating annual grasses listed above. All use precautions and restrictions on the BETANEX and BETAMIX master labels must be followed.

**Table 13. TANK MIX APPLICATION OF TAPOUT AND FUNGICIDES FOR CONTROL OF GRASS WEEDS AND DISEASES IN SUGAR BEET**

| PRODUCT <sup>(1)</sup> | APPLICATION RATES/ACRE <sup>(2)</sup> |                   | ADJUVANT         |
|------------------------|---------------------------------------|-------------------|------------------|
|                        | ANNUAL GRASSES                        | PERENNIAL GRASSES |                  |
| TAPOUT                 | 9 to 12 fl. oz.                       | 12 to 24 fl. oz.  | NIS at 0.25% v/  |
| +<br>EMINENT®          | Refer to label                        | Refer to label    |                  |
| TAPOUT                 | 9 to 12 fl. oz.                       | 12 to 24 fl. oz.  | NIS at 0.25% v/v |
| +<br>HEADLINE®         | Refer to label                        | Refer to label    |                  |
| TAPOUT                 | 9 to 12 fl. oz.                       | 12 to 24 fl. oz.  | NIS at 0.25% v/v |
| +<br>GEM™              | Refer to label                        | Refer to label    |                  |

- 1) Refer to TAPOUT and fungicide label for rates and weeds and diseases controlled.

- 2) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of **TAPOUT** alone (without a tank mix fungicide) according to the appropriate size and rate recommendations.
- 3) Always use a CPDA Certified Adjuvant when possible.

**Table 14. TANK MIX APPLICATION OF TAPOUT AND INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN ALFALFA, COTTON, MINT, PEANUT (INCLUDING PERENNIAL), SOYBEAN AND SUNFLOWER**

| PRODUCT <sup>(1)</sup>                                  | APPLICATION RATES/ACRE <sup>(2)</sup>                            |   | ADJUVANT RECOMMENDATION                    | CROPS                  |        |                       |        |         |           |
|---|--|---|--|------------------------|--------|-----------------------|--------|---------|-----------|
|   | ANNUAL GRASSES   | PERENNIAL GRASSES   |  | Alfalfa <sup>(3)</sup> | Cotton | Mint <sup>(3,4)</sup> | Peanut | Soybean | Sunflower |
| <b>TAPOUT</b><br>+<br>Asana® XL                         | 9 to 12 fl. oz.<br>+<br>Refer to Asana XL label                  | 12 to 24 fl. oz.<br>+<br>Refer to Asana XL label                  | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A |                        |        |                       |        | X       | X         |
| <b>TAPOUT</b><br>+<br>Baythroid®                        | 9 to 12 fl. oz.<br>+<br>Refer to Baythroid label                 | 12 to 24 fl. oz.<br>+<br>Refer to Baythroid label                 | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A | X                      |        |                       |        | X       |           |
| <b>TAPOUT</b><br>+<br>Danitol® 2.4 EC                   | 9 to 12 fl. oz.<br>+<br>10 2/3 to 16 fl. oz                      | 12 to 24 fl. oz.<br>+<br>10 2/3 to 16 fl. oz                      | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs/A  |                        | X      |                       | X      |         |           |
| <b>TAPOUT</b><br>+<br>Dimethoate                        | 9 to 12 fl. oz.<br>+<br>Refer to Dimethoate label                | 12 to 24 fl. oz.<br>+<br>Refer to Dimethoate label                | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A | X                      |        |                       |        | X       |           |
| <b>TAPOUT</b><br>+<br>Lorsban®                          | 9 to 12 fl. oz.<br>+<br>Refer to Lorsban label                   | 12 to 24 fl. oz.<br>+<br>Refer to Lorsban label                   | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A | X                      |        |                       |        | X       |           |
| <b>TAPOUT</b><br>+<br>Orthene® 75 S<br>or<br>Orthene 97 | 9 to 12 fl. oz.<br>+<br>0.33 to 1.33 lbs.<br>or<br>0.25 to 1 lb. | 12 to 24 fl. oz.<br>+<br>0.33 to 1.33 lbs.<br>or<br>0.25 to 1 lb. | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A |                        | X      | X                     | X      | X       |           |
| <b>TAPOUT</b><br>+<br>Orthene 90 S                      | 9 to 12 fl. oz.<br>+<br>0.25 to 1 lb.                            | 12 to 24 fl. oz.<br>+<br>0.25 to 1 lb.                            | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A |                        | X      | X                     | X      | X       |           |
| <b>TAPOUT</b><br>+<br>Pounce®                           | 9 to 12 fl. oz.<br>+<br>Refer to Pounce label                    | 12 to 24 fl. oz.<br>+<br>Refer to Pounce label                    | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs/A  | X                      |        | X                     |        |         |           |
| <b>TAPOUT</b><br>+<br>Warrior®                          | 9 to 12 fl. oz.<br>+<br>Refer to Warrior label                   | 12 to 24 fl. oz.<br>+<br>Refer to Warrior label                   | NIS at 0.25% v/v<br>+<br>AMS at 2.5 lbs./A |                        |        | X                     | X      |         |           |

- 1) Refer to **TAPOUT** and insecticide label for rates and weeds and insects controlled.
- 2) If grass regrowth occurs, or an additional flush of new grass emerges, make a second application of **TAPOUT** alone (without a tank mix insecticide) according to the appropriate size and rate recommendations.
- 3) Certain insecticides may cause temporary phytotoxic symptoms on alfalfa and mint foliage. Refer to the insecticide label for further information. It is suggested that prior to using any of these insecticide/herbicide tank mixtures, that a small area of the field be treated first and observations for crop injury be made prior to treating the whole field.
- 4) The **TAPOUT** rate should be 9 to 12 fl. oz./A for annual grass control in baby mint, minimum of 12 fl. oz./A for annual grass control in established mint and 16 to 32 fl. oz./A for perennial grass control.
- 5) Always use a CPDA Certified Adjuvant when possible.

**FALLOW LAND**

**DIRECTIONS FOR USE**

**TAPOUT** may be used to control annual and perennial grasses in land that has been left fallow the previous year and other non-producing agricultural areas. Apply **TAPOUT** at 12 to 16 fl. oz./A for annual grasses and 16 to 32 fl. oz./A for perennial grasses. When both grass and broadleaf weeds are the target pest, **TAPOUT** may be tank mixed with 2,4-D ester or BANVEL® SGF for broad spectrum control. When both annual and perennial grasses occur in the same field, use a minimum of 16 fl oz./A **TAPOUT** rate.

• **PRODUCT INFORMATION:**

- Use a minimum spray volume of 5 gals./A for aerial applications and 15 gals./A for ground applications. Apply only to actively growing grasses when the first grass reaches the specified weed height as specified by the Directions for Annual and Perennial Grasses section of this label.
- Annual grasses that emerge after the **TAPOUT** application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than 1 application in non-tilled areas.
- Do not apply to grasses that have tillered, formed seedheads or exceeded specified growth stage.
- Do not use flood jet nozzles.
- Do not apply to drought stressed grasses.
- Do not mow area for 2 weeks prior to or after the **TAPOUT** application.

**TABLE 15. TAPOUT IN TANK MIXES TO CONTROL ANNUAL AND PERENNIAL GRASSES IN FALLOW LAND**

| PRODUCT   | APPLICATION RATES/ACRE <sup>(1)</sup>  |   | ADJUVANT RECOMMENDATION                                    |  |
|---|--|---|--|--|
|   | ANNUAL GRASSES   | PERENNIAL GRASSES   | GROUND   | AIR  |
| <b>TAPOUT</b><br>+<br>2,4-D ester<br>or<br>BANVEL SGF | 12 to 16 fl. oz.<br>+<br>0.5 lb./A<br>or<br>See BANVEL SGF<br>label for rates. | 16 to 32 fl. oz.<br>+<br>0.5 lb./A<br>or<br>See BANVEL SGF<br>label for rates | NIS at 0.25% v/v or<br>COC at 1%<br>+<br>AMS at 2.5 lbs./A | NIS at 0.25% v/v or<br>COC at 1%<br>+<br>AMS at 17 lbs./100<br>gals. |

- 1) Refer to **TAPOUT** label for weed height and species control. Review BANVEL SGF and 2,4-D labels for crop restrictions, use rates and weeds controlled.
- 2) Always use a CPDA Certified Adjuvant when possible.

| DIRECTIONS FOR GRASS SUPPRESSION IN NON-CROP AREAS WITH TAPOUT                            |   |                       |                        |
|---|---|-----------------------|------------------------|
| GRASS SPECIES   | WEED STAGE  | APPLICATION RATES     |                        |
|   |   | MINIMUM RATE fl. oz/A | MAXIMUM RATE fl. oz./A |
| Annual and perennial grasses that exceed height claimed for control on height chart above | Up to and including grasses in the seed head stage. | 24                    | 32                     |

- Do not apply as part of a tank mix when applying **TAPOUT** for grass suppression.
- Add a crop oil concentrate at 1 qt./A by ground to the finished spray volume.
- Always use a CPDA Certified Adjuvant when possible.

**TABLE 16. TAPOUT FOR THE CONTROL AND/OR SUPPRESSION OF TALL FESCUE IN NATIVE PRAIRIE WARM-SEASON GRASS RESTORATION PROJECTS**

| PRODUCT       | PRODUCT RATE       | GRASS WEEDS CONTROLLED/SUPPRESSED |                            | WEED STAGE                              |
|---------------|--------------------|-----------------------------------|----------------------------|---|
|               |                    | Common Name                       | Scientific Name            |   |
| <b>TAPOUT</b> | 12 to 16 fl. oz./A | Tall Fescue                       | <i>Festuca arundinacea</i> | 4 to 6 inches tall (40 to 60% green-up) |

**Adjuvant:** TAPOUT must be applied with non-ionic surfactant at 0.25% v/v, plus a spray grade ammonium sulfate at 2.5 to 4 lbs./A.

**Recommended Mixing Order:** Thoroughly mix spray grade ammonium sulfate in water, add TAPOUT, then add non-ionic surfactant.

Always use a CPDA Certified Adjuvant when possible.

**SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS**

Burn or mow fields a minimum of 3 weeks prior to application to remove excess crop residue. Apply in the spring, at 40 to 60% tall fescue green-up, prior to emergence of warm-season grasses. Do not mow area for 2 weeks after the **TAPOUT** application.

Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 PSI at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood jet nozzles.

Apply only to fields that have warm-season grasses established for 2 years. Applications of **TAPOUT** to emerged warm-season grasses may cause injury. Do not apply to warm-season grasses grown for seed.

Do not graze treated fields or feed treated forage and or hay to livestock.

**NOTE:** **TAPOUT** applications are most effective if applied when average nighttime temperatures are consistently greater than or equal to 47 degrees Fahrenheit.

**TABLE 17. TAPOUT FOR THE SUPPRESSION OF TALL FESCUE SEED-HEADS IN NON- PRODUCING AGRICULTURAL AREAS**

| PRODUCT       | PRODUCT RATES    | SUPPRESSION  | APPLICATION TIMING   |
|---------------|------------------|--|--|
| <b>TAPOUT</b> | 3 to 4 fl. oz./A | Tall Fescue Seed-Heads<br>( <i>Festuca arundinacea</i> ) | (50 to 90% Tall Fescue green-up in the spring) or 3 weeks prior to dormancy in the fall. |

**ADJUVANT:** **TAPOUT** must be applied with crop oil concentrate at 1 qt. /A, plus a spray grade ammonium sulfate at 2.5 to 4 lb. /A.

**Recommended Mixing Order:** Thoroughly mix spray grade ammonium sulfate in water, add **TAPOUT**, then add crop oil concentrate.

**Note:** Use crop oil concentrate at 2 pts. /A with fall applications.

**SPECIAL APPLICATION INSTRUCTIONS/PRECAUTIONS**

- Apply at 50 to 90% tall fescue green-up.
- Use the higher **TAPOUT** rate if less tall fescue green matter is present.
- Do not mow area for 2 weeks after the **TAPOUT** application.
- Apply in a minimum of 15 to 20 gals of water per acre at a spray pressure of 40 to 60 psi at the nozzle. Apply using flat fan or hollow cone nozzles. Do not use flood nozzles.
- 2,4-D ester, **TORDON® 22K**, **GRAZON® P+D** or **CROSSBOW®** maybe added to this tank mix for broadleaf control (see 2,4-D ester label for weeds controlled)
- Do not graze treated fields or feed treated forage and or hay to livestock.

**DIRECTIONS FOR USE IN ORNAMENTALS**

For ornamental plant uses, **TAPOUT** can be used to control labeled grass weeds in greenhouses, lathhouses, shadehouses, and around outdoor ornamentals, including nurseries, parks, roadside plantings, and structure landscapes.

**IMPORTANT**

**TAPOUT** successfully controls weeds in newly transplanted and established non-grassy ornamentals. Plant tolerance to **TAPOUT** at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of **TAPOUT** has investigated the safety factor to ornamental plants not listed on the label.

The following plants have shown a tolerance for TAPOUT applications:

**ORNAMENTAL TREES**

| COMMON NAME           | SCIENTIFIC NAME          | COMMON NAME         | SCIENTIFIC NAME                |
|-----------------------|--------------------------|---------------------|--------------------------------|
| Alder, Red            | <i>Alnus rubra</i>       | Goldon Chain Tree   | <i>Laburnum anagyroides</i>    |
| Ash                   | <i>Fraxinus</i> spp.     | Maples              | <i>Acer</i> spp.               |
| Basswood              | <i>Tiia</i> spp.         | Mulberry, White     | <i>Morus alba</i>              |
| Birch, European White | <i>Betula pendula</i>    | Oaks                | <i>Quercus</i> spp.            |
| Birch, River          | <i>Betula nigra</i>      | Olive, Wild         | <i>Elaea gnus angustifolia</i> |
| Birch, White          | <i>Betula papyrifera</i> | Redbud, Eastern     | <i>Cercis canadensis</i>       |
| Crabapple, Flowering  | <i>Malus halliana</i>    | Sweet Gum, American | <i>Liquidambar styraciflua</i> |
| Dogwood, Flowering    | <i>Cornus florida</i>    |                     |                                |

**GROUND COVERS**

| COMMON NAME        | SCIENTIFIC NAME               |
|--------------------|-------------------------------|
| Bugleweed, Carpet  | <i>Ajuga reptans</i>          |
| Ivy, English       | <i>Hedera helix</i>           |
| Japanese Spurge    | <i>Pachysandra terminalis</i> |
| Lilyturf           | <i>Liriope muscari</i>        |
| Moneywort          | <i>Lysimachia nummulana</i>   |
| Mondo Grass, White | <i>Ophiopogon jaburan</i>     |
| Mondo Grass Dwarf  | <i>Ophiopogo njaponicus</i>   |
| Periwinkle, Lesser | <i>Vinca minor</i>            |

**GARDEN FLOWERS AND PLANTS**

\*slight foliage or flower speckling has been observed on these species.

| COMMON NAME             | SCIENTIFIC NAME                | COMMON NAME     | SCIENTIFIC NAME              |
|-------------------------|--------------------------------|-----------------|------------------------------|
| Ageratum                | <i>Ageratum</i> spp.           | Jasmine Tobacco | <i>Nicotiana alata</i>       |
| Alyssum*, Sweet         | <i>Lobularia maritima</i>      | Loosestrife     | <i>Lythrum salicaria</i>     |
| Asparagus Fern          | <i>Asparagus setaceus</i>      | Marigold        | <i>Tagetes</i> spp.          |
| Bleeding Heart          | <i>Dicentra spectabilis</i>    | Partridgeberry  | <i>Mitchella repens</i>      |
| Cast Iron Plant         | <i>Aspidistra elatior</i>      | Petunia*        | <i>Petunia hybrida</i>       |
| Chrysanthemum           | <i>Chrysanthemum</i> spp.      | Phlox           | <i>Phlox</i> spp.            |
| Cinquefoil              | <i>Potentilla</i> spp.         | Pinks           | <i>Dianthus</i> spp.         |
| Coleus                  | <i>Coleus</i> spp.             | Portulaca       | <i>Portulaca grandiflora</i> |
| Coralbells              | <i>Heuchera san guinea</i>     | Salvia          | <i>Salvia</i> spp.           |
| Cranesbill              | <i>Geranium</i> spp.           | Saxifrage       | <i>Saxifraga</i> spp.        |
| Dahlia                  | <i>Dahlia</i> spp.             | Sedum           | <i>Sedum</i> spp.            |
| Daisy, Trailing African | <i>Osteospermum fruticosum</i> | Selloum         | <i>Philodendron selloum</i>  |
| Daylily                 | <i>Hemerocallis</i> spp.       | Snapdragon*     | <i>Antirrhinum majus</i>     |
| Dusty Miller            | <i>Seneclo cineraria</i>       | Sweet Flag      | <i>Acorus gramineus</i>      |
| Euonymus                | <i>Euonymus</i> spp.           | Tickseed        | <i>Coreopsis grandiflora</i> |
| Gaza nia                | <i>Gazania</i> spp.            | Touch-Me-Not    | <i>Impatiens</i> spp.        |
| Geranium, House         | <i>Pelargonium hortorum</i>    | Verbena         | <i>Verbena</i> spp.          |
| Heather, False          | <i>Cuphea hyssopifolia</i>     | Violet          | <i>Viola</i> spp.            |
| Hosta                   | <i>Hosta fortunei</i>          | Yarrow, Common  | <i>Achillea millefolium</i>  |
| Iris                    | <i>Iris</i> spp.               | Zinnia          | <i>Zinnia elecans</i>        |



**SHRUBS**

\*Slight foliage or flower speckling has been observed on these species.

| COMMON NAME        | SCIENTIFIC NAME             | COMMON NAME               | SCIENTIFIC NAME                    |
|--------------------|-----------------------------|---------------------------|------------------------------------|
| Abelia             | <i>Abelia</i> spp.          | Jasmine                   | <i>Jasminum</i> spp.               |
| Anise, Purple      | <i>Illicium floridanum</i>  | Jasmine, Asiatic          | <i>Trachelospermum asiaticum</i>   |
| Aucuba             | <i>Aucuba</i> spp.          | Jasmine, Star             | <i>Trachelospermum jasminoides</i> |
| Azalea*            | <i>Rhododendron</i> spp.    | Juniper                   | <i>Juniperus</i> spp.              |
| Bamboo             | <i>Bambusa</i> spp.         | Lantana                   | <i>Lantana</i> spp.                |
| Barberry, Japanese | <i>Berberis thunbergii</i>  | Nandina* Bamboo, Heavenly | <i>Nandinia domestica</i>          |
| Barberry, Magellan | <i>Berberis buxifolia</i>   | Oleander, Common          | <i>Nerium oleander</i>             |
| Bayberry           | <i>Myrica pensylvanica</i>  | Oregon Grape              | <i>Mahonia aquifolium</i>          |
| Bottlebrush        | <i>Callistemon citrinus</i> | Photinia                  | <i>Photinia</i> spp.               |
| Boxwood, Common    | <i>Buxus sempervirens</i>   | Pittosporum               | <i>Pittosporum</i> spp.            |
| Camellia, Common   | <i>Camellia japonica</i>    | Podocarpus                | <i>Podocarpus</i> spp.             |
| Candytuft          | <i>Iberis sempervirens</i>  | Privet                    | <i>Ligustrum</i> spp.              |
| Cleyera            | <i>Cleyera japonica</i>     | Pyracantha                | <i>Pyracantha</i> spp.             |
| Coralberry         | <i>Ardisia crenata</i>      | Rhododendron              | <i>Rhododendron</i> spp.           |
| Crape Myrtle       | <i>Lagerstroemia indica</i> | Rose                      | <i>Rosa</i> spp.                   |
| Coyote Brush       | <i>Baccharis pilularis</i>  | Spirea                    | <i>Spiraea bumalda</i>             |
| Fig, Creeping      | <i>Ficus pumila</i>         | Sweet Olive               | <i>Osmanthus fragrans</i>          |
| Gardenia           | <i>Gardenia</i> spp.        | Viburnum                  | <i>Viburnum</i> thus               |
| Holly              | <i>Ilex</i> spp.            | Wisteria                  | <i>Wisteria</i> spp.               |
| Honeysuckle        | <i>Lonicera</i> spp.        | Yellow Sage/Shrub         | <i>Lantana camara</i>              |
| Indian Hawthorn    | <i>Raphiolepis indica</i>   | Verbena                   |                                    |

**DIRECTIONS FOR ANNUAL GRASSES IN ORNAMENTALS**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

| GRASS SPECIES         | SCIENTIFIC NAME                 | WEED* HEIGHT (inches) | APPLICATION RATES                     |                             |
|-----------------------|---------------------------------|-----------------------|---------------------------------------|-----------------------------|
|                       |                                 |                       | MINIMUM RATE fl. oz./A <sup>(1)</sup> | MAXIMUM RATE <sup>(2)</sup> |
| Barnyardgrass         | <i>Echinochloa crus-galli</i>   | 2 to 8                | 12                                    | 32                          |
| Broadleaf Signaigrass | <i>Brachiaria platyphylla</i>   | 2 to 6                | 12                                    | 32                          |
| Brome                 |                                 |                       |                                       |                             |
| California            | <i>Bromus carinatus</i>         | 2 to 6                | 12                                    | 32                          |
| Cheat                 | <i>Bromus secalinus</i>         | 2 to 6                | 12                                    | 32                          |
| Downy                 | <i>Bromus tectorum</i>          | 2 to 6                | 12                                    | 32                          |
| Ripgut                | <i>Bromus diandrus</i>          | 2 to 6                | 12                                    | 32                          |
| Canarygrass           | <i>Phalaris canariensis</i>     | 1 to 4                | 12                                    | 32                          |
| Crabgrass             |                                 |                       |                                       |                             |
| Hairy                 | <i>Digitaria adscendens</i>     | 2 to 6**              | 12                                    | 32                          |
| Large                 | <i>Digitaria sanguinalis</i>    | 2 to 6**              | 12                                    | 32                          |
| Smooth                | <i>Digitaria ischaemum</i>      | 2 to 6**              | 12                                    | 32                          |
| Southern              | <i>Digitaria ciliaris</i>       | 2 to 6**              | 12                                    | 32                          |
| Crowfootgrass         | <i>Dactyloctenium aegyptium</i> | 2 to 6**              | 12                                    | 32                          |
| Fall Panicum          | <i>Panicum dichotomiflorum</i>  | 2 to 8                | 12                                    | 32                          |
| Field Sandbur         | <i>Cenchrus incertus</i>        | 2 to 6                | 12                                    | 32                          |
| Foxtail               |                                 |                       |                                       |                             |
| Giant                 | <i>Setaria faberi</i>           | 2 to 12               | 12                                    | 32                          |
| Green                 | <i>Setaria vi ridis</i>         | 2 to 8                | 12                                    | 32                          |

|                         |                                |          |    |    |
|-------------------------|--------------------------------|----------|----|----|
| Yellow                  | <i>Setaria glauca</i>          | 2 to 8   | 12 | 32 |
| Goosegrass              | <i>Eleusine indica</i>         | 2 to 6** | 12 | 32 |
| Itchgrass               | <i>Rottboellia cochin</i>      | 2 to 6   | 12 | 32 |
| Junglerice              | <i>Echinochloa colona</i>      | 2 to 6   | 12 | 32 |
| Lovegrass (Stinkgrass)  | <i>Era grostis cilianensis</i> | 2 to 6   | 12 | 32 |
| Rabbitsfootgrass        | <i>Polypogon monspeliensis</i> | 1 to 4   | 12 | 32 |
| Red Rice                | <i>Oryza sativa</i>            | 1 to 3   | 12 | 32 |
| Rygrass                 |                                |          |    |    |
| Hardy                   | <i>Lolium remotum</i>          | 2 to 6   | 12 | 32 |
| Italian                 | <i>Lolium multiflorum</i>      | 2 to 6   | 12 | 32 |
| Seedling Johnsongrass   | <i>Sorghum halepense</i>       | 4 to 10  | 12 | 32 |
| Shattercane             | <i>Sorghum bicolor</i>         | 6 to 18  | 12 | 32 |
| Southwestern Cupgrass   | <i>Eriochloa gracilis</i>      | 2 to 6   | 12 | 32 |
| Sprangletop             |                                |          |    |    |
| Amazon                  | <i>Leptochloa panicoides</i>   | 2 to 6   | 12 | 32 |
| Bearded                 | <i>Leptochloa fascicularis</i> | 2 to 6   | 12 | 32 |
| Mexican                 | <i>Leptochloa uninervia</i>    | 2 to 6   | 12 | 32 |
| Red                     | <i>Leptochloa fihiformis</i>   | 2 to 6   | 12 | 32 |
| Texas Panicum           | <i>Panicum texanum</i>         | 2 to 6   | 12 | 32 |
| Volunteer Cereals       |                                |          |    |    |
| Barley                  | <i>Hordeum vulgare</i>         | 2 to 6   | 12 | 32 |
| Oats                    | <i>Avena sativa</i>            | 2 to 6   | 12 | 32 |
| Rye                     | <i>Secale cereale</i>          | 2 to 6   | 12 | 32 |
| Wheat                   | <i>Trifolium aestivum</i>      | 2 to 6   | 12 | 32 |
| Volunteer Corn          | <i>Zea mays</i>                | 4 to 12  | 12 | 16 |
| Volunteer Corn          | <i>Zea mays</i>                | 12 to 24 | 12 | 32 |
| Volunteer Grain Sorghum | <i>Sorghum bicolor</i>         | 8 to 12  | 12 | 32 |
| Wild Oats               | <i>Avena fatua</i>             | 2 to 6   | 12 | 32 |
| Wild Proso Millet       | <i>Panicum miliaceum</i>       | 2 to 10  | 12 | 32 |
| Witchgrass              | <i>Panicum capilare</i>        | 2 to 8   | 12 | 32 |
| Woolly Cupgrass         | <i>Eriochloa villosa</i>       | 2 to 8   | 12 | 32 |

\*Generally occurs between 3-leaf stage and tillering.

\*\*Length of lateral growth.

(1) 16 fl. oz./A = approximately 0.4 fl. oz./1000 sq. ft.

(2) 32 fl. oz./A = approximately 0.8 fl. oz./1000 sq. ft.

Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Always use a CPDA Certified Adjuvant when possible.

| DIRECTIONS FOR ANNUAL BLUEGRASS CONTROL WITH TAPOUT IN ORNAMENTALS   |            |                           |                           |
|--|------------|---------------------------|---------------------------|
| GRASS SPECIES  | WEED STAGE | APPLICATION RATES         |                           |
|  |            | MINIMUM RATE<br>fl. oz./A | MAXIMUM RATE<br>fl. oz./A |
| Annual Bluegrass ( <i>Poa annua</i> )  | To 4-leaf  | 12                        | 32                        |
| Apply under favorable soil moisture and humidity that exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s). |            |                           |                           |
| Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.  |            |                           |                           |
| Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.  |            |                           |                           |
| Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v).  |            |                           |                           |

43/45

**DIRECTIONS FOR PERENNIAL GRASSES IN ORNAMENTALS**

- Apply only to actively growing grasses at specified weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the specified growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

| GRASS SPECIES                                     | WEED HEIGHT<br>(inches) | APPLICATION RATES                           |   |
|---|-------------------------|---|---|
|   |                         | MINIMUM<br>RATE<br>fl. oz./A <sup>(1)</sup> | MAXIMUM<br>RATE <sup>(2)</sup><br>fl. oz./A |
| Bermudagrass ( <i>Cynodon dactylon</i> )          | 3 (or up to 6' runners) | 12  | 32  |
| First Application                                 | 3 (or up to 6' runners) | 12  | 32  |
| Repeat Application(s) (if regrowth occurs)        |                         |   |   |
| Foxtail Barley ( <i>Hordeum jubatum</i> )         |                         |   |   |
| First Application                                 | 2 to 6                  | 12  | 32  |
| Repeat Application(s) (if regrowth occurs)        | 2 to 6                  | 12  | 32  |
| Quackgrass ( <i>Elytrigia repens</i> )            |                         |   |   |
| First Application                                 | 4 to 8                  | 12  | 32  |
| Repeat Application(s) (if regrowth occurs)        | 4 to 8                  | 12  | 32  |
| Rhizome Johnsongrass ( <i>Sorghum halepense</i> ) |                         |   |   |
| First Application                                 | 12 to 24                | 2   | 32  |
| Repeat Application(s) (if regrowth occurs)        | 6 to 18                 | 9   | 16  |
| Wirestem Muhly ( <i>Muhlenbergia frondosa</i> )   |                         |   |   |
| First Application                                 | 4 to 8                  | 12  | 32  |
| Repeat Application(s) (if regrowth occurs)        | 4 to 8                  | 12  | 32  |

<sup>(1)</sup>16 fl. oz./A = approximately 0.3 fl. oz./1000 sq. ft.

<sup>(2)</sup>32 fl. oz./A = approximately 0.8 fl. oz./1000 sq. ft.

Add non-ionic surfactant containing at least 80% active ingredient at the rate of 1 pt. per 50 gals. (0.25% v/v). Always use a CPDA Certified Adjuvant when possible.

**IMPORTANT**

Plant tolerance to **TAPOUT** at labeled rates has been found to be acceptable for the indicated genera and species listed below. Due to variability within species, crop growth stage, environmental conditions, and application techniques, it is recommended that the user determine if the herbicide can be used safely on a few plants prior to widespread application. Neither the seller nor the manufacturer of **TAPOUT** have investigated the safety factor to plants not listed on the label.

**NON-BEARING FOOD CROPS**

**TAPOUT SHOULD NOT BE APPLIED TO NON-BEARING FRUIT OR NUT CROPS WHICH ARE GROWN FOR ROOT STOCK.**

Crop injury to non-bearing fruit and nut crops can occur if **TAPOUT** is improperly applied. **TAPOUT** should not be applied directly over the top of these plant types. Instead spray should be directed at the base of the plant where grassy weeds are growing near the ground.

Non-bearing fruit and nut crops are plants which will not bear fruit or nuts for at least one year following **TAPOUT** application.

| COMMON NAME   | SCIENTIFIC NAME                            |
|---------------|--|
| Apples        | <i>Milus spp</i>                           |
| Berries       | <i>Vaccinium spp.</i><br><i>Rubus spp.</i> |
| Cherry, Sweet | <i>Prunus avium</i>                        |
| Citrus Fruits | <i>Citrus spp</i>                          |
| Grapes        | <i>Vitis spp</i>                           |
| Olives        | <i>Olea spp</i>                            |
| Peach         | <i>Prunus persica</i>                      |
| Pears         | <i>Pyrus communis</i>                      |
| Prunes        | <i>Prunus spp</i>                          |
| Stone Fruits  | <i>Prunus spp</i>                          |
| Strawberries  | <i>Fragaria spp.</i>                       |
| Tree Nuts     |  |
| Almond        | <i>Prunus triloba</i>                      |
| Filbert       | <i>Coryluss maxima</i>                     |
| Pecan         | <i>Carya illiinoensis</i>                  |
| Pistachio     | <i>Pistacia vera</i>                       |
| Walnut        | <i>Juglans spp.</i>                        |

**CONIFER TREES**

TAPOUT can be used to control labeled grasses in Christmas tree farms, conifer nurseries and conifer plantations (but not in forests).

| COMMON NAME               | SCIENTIFIC NAME              |
|---------------------------|------------------------------|
| Arborvitae, American      | <i>Thuja occidentalis</i>    |
| Cedars                    | <i>Cedrus spp.</i>           |
| Cypress                   | <i>Taxodium spp.</i>         |
| Fir, Douglas              | <i>Pseudotsuga menziesll</i> |
| Firs                      | <i>Abiesspp</i>              |
| Hemlock, Canadian/Eastern | <i>Tsuga canadensis</i>      |
| Hemlock, Western          | <i>Tsuga heterophylla</i>    |
| Pines                     | <i>Pinus spp</i>             |
| Spruces                   | <i>Picea spp.</i>            |
| Yew                       | <i>Taxus spp.</i>            |

**NON-CROP OR NON-PLANTED AREAS**

The following areas are considered non-crop or non-planted areas: Rights-of-way including railroads, highways, roads, dividers, medians, pipelines, public utility lines, pumping stations, transformer stations and substations. Around airports, electric utilities, commercial buildings, manufacturing plants, storage yards, rail yards, fence lines, parkways, and post-harvest croplands. Also beneath greenhouse benches and around golf courses.

**CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES**

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

45/45

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

*Cobra, Phoenix and Resource* are trademarks and registered trademarks of Valent U.S.A. Corporation  
*Danitol* is a registered trademark of Sumitomo Chemical Co., Ltd.  
*Orthene* is a registered trademark of OMS Investments, Inc. *Orthene 97* is U.S. Patent 6,013,272  
*Asana, Harmony, Synchrony, STS and UpBeet* are registered trademarks of E.I. du Pont de Nemours and Company  
*Banvel, Clearfield, Headline, Pursuit and Raptor* are registered trademarks of BASF Corporation  
*Basagran* is a registered trademark of BASF AG  
*Baythroid* is a registered trademark of Bayer Corporation  
*Betamix, Betanex, Bucril, Gem, Liberty, Liberty Link and Progress* are registered trademarks of Bayer  
*Bronate and Bronate Advanced* are registered trademarks of Bayer Crop Science  
*Crossbow, Curtail, FirstRate, Frontrow, Grazon, Lorsban, Stinger and Tordon* are registered trademarks of Dow AgroSciences LLC  
*Eminent* is a registered trademark of ISAGRO S.P.A.  
*Flexstar and Warrior* are registered trademarks of Syngenta Corporation  
*Pounce* is a registered trademark of FMC Corporation  
*Roundup Ready* is a registered trademark of Monsanto Company  
*Ultra Blazer and Storm* are registered trademarks of United Phosphorus, Inc.  
*TAPOUT* is a registered trademark of Helena Holding Company.  
*Danitol* is a restricted use pesticide

---

© Helena Holding Company, 2008.