



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 6, 2023

Bill Washburn
Registration Manager
Helena Agri-Enterprises, LLC
d/b/a Helena Chemical Company
225 Schilling Blvd., Suite 300
Collierville, TN 38017

Subject: Registration Review Label Mitigation for Permethrin
Product Name: PRO-MATE UL 4-4
EPA Registration Number: 5905-587
Application Date: 11/6/2020
Decision Number: 567740

Dear Bill Washburn:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

Page 2 of 2
EPA Reg. No. 5905-587
Decision No. 567740

approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Carolyn Smith by phone at (202) 566-2273, or via email at smith.carolyn@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kelly Sherman", with a long horizontal flourish extending to the right.

Kelly Sherman
Chief, Risk Management & Implementation Branch III
Pesticide Re-evaluation Division
Office of Pesticide Programs

Enclosure

PRO-MATE UL 4-4


For use only by federal, state, tribal or local government officials responsible for public health or vector control or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control applications or by persons under their direct supervision.

A Ready to Use, Quick Knockdown, Low Odor, Non-Corrosive Synergized Synthetic Pyrethroid for the control of Adult Mosquitoes, In Residential, Municipal and Recreational areas Provides quick knockdown and effective control of mosquitoes, flies, black flies, gnats, biting and non-biting midges, as well as other biting flies.

ACTIVE INGREDIENTS:

Permethrin 4.0%
Piperonyl Butoxide..... 4.0%

OTHER INGREDIENTS*: 92.0%
TOTAL 100.0%



*Contains Petroleum Distillate
Contains 0.2783 lb. Permethrin a.i. and 0.2783 lb. Piperonyl Butoxide a.i. per gallon.

KEEP OUT OF REACH OF CHILDREN	
CAUTION	
PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.	
FIRST AID	
If swallowed	<ul style="list-style-type: none"> Immediately call a Poison Control Center or physician for treatment advice. Do not induce vomiting unless told to do so by the Poison Control Center or physician. Do not give any liquid to the person. Do not give anything by mouth to an unconscious or convulsing person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or physician for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or physician for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a Poison Control Center or physician for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
For Chemical Spill, Leak, Fire or Exposure, Call: CHEMTREC 1-800-424-9300	
NOTE TO PHYSICIAN: Contains Petroleum Distillates – vomiting may cause aspiration pneumonia.	
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS	

EPA Reg. No. 5905-587
EPA Est. No.

NET CONTENTS:
AD 061912

Manufactured for
Helena Agri-Enterprises, LLC
225 Schilling Boulevard, Suite 300
Collierville, TN 38017

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. Mixers, loaders, applicators and other handlers (including during mixing/loading, repair and cleaning of equipment and disposal of pesticide) must wear the following:

- Long-sleeved shirt and long pants
- Shoes plus socks,
- Chemical-resistant gloves made of barrier laminate, viton, neoprene rubber or nitrile rubber.
- Chemical-resistant apron for mixers/loaders, persons cleaning equipment and persons exposed to the concentrate.

In addition to the above PPE, mixers and loaders supporting aerial applications to forests must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

See engineering controls for additional requirements.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls:

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

User Safety Recommendations-

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to fish and aquatic invertebrates. Before making the first application in a season, it is advisable to consult with the state or tribal agency with primary responsibility for pesticide regulation to determine if permits or other regulatory requirements exist.

Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material away from the water in order to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or washwaters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use only by federal, state, tribal or local government officials responsible for public health or vector control or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control applications or by persons under their direct supervision.

RESISTANCE MANAGEMENT

For resistance management, Pro-Mate UL 4-4 contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to Pro-Mate UL 4-4 and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of Pro-Mate UL 4-4 or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Helena Agri-Enterprises at www.helenaagri.com

MANDATORY SPRAY DRIFT MANGEMENT

- Apply when ground wind speeds are equal to or greater than 1 mph.
- All types of applications should be conducted when temperatures at ground level are at or above 50°F.

For Ground Applications:

- Create an optimum swath when possible. An optimum swath width can be achieved when Pro-Mate UL 4-4 is applied from a truck that is being driven perpendicular to the wind direction. Direct the spray head of equipment to ensure even distribution of the spray cloud throughout the area.
- FOR BEST RESULTS treat when mosquitoes or insects are most active and weather conditions are conducive to keeping the spray cloud in the air column close to the ground.
- An inversion of air temperatures and a light breeze is preferable. Application during the cooler hours of the night or early morning is recommended.

For Aerial Applications:

- Do not apply by fixed wing aircraft at a nozzle height less than 100 feet (30.5 m) above ground or canopy, or by helicopter at a height less than 75 feet (22.9 m) above the ground or canopy, unless specifically approved by the state or tribe based on public health needs.

Adult mosquito control applications should be limited to trained personnel.

- For use only by federal, state, tribal or local government officials responsible for public health or vector control or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory Agency to perform adult mosquito control applications, or by persons under their supervision, or as allowed by state regulations for persons treating private property.
- This pesticide is toxic to aquatic organisms. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to aquatic organisms.
- Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material beyond the body of water to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or wash waters.
- Before making the first application in a season, it is advisable to consult with the state or tribal agency with primary responsibility for pesticide regulation to determine if other regulatory requirements exist.
- Do not treat a site with more than 0.007 pounds of permethrin or 0.007 pounds piperonyl butoxide of each a.i., per acre in a single application or in any 24-hour period. Do not exceed 0.18 pounds permethrin or 0.18 pounds piperonyl butoxide of a.i. in any site in one year. More frequent applications may be made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease-causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

Precautions and Use Restrictions

- Apply this product only as specified on this label.
- Apply this product at temperatures greater than 50°F and when wind speed equals or exceeds 1 mph.
- Not for use in metered release systems or animal premise automatic misting systems.
- Not for use in outdoor residential misting systems.
- Do not spray this product on or allow it to drift onto cropland (other than those listed) or potable water supplies.
- In the treatment of corrals, feedlots, animal confinements/houses, swine lots, poultry ranges and zoos, cover any exposed drinking water, drinking fountains and animal feed before application.
- For hand carried and/or lawn mower foggers: Do not fog wet foliage. Do not fog when wind conditions exceed 10 miles per hour. Only use a clean, well maintained and properly calibrated fogger, known to give excellent performance. Fog down wind.
- Do not apply by fixed wing aircraft at a height less than 100 feet, or by helicopter at a height less than 75 feet unless specifically approved by the state or tribe based on public health needs.
- Apply when wind speed is greater than 1 mph.
- Do not make applications during rain.
- Except when applying to storm drains, drainage pipes and pipe chases as instructed on this label, do not apply directly to sewers and drains or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur.
- Do not use in dairy barns or facilities.

Do not retreat site more than once in 3 days. Do not exceed 25 applications at 0.007 lbs. permethrin per acre per application or 0.007 lbs piperonyl butoxide per acre per application, or 0.18 lbs. of permethrin per acre per season or 0.18 lbs piperonyl butoxide per acre per year. More frequent treatments may be made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

Pro-Mate UL 4-4 is recommended for application as an Ultra Low Volume (ULV) Thermal or Non-thermal aerosol (cold fog) for control of adult mosquitoes and flies in residential, recreational, industrial, urban and other areas where these insect pests are found. Pro-Mate UL 4-4 provides effective control of mosquitoes and flies in areas such as parks, campsites, woodlands, athletic fields, golf courses, residential areas, municipalities, roadsides, animal quarters, barns, playgrounds, and non- crop waste areas. Treat when mosquitoes or insects are most active and weather conditions are conducive to keeping the fog close to the ground, e.g. cool temperatures and ground wind speeds are not greater than 10 mph. Applications during the cool hours of night or early morning are usually preferable. Repeat treatment when recommended by Public Health officials, or Mosquito Abatement Districts.

Pro-Mate UL 4-4 may be applied over specific growing crops and range grasses prior to harvest for the control of adult mosquitoes and biting flies with or adjacent to those areas. Application can only be made where the following crops are present:

Alfalfa, Almonds, Apples, Artichoke, Asparagus, Avocado, Broccoli, Brussel sprouts, Cabbage, Cantaloupe, Cauliflower, Celery, Cherry, Corn (field and pop), Corn (sweet), Eggplant, Filberts (Hazelnuts), Garlic, Horseradish, Lettuce, Onions (dry bulb only), Papayas (FL only), Peaches/Nectarines, Pears, Bell peppers, Pistachios, Potatoes, Soybeans, Spinach, Tomatoes, and Walnuts.

Cucurbit vegetables, including Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of Cucumis melo including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of Citrullus lanatus).

Leafy Vegetables, including Leafy vegetables (except brassica) (Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Envide (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), , Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)).

Do not allow spray treatment to drift on cropland, (other than crops listed) or water supplies. In treatment of corrals, feedlots, animal confinements/houses, swine lots and zoos, cover any exposed drinking water, drinking fountains and animal feed before application

Spray Droplet Size Determination

Ground-Based Application: Spray equipment must be adjusted so that the volume median diameter is less than 30 microns ($Dv\ 0.5 < 30\ \mu m$) and that 90% of the spray is contained in droplets smaller than 50 microns ($Dv\ 0.9 < 50\ \mu m$). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using laser based equipment to produce acceptable droplet spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

Aerial Application made at ≤ 200 feet above ground elevation: Spray equipment must be adjusted so that the volume median diameter produced is less than 60 microns ($Dv\ 0.5 < 60\ \mu m$) and that 90% of the spray is contained in droplets smaller than 100 microns ($Dv\ 0.9 < 100\ \mu m$). The effects of flight speed and, for non-rotary nozzles, nozzle angle on the droplet size spectrum must be considered. Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

Aerial Application made at > 200 feet above ground elevation: Spray equipment must be adjusted so that the volume median diameter produced is less than 70 microns ($Dv\ 0.5 < 70\ \mu m$) and that 90% of the spray is contained in droplets smaller than 145 microns ($Dv\ 0.9 < 145\ \mu m$). The effects of flight speed and, for non-rotary nozzles, nozzle angle on the droplet size spectrum must be considered. Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

GROUND APPLICATION

ULV NON-THERMAL AEROSOL (COLD FOG): To control adult mosquitoes, flies, and gnats (blackflies) apply Pro-Mate UL 4-4 using truck mounted ground ULV (cold fog) equipment with a swath width of 300 feet in areas such as residential, recreational, industrial areas. Apply when insects are most active and conditions are conducive to keeping the fog close to the ground. Apply the undiluted product at flow rates of 4.092 to 19.092 fluid oz. per minute at a truck speed of 10 mph. These rates are equivalent to 0.001 to 0.007 pounds of permethrin a.i. and 0.001 to 0.007 pounds of piperonyl butoxide a.i. per acre. If a different vehicle speed is used, adjust the flow rate accordingly. An accurate flow control system must be used to ensure proper flow rate. Use of the maximum flow rates and application rates on this label to treat heavy vegetation or when target insect populations are high. Do not exceed the highest flow rates specified by the manufacturer. Pro-Mate UL 4-4 may also be applied by diluting with a light mineral oil or another suitable solvent, applied so as not to exceed 0.007 lbs. a.i. per acre using a swath width of 300 feet for calculation. Refer to Chart 2 for suggested rates for diluted use. Pro-Mate UL 4-4 may be used to treat outdoor aggregation and harborage sites including storm drains, underground drainage pipes, and pipe chases.

THERMAL AEROSOL APPLICATIONS: For use in thermal fogging equipment such as truck-mounted thermal fog generators or other equipment capable of making thermal fogging applications. Pro-Mate UL 4-4 may be applied either diluted or undiluted. If diluted, use a light mineral oil, or another suitable non-phytotoxic diluent. Do not exceed the maximum rate of 0.007 lbs. a.i. per acre. May be applied with truck-mounted equipment at speeds of 5 mph to 20 mph. Use a swath width of 300 feet for calculation. For use with handheld, backpack or lawn mower thermal fogging equipment do not exceed the maximum rate of 0.007 lbs. a.i. per acre. Use a walking speed of 2 mph (3.2 kph) and a swath width of 300 feet for calculations. May be used to treat aggregation and harborage sites

such as abandoned buildings, warehouses, storm drains, underground drainage pipes, pipe chases and other areas where adult mosquitoes find harborage.

BARRIER APPLICATIONS: For use in non-thermal ULV. motorized, portable, backpack, handheld or truck-mounted fogging equipment or mist blower equipment, apply Pro-Mate UL 4-4 with equipment adjusted to deliver ULV particles of 50 to 100 volume median diameter. Pro-Mate UL 4-4 may be applied undiluted or diluted (see Application Rates Charts on this label). If diluted, use a light mineral oil, or another suitable, non-phytotoxic diluent. For truck-mounted applications, use not more than 0.007 lbs. permethrin per acre. Use a 300 ft. (15.2 m) swath width (see application rate charts on this label) for calculations. For backpack or handheld applications, use not more than 0.007 lbs. a.i. per acre. Use a 300 ft. swath width and a walking speed of 2 mph (3.2 kph) for calculations. Direct spray onto foliage at the perimeter of areas from which mosquitoes or flies are to be excluded or direct spray into harborage areas where adult mosquitoes or flies may be found. May be used to treat aggregation and harborage sites such as abandoned buildings, residential buildings, warehouses, storm drains, underground drainage pipes and other areas where adult mosquitoes find harborage.

AERIAL APPLICATION

Use aircraft specially equipped and capable of applying Ultra-Low Volumes of Pro-Mate UL 4-4. Applications can be made diluted or undiluted but cannot exceed the specified rates of 0.007 lbs. permethrin per acre and 0.007 lbs. piperonyl butoxide per acre, equivalent to 3.1819 fl. oz. of undiluted Pro-Mate UL 4-4 per acre. Applications should be made at an altitude of 100 to 300 feet. Apply at airspeed sufficient to deliver the appropriate amount of active ingredient per acre. The minimum height (from the ground) for aerial application of this product is 100 feet. Treat when insects are most active and meteorological conditions are conducive to keeping the spray cloud close to the ground. In order to compensate for windy conditions and ensure drift onto the target area aerial application with aircraft equipped with Global Positioning Systems (GPS) is recommended.

PROHIBITION ON AERIAL USE IN FLORIDA: Not for aerial application in the State of Florida UNLESS specifically authorized by the Bureau of Entomology and Pest Control, Florida Department of Agriculture and Consumer Services.

CHART NO. 1 – FOR USE UNDILUTED

Conditions and Rates To Use Pro-Mate UL 4-4 Undiluted for Mosquito Control						
Pounds Per Acre		Application Rates				Fluid Ounces and Milliliters Per Acre
		Fluid Ounces per Minute				
Permethrin	PBO	5 mph	10 mph	15 mph	20 mph	
0.007	0.007	9.546	19.092	28.637	38.183	3.1819(94.1 ml)
0.0035	0.0035	4.773	9.545	14.318	19.091	1.5909 (47.05 ml)
0.001	0.001	1.363	2.728	4.092	5.455	0.4546 (13.44 ml)

CHART NO. 2 – DILUTE 1 PART WITH 2 PARTS MINERAL OIL

Amounts to Use After Diluting 1 Part Pro-Mate UL 4-4 with 2 Parts Mineral Oil						
Lbs. of A.I. Per Acre		Application Rates for Fluid Ounces of Diluted Product				Diluted Fl. Oz. (ml) to Apply Per Acre
		Per Minute Per MPH				
Permethrin	PBO	5 mph	10 mph	15 mph	20 mph	
0.007	0.007	26.250	52.50	78.75	105.00	8.75 (258.76 ml)
0.0035	0.0035	3.125	26.25	39.38	52.50	4.38 (129.53 ml)
0.0010	0.0010	3.750	7.50	11.25	15.00	1.25 (36.96 ml)

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original labeled container in a cool, dry, locked place designated for such insecticides and out of reach of children. In case of spillage, soak up with absorbent material such as sawdust, or fuller's earth, sweep up and place in a labeled container and dispose of as follows.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not refill or reuse this container. Clean container promptly after emptying. [For Containers 5 gallons or less] Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with mineral oil (or other solvent used when diluting the product) and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. [For Containers greater than 5 gallons] Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with mineral oil (or other solvent used when diluting the product). Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Helena Agri-Enterprises, LLC (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Agri-Enterprises, LLC's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.