



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505T)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number: **5905-670** Date of Issuance: **2/11/26**

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Omni Brand Lambda-Cy 250 CS

**Name and Address of Registrant (include ZIP Code):**

Helena Agri-Enterprises, LLC d/b/a Helena Chemical Company  
225 Schilling Boulevard, Suite 300  
Collierville, Tennessee 38017

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

*Continues page 2*

**Signature of Approving Official:**

Date:

**2/11/26**

Jacquelyn Herrick, Product Manager 03  
IVB1, Registration Division (7505T)

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 5905-670."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 05/28/2024

If you have any questions, please contact Hester Dingle at 202-566-2596 or at [dingle.hester@epa.gov](mailto:dingle.hester@epa.gov).

Enclosure

**RESTRICTED USE PESTICIDE  
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS**

For retail sale to and use only by certified applicators, or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

**Lambda-Cyhalothrin** **Group** **3A** **Insecticide**

# **OMNI BRAND LAMBDA-CY 250 CS**

**Active Ingredient:**

**(% BY WT)**

Lambda-Cyhalothrin <sup>1</sup>:

[1α(S\*)3α(Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate..... 23.7%

Other Ingredients: ..... 76.3%

Total: ..... 100.0%

<sup>1</sup> Synthetic pyrethroid

Contains petroleum distillate.

**Omni Brand Lambda-Cy 250 CS** contains 2.08 lbs of active ingredient per gallon or 250 grams of active ingredient per liter and is a capsule suspension.

**KEEP OUT OF REACH OF CHILDREN  
WARNING / AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

**FIRST AID**

If swallowed	<ul style="list-style-type: none"><li>Immediately call a poison control center or doctor.</li><li><b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li><b>DO NOT</b> give any liquid to the person.</li><li><b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
If on skin or clothing	<ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
If in eyes	<ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>Call a poison control center or doctor for treatment advice</li></ul>
If inhaled	<ul style="list-style-type: none"><li>Move person to fresh air.</li><li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>Call a poison control center or doctor for treatment advice</li></ul>

**NOTE TO PHYSICIAN**

Contains petroleum distillate - vomiting may cause aspiration pneumonia.


**HOT LINE NUMBER**

In case of an emergency involving this product and for 24-hour medical emergency assistance (human or animal) and chemical emergency assistance (spill, leak, fire, or accident), call CHEMTREC at 1-800-424-9300.

[See] Additional Precautions and Directions for Use inside Product Booklet

EPA Reg. No. 5905-X

EPA Est. No. \_\_\_\_\_



NET CONTENTS: \_\_\_\_\_

Manufactured for  
Helena Agri-Enterprises, LLC  
225 Schilling Boulevard, Suite 300  
Collierville, TN 38017

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**  
**WARNING / AVISO**

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if absorbed through skin or inhaled. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Avoid breathing vapor or spray mist. Wear protective clothing, gloves, eyewear (goggles, face shield, or safety glasses) as indicated under **Personal Protective Equipment**. Remove and wash contaminated clothing before reuse.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hrs. after exposure and may last 2 – 30 hrs., without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

**Personal Protective Equipment (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of barrier laminate, nitrile rubber ( $\geq 14$  mls), neoprene rubber ( $\geq 14$  mls), or Viton ( $\geq 14$  mls)
- Shoes plus socks
- Protective eyewear

In addition to the above, for foliar broadcast spray treatment using a mechanically pressurized handgun in nurseries (conifer and deciduous trees), mixers, loaders, and applicators must also wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 lb ai per acre, and treating 1200 acres (or more) per day, must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

**Environmental Hazards**

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife.

For terrestrial uses: **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

## NON-TARGET ORGANISM ADVISORY STATEMENT

This product is highly toxic to bees and other pollinating insects exposed to direct treatment or residues in blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.**

### Physical and Chemical Hazards

Combustible liquid. **DO NOT** use or store near heat or open flame. **DO NOT** mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

### RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

**DO NOT ALLOW PRODUCT TO FREEZE.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours for the following activities:

- Hand detasseling or mechanically assisted detasseling of field corn grown for seed.
- Hand detasseling or mechanically assisted detasseling of pop corn grown for seed.
- Hand detasseling or mechanically assisted detasseling of sweet corn grown for seed.
- Hard harvesting of sweet corn grown for grain.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ( $\geq 14$  mls), neoprene rubber ( $\geq 14$  mls), or Viton ( $\geq 14$  mls)
- Shoes plus socks
- Protective eyewear

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

## USE DIRECTIONS

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gal. per acre by air or 10 gal. per acre by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher label rates may improve initial and residual control. Remove chemical extraction probes (also known as "stingers") used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

For cutworm control, **Omni Brand Lambda-Cy 250 CS** may be applied before, during, or after planting. For soil incorporated applications, use higher label rates for improved control.

## USE RESTRICTIONS

**DO NOT** apply as foliar broadcast application using a mechanically pressurized handgun on: Brassica (head and stem), Cucurbit Vegetables, Fruiting Vegetables, Garlic, Legume Vegetables, Lettuce (head and leaf), Onion (dry bulb), Tobacco, Tuberous and Corm Vegetables.

## RESISTANCE MANAGEMENT

For resistance management, **Omni Brand Lambda-Cy 250 CS** contains a Group 3A insecticide/acaricide. Any insect population may contain individuals naturally resistant to **Omni Brand Lambda-Cy 250 CS** and other Group 3A insecticides/acaricides. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Rotate the use of **Omni Brand Lambda-Cy 250 CS** or other Group 3A insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Helena Agri-Enterprises, LLC at 901-761-0050.

## MANDATORY SPRAY DRIFT MANAGEMENT

### Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets (ASABE

S641).

- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use  $\frac{3}{4}$  swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

#### **Airblast Applications:**

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

#### **Ground Boom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a Medium or coarser droplet size (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

#### **Boomless Ground Applications:**

- Applicators are required to select nozzle and pressure that deliver a Medium or coarser droplet size (ASABE S572) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.  
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

#### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

#### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

#### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

#### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **HANDHELD TECHNOLOGY APPLICATIONS**

Take precautions to minimize spray drift.

### **VEGETATIVE FILTER STRIPS**

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing **Omni Brand Lambda-Cy 250 CS** onto fields where a maintained vegetative filter strip of at least 25 feet exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
  - For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
  - The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
  - Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
  - A functional terrace system is maintained on the area of application.
  - Water and sediment control basins for the area of application are functional and maintained.
  - The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers:

*Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services.*

<https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>

### **Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)**

**DO NOT** apply within 25 feet of aquatic habitats (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

### **Buffer Zone for ULV Aerial Application**

**DO NOT** apply within 450 feet of aquatic habitats (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

### **Buffer Zone for Non-ULV Aerial Application**

**DO NOT** apply within 150 feet of aquatic habitats (including, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

In New York State, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh, or stream that drains into a coastal salt marsh, for both aerial and ground application. For aerial applications, the 25 ft. vegetated, non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

## CHEMIGATION

### Sprinkler Irrigation Application

Only apply **Omni Brand Lambda-Cy 250 CS** at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations (within the confines of this label) on adjuvant or diluent types, rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with **Omni Brand Lambda-Cy 250 CS** applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period:

Apply by injecting the specified rate of **Omni Brand Lambda-Cy 250 CS** into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1 - 0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to ensure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of **Omni Brand Lambda-Cy 250 CS** for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that **Omni Brand Lambda-Cy 250 CS** be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption, if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Removable chemical extraction probes (also known as "stingers") used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

### Use Precautions: Sprinkler Irrigation Application

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). **DO NOT** apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label- prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve. Vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water- source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick- closing check valve to prevent the

flow of fluid back toward the injection pump.

- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Any alternatives to the above required safety devices must conform to the list of EPA-approved alternative devices.
- M. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment or non-uniform distribution of treated water.
- N. **DO NOT** apply through chemigation systems connected to public water systems.

Following best management practices can help reduce risk to terrestrial pollinators. Examples of best management practices include applying pesticides in the evening and at night when pollinators are not foraging and checking to confirm hive locations before spraying. For additional resources on pollinator best management practices, visit <https://www.epa.gov/pollinator-protection/find-best-management-practices-protect-pollinators>.

**Managed pollinator protection plans** are developed by states/tribes to promote communication between growers, landowners, farmers, beekeepers, pesticide users, and other pest management professionals to reduce exposure of bees to pesticides. If available, visit state plans for additional information on how to protect pollinators.

#### How to Report Bee Kills

It is recommended that users contact both the state lead agency and the U.S. Environmental Protection Agency to report bee kills due to pesticide application. Bee kills can be reported to EPA at [beekill@epa.gov](mailto:beekill@epa.gov). To contact your state lead agency, see the current listing of state pesticide regulatory agencies at the National Pesticide Information Center's website: [http://npic.orst.edu/reg/state\\_agencies.html](http://npic.orst.edu/reg/state_agencies.html).

#### Rate Conversion Chart

Lb. A.I. Per Acre	Fl. Oz. Per Acre	Pints Per Acre	Treated Acres Per Gallon
0.015	0.96	0.06	133
0.02	1.28	0.08	100
0.025	1.60	0.10	80
0.03	1.92	0.12	67
0.035	2.24	0.14	57
0.04	2.56	0.16	50

**SPECIFIC USE DIRECTIONS  
AGRICULTURAL USES**

**ALFALFA AND ALFALFA GROWN FOR SEED**

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Alfalfa Caterpillar		
Army Cutworm		
Cutworm species		
Green Cloverworm		
Leafhopper species	0.015 - 0.025	0.96 - 1.60
Looper species		
Three-cornered Alfalfa Hopper		
Velvetbean Caterpillar		
Webworm species		
Alfalfa Seed Chalcid (Adult)		
Alfalfa Weevil		
Armyworm		
Bean Leaf Beetle (Adult)		
Blister Beetle species		
Blue Alfalfa Aphid		
Clover Leaf Weevil species		
Clover Root Borer (Adult)		
Clover Root Curculio species (Adult)		
Clover Stem Borer (Adult)		
Corn Earworm		
Cowpea Aphid		
Cowpea Curculio (Adult)		
Cowpea Weevil (Adult)		
Cucumber Beetle species (Adult)		
Egyptian Alfalfa Weevil		
Fall Armyworm <sup>1</sup>	0.02 - 0.03	1.28 - 1.92
Grape Colaspis (Adult)		
Grasshopper species		
Green June Beetle (Adult)		
Green Peach Aphid <sup>3</sup>		
Japanese Beetle (Adult)		
Meadow Spittlebug		
Mexican Bean Beetle		
Pea Aphid		
Pea Weevil (Adult)		
Plant Bug species including Lygus species <sup>3</sup>		
Spotted Alfalfa Aphid		
Stink Bug species		
Sweet Clover Weevil (Adult)		
Thrips species <sup>4</sup>		
Western Yellowstriped Armyworm		
Whitefringed Beetle species (Adult)		
Yellowstriped Armyworm		
Beet Armyworm <sup>1,3</sup>		
Blotch Leafminer <sup>3</sup>	0.03	1.92
Spider Mites <sup>2</sup>		

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

<sup>4</sup> Does not include Western Flower Thrips.

## Application Instructions for Alfalfa and Alfalfa Grown for Seed

- Apply as required by scouting. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gal. per acre by air or 10 gal. per acre by ground. When foliage is dense and/or pest populations are high 5-10 gal. per acre by air or 20 gal. per acre by ground and higher label rates are recommended. Use higher label rates for increased residual control.
- Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. **DO NOT** apply directly to bee shelters.
- **DO NOT** apply more than 0.03 lb. a.i. (1.92 fl. oz. or 0.12 pt. of product) per acre per cutting.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.
- **DO NOT** apply within 1 day of harvest for forage or within 7 days of harvest for hay.

## CANOLA

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Armyworm species		
Cabbage Seedpod Weevil		
Cutworm species		
Diamondback Moth	0.015 - 0.03	0.96 - 1.92
Flea Beetle		
Grasshoppers		
Looper species		
Lygus Bug		
Cabbage Aphid	0.03	1.92

## Application Instructions for Canola

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply a minimum of 2 gal. of water per acre.
- **DO NOT** apply within 7 days of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.09 lb. a.i. (5.76 fl. oz. or 0.36 pt. of product) per acre per year.

## CEREAL GRAINS:

### CORN (at Plant): Field Corn, Popcorn, Seed Corn, Sweet Corn

Target Pest	Rate	
	lb. a.i. / A	fl. oz. / A
Corn Rootworm Larvae: Mexican Northern Southern Western		
Cutworm species	0.005 lbs. a.i. per 1000 ft. of row <sup>2</sup>	0.33 fl. oz. per 1000 ft. of row <sup>2</sup>
Lesser Cornstalk Borer		
Red Imported Fire Ant <sup>1</sup>		
Seedcorn Beetle		
Seedcorn Maggot		
White Grub species		
Wireworm species		

### Application Instructions for CORN (at Plant): Field Corn, Popcorn, Seed Corn, Sweet Corn

- Banded Applications - Apply at planting as a 5-7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel.
- In-Furrow Applications - Apply into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel.
- Apply a minimum of 3 gal. finished spray per acre.
- **DO NOT** harvest or graze livestock or cut treated crops for feed within 21 days of at plant application.
- **DO NOT** apply more than 0.09 lb. a.i. (5.76 fl. oz. of product) per acre per crop at plant.
- **DO NOT** apply more than 0.032 lb. a.i. (2.0 fl. oz. of product) per acre in a single application.
- For field corn, popcorn, and seed corn **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per crop from at plant and foliar applications. For sweet corn **DO NOT** apply more than 0.48 lb. a.i. (30.72 fl. oz. or 1.92 pt. of product) per acre per crop from at plant and foliar applications.
- **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours if involved in one of the following activities –
  - Hand detasseling or mechanically assisted detasseling of field corn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of popcorn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of sweet corn grown for seed.
  - Hand detasseling of sweet corn grown for grain.

<sup>1</sup>Suppression only.

<sup>2</sup> lbs. a.i. and fl. oz./A of Omni Brand Lambda-Cy 250 CS Applied at 0.33 fl. oz./1000 ft. of Row for Various Row Spacings						
Row Spacing	40"	38"	36"	34"	32"	30"
linear ft./A	13.068	13.756	14.520	15.374	16.335	17.424
lbs. a.i./A	0.067	0.07	0.075	0.079	0.084	0.09
fl. oz./A	4.3	4.55	4.8	5.05	5.4	5.75

## CEREAL GRAINS:

### CORN (Foliar): Field Corn, Popcorn, Seed Corn

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Com Earworm <sup>1</sup> Cutworm species Green Cloverworm Meadow Spittlebug Western Bean Cutworm <sup>1</sup>	0.015 - 0.025	0.96 - 1.60
Armyworm <sup>2</sup> Bean Leaf Beetle Bird Cherry-Oat Aphid <sup>3</sup> Cereal Leaf Beetle	0.02 - 0.03	1.28 - 1.92

Com Leaf Aphid <sup>3</sup> Com Rootworm Beetle (Adult): Mexican Northern Southern Western English Grain Aphid <sup>3</sup> European Corn Borer <sup>1</sup> Fall Armyworm <sup>2</sup> Flea Beetle species Grasshopper species Hop Vine Borer <sup>1</sup> Japanese Beetle (Adult) Lesser Cornstalk Borer Sap Beetle (Adult) Seed corn Beetle Southwestern Corn Borer <sup>1</sup> Stalk Borer <sup>1</sup> Stink Bug species Tobacco Budworm <sup>1,4</sup> Webworm species Yellowstriped Armyworm <sup>2</sup>		
Beet Armyworm <sup>4</sup> Chinch Bug Greenbug <sup>3,4</sup> Mexican Rice Borer <sup>1</sup> Rice Stalk Borer <sup>1</sup> Southern Corn Leaf Beetle <sup>3</sup> Sugarcane Borer <sup>1</sup>	0.03	1.92

<sup>1</sup> For control before the larva bores into the plant stalk or ear.

<sup>2</sup> Use higher label rates for large larvae.

<sup>3</sup> Suppression only.

<sup>4</sup> See **Resistance** statement under **Use Directions**.

#### Application Instructions for CORN (Foliar): Field Corn, Popcorn, Seed Corn

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds or other locally recommended methods.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gal. of water per acre.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5-day intervals if needed. **Omni Brand Lambda-Cy 250 CS** may only suppress heavy infestations and/or subsequent migrations.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a maximum of 0.03 lb. a.i. (1.92 fl. oz. of product) per acre per application.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. **DO NOT** feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- **DO NOT** apply more than 0.032 lb. a.i. (2.0 fl. oz. of product) per acre in a single application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per crop from a plant and foliar application.
- **DO NOT** apply more than 0.06 lb. a.i. (3.84 fl. oz. or 0.24 pt. of product) per acre after silk initiation. **DO NOT** apply more than 0.03 lb. a.i. (1.92 fl. oz. or 0.12 pt. of product) per acre after corn has reached the milk stage (yellow kernels with milky fluid).
- **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours if involved in one of the following activities –
  - Hand detasseling or mechanically assisted detasseling of field corn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of popcorn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of sweet corn grown for seed.
  - Hand detasseling of sweet corn grown for grain.

**CEREAL GRAINS:**  
**CORN (Foliar): Sweet Corn**

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Aphid Species <sup>2,3</sup>		
Armyworm <sup>1</sup>		
Aster Leafhopper		
Beet Armyworm <sup>1,3</sup>		
Chinch Bug		
Common Cornstalk Borer Corn		
Earworm		
Corn Rootworm Beetle (Adult): Mexican		
Northern Southern Western		
Cutworm species		
European Com Borer		
Fall Armyworm <sup>1</sup>	0.02 - 0.03	1.28 - 1.92
Flea Beetle species		
Grasshopper species		
Japanese Beetle (Adult)		
Sap Beetle (Adult)		
Southern Armyworm <sup>1</sup>		
Southwestern Com Borer		
Spider Mite species <sup>2</sup>		
Stink Bug species		
Tarnished Plant Bug		
Webworm species		
Western Bean Cutworm		
Yellowstriped Armyworm <sup>1</sup>		
Corn Silkfly (Adult) <sup>2</sup>	0.03	1.92

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

**Application Instructions for CORN (Foliar): Sweet Corn**

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds or other locally recommended methods and target application for control before insects enter the stalk or ear.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gal. of water per acre.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 0.025 lb. a.i. (1.60 fl. oz. of product) per acre.
- **DO NOT** apply within 1 day of harvest.
- **DO NOT** allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. **DO NOT** feed treated corn fodder or silage to meat or dairy animal within 21 days after last treatment.
- **DO NOT** apply more than 0.032 lb. a.i. (2.0 fl. oz. of product) per acre in a single application.
- **DO NOT** apply more than 0.48 lb. a.i. (30.72 fl. oz. or 1.92 pt. of product) per acre per crop from all plant and foliar applications.
- **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours if involved in one of the following activities –
  - Hand detasseling or mechanically assisted detasseling of field corn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of popcorn grown for seed.
  - Hand detasseling or mechanically assisted detasseling of sweet corn grown for seed.
  - Hand detasseling of sweet corn grown for grain.

## CEREAL GRAINS: RICE, WILD RICE

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Bird Cherry-Oat Aphid		
Cinch Bug		
Fall Armyworm		
Grasshopper species		
Greenbug		
Leafhopper species	0.025 - 0.04	1.60 - 2.56
Rice Stink Bug		
Riceworm		
Rice Water Weevil (Adult)		
Sharpshooter species		
True Armyworm		
Yellow Sugarcane Aphid		
Yellowstriped Armyworm		
European Corn Borer <sup>1</sup>		
Mexican Rice Borer <sup>1</sup>		
Rice Seed Midge <sup>1</sup>	0.03 - 0.04	1.92 - 2.56
Rice Stalk Borer <sup>1</sup>		
Sugarcane Borer <sup>1</sup>		

<sup>1</sup> For control before the larvae bores into the plant stalk.

### Application Instructions for RICE, WILD RICE

- Apply as required by scouting. Base timing and frequency of application upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting.
- **Omni Brand Lambda-Cy 250 CS** can be used when propanil products are being used for weed control.
- Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water (or a total carrier volume) per acre but ensure sufficient volume is used to provide adequate coverage. In addition, adding an emulsifiable crop oil (e.g., 1 pt. per acre) when lower aerial application volumes are used is recommended to help improve coverage, reduce evaporation, and improve efficacy.
- For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. **DO NOT** exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- California: In addition to above directions for control of rice water weevil in water seeded rice, **Omni Brand Lambda-Cy 250 CS** may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field.
- Greenbug is known to have many biotypes. **Omni Brand Lambda-Cy 250 CS** may only provide suppression. If satisfactory control is not achieved with the first application of **Omni Brand Lambda-Cy 250 CS**, a resistant biotype may be present. Use alternate chemistry for control.
- For control of stem borers, scout fields, when rice growth is near panicle differentiation, for early symptoms of damaging populations exhibited as discoloration (orange-tan) around the junction of the leaf sheath and leaf blade which is caused by feeding of young larvae within the sheath. Applications must be made before larvae bore into rice stems. Make the first application at panicle differentiation to 2 inch panicle for partial control. Make the second application at boot to heading for maximum control. All rice varieties are susceptible to stem borer damage, but Cocodrie and Priscilla are particularly susceptible.
- Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 lb. ai. per acre, and treating 1200 acres (or more) per day must wear dust-mist respirator.

- **DO NOT** release flood water within 7 days of an application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.
- **DO NOT** apply more than 0.04 lb. a.i. (2.56 fl. oz. or 0.16 pt. of product) per acre within 21 to 27 days of harvest.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** use treated rice fields for the aquaculture of edible fish and crustacea.
- **DO NOT** apply as an ultra-low volume (ULV) spray.

## CEREAL GRAINS: SORGHUM (Grain)

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Cutworm species	0.015 - 0.02	0.96 - 1.28
Sorghum Midge		
Armyworm		
Beet Armyworm <sup>3</sup>		
Corn Earworm		
European Corn Borer <sup>2</sup>		
Fall Armyworm <sup>1</sup>		
Flea Beetle species	0.02 - 0.03	1.28 - 1.92
Grasshopper species		
Lesser Cornstalk Borer <sup>2</sup>		
Southwestern Corn Borer <sup>2</sup>		
Stink Bug species		
Webworm species		
Yellowstriped Armyworm <sup>1</sup>		
Chinch Bug		
Mexican Rice Borer <sup>2</sup>	0.03	1.92
Rice Stalk Borer <sup>2</sup>		
Sugarcane Borer <sup>2</sup>		

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> For control before the larva bores into the plant stalk.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

### Application Instructions for SORGHUM (Grain)

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gal. of water per acre.
- For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3 - 5-day intervals if needed. **Omni Brand Lambda-Cy 250 CS** may only suppress heavy infestations and/or subsequent migrations. .
- **DO NOT** apply more than 0.08 lb. a.i. (5.12 fl. oz. or 0.32 pt. of product) per acre per year.
- **DO NOT** apply more than 0.06 lb. a.i. (3.84 fl. oz. or 0.24 pt. of product) per acre per year after crop emergence.
- **DO NOT** apply more than 0.02 lb. a.i. (1.28 fl. oz. or 0.08 pt. of product) per acre per year once crop is in soft dough stage.
- **DO NOT** apply within 30 days of harvest.

## CEREAL GRAINS: BUCKWHEAT, BARLEY, OATS, RYE, TRITICALE, WHEAT, WHEAT HAY

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Army Cutworm		
Cutworm species	0.015 - 0.025	0.96 - 1.60
Armyworm		
Bird Cherry-Oat Aphid <sup>1</sup>	0.02 - 0.03	1.28 - 1.92
Cereal Leaf Beetle		

English Grain Aphid <sup>1</sup> Fall Armyworm Flea Beetle species Grasshopper species Hessian Fly Orange Blossom Wheat Midge Russian Wheat Aphid <sup>1</sup> Stink Bug species Yellowstriped Armyworm		
Grass Sawfly	0.025 – 0.03	1.60 - 1.92
Chinch Bug Corn Leaf Aphid <sup>2</sup> Greenbug <sup>1,3</sup> Mite species <sup>2</sup>	0.03	1.92

<sup>1</sup> Best control is obtained before insects begin to roll leaves. Once crop has started to boot, **Omni Brand Lambda-Cy 250 CS** may provide suppression only. Higher label rates and increased coverage will be necessary.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

<sup>4</sup> Make applications when adults emerge.

#### **Application Instructions for BUCKWHEAT, BARLEY, OATS, RYE, TRITICALE, WHEAT, WHEAT HAY**

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- For chinch bug control, repeat applications at 3 - 5-day intervals if needed. **Omni Brand Lambda-Cy 250 CS** may only suppress heavy infestations and/or migrations.
- Greenbug is known to have many biotypes. **Omni Brand Lambda-Cy 250 CS** may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.
- **DO NOT** apply within 30 days of harvest.
- **DO NOT** allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment. **DO NOT** feed treated straw to meat or dairy animals within 30 days after the last treatment.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.06 lb. a.i. (3.84 fl. oz. or 0.24 pt. of product) per acre per year.

#### **COLE CROPS (HEAD AND STEM BRASSICA):**

**Broccoli; Brussels Sprouts; Cabbage; Cauliflower; Cavolo Broccolo; Chinese Broccoli (gai lon); Chinese Cabbage (napa); Chinese Mustard (gai choy); Kohlrabi**

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Alfalfa Looper Cabbage Looper Cabbage Webworm Cutworm species Imported Cabbageworm Southern Cabbageworm	0.015 - 0.025	0.96 - 1.60
Aphid species <sup>2,3</sup> Armyworm Beet Armyworm <sup>1,3</sup> Corn Earworm Diamondback Moth <sup>3</sup> Fall Armyworm <sup>1</sup> Flea Beetle species Grasshopper species Japanese Beetle (Adult) Leafhopper species Meadow Spittlebug	0.02 - 0.03	1.28 - 1.92

Plant Bug species including Lygus species <sup>3</sup> Spider Mite species <sup>2</sup> Stink Bug species Thrips species <sup>2</sup> Vegetable Weevil (Adult) Whitefly species <sup>2,3</sup> Yellowstriped Armyworm		
---	--	--

<sup>1</sup> For control of first and second instar only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

#### Application Instructions for COLE CROPS (HEAD AND STEM *BRASSICA*)

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply within 1 day of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.24 lb. a.i. (15.36 fl. oz. or 0.96 pt. of product) per acre per year.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on *Brassica* (head and stem) Cole Crops.

#### COTTON

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Cutworm species		
Soybean Thrips	0.015 - 0.02	0.96 - 1.28
Tobacco Thrips		
Cabbage Looper		
Cotton Fleahopper		
Cotton Leafperforator		
Cotton Leafworm	0.02 - 0.03	1.28 - 1.92
Lygus Bug species <sup>3</sup>		
Pink Bollworm		
Saltmarsh Caterpillar		
Bandedwing Whitefly <sup>2,3</sup>		
Beet Armyworm <sup>1,3</sup>		
Boll Weevil		
Brown Stink Bug		
Cotton Aphid <sup>2,3</sup>		
Cotton Bollworm		
European Com Borer	0.025 - 0.04	1.60 - 2.56
Fall Armyworm		
Green Stink Bug		
Southern Green Stink Bug		
Sweetpotato Whitefly <sup>2,3</sup>		
Tobacco Budworm <sup>3</sup>		
Twospotted Spider Mite <sup>2</sup>		

<sup>1</sup> For control of first and second instar only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

#### Application Instructions for COTTON

- Apply as required by scouting, usually at intervals of 5 - 7 days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage.

- Applications may also be made with equipment adapted and calibrated for ULV sprays. **Omni Brand Lambda-Cy 250 CS** may be mixed with once-refined vegetable oil and applied in a minimum of at least one qt. of finished spray/A.
- Under light bollworm/budworm infestation levels, 0.02 lb. a.i. (1.28 fl. oz. of product) per acre may be applied in conjunction with intense field monitoring.
- For boll weevil control spray on a 3 - 5 day schedule.
- When applied according to label directions for control of cotton bollworm and tobacco budworm, **Omni Brand Lambda-Cy 250 CS** also provides ovicidal control of unhatched *Heliothine* species eggs.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** graze livestock in treated areas.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.2 lb. a.i (12.8 fl. oz. or 0.8 pt. of product) per acre per year.
- **DO NOT** make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.

## CUCURBIT VEGETABLES:

Chayote (fruit); Chinese Waxgourd (Chinese preserving melon); Citron Melon; Cucumber Gherkin; Gourd (edible), *Lagenaria* species - includes: hyotan, cucuzza, *Luffa acutangula*, *L. cylindrical* - includes: hechima, Chinese okra; *Momordica* species - includes: balsam apple, balsam pear, bitter melon, Chinese cucumber; Muskmelon (hybrids and/or cultivars of *Cucumis me/o*)- includes: true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon; Pumpkin; Squash, summer (*Cucurbita pepo* var. *melopepo*)- includes: crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini; Squash, winter (*Cucurbita maxima*; *C. moschata*)- includes butternut squash, calabaza, hubbard squash (*C. mixta*; *C. pepo*) - includes: acorn squash, spaghetti squash; Watermelon - includes: hybrids and/or varieties of *Citrullus lanatus*

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Armyworm species <sup>1</sup>		
Blister Beetle species		
Cabbage Looper		
Com Earworm		
Cricket species		
Cucumber Beetle species (adults)		
Cutworm species		
Flea Beetle species		
Grasshopper species		
June Beetle species		
Leaffooted Bug		
Leafhopper species		
Lugus Bug species <sup>1</sup>	0.02 - 0.03	1.28 - 1.92
Melonworm		
Pickleworm		
Plant Bug species		
Rindworm species complex		
Saltmarsh Caterpillar		
Squash Beetle		
Squash Bug species		
Squash Vine Borer species		
Stink Bug species		
Thrips species <sup>1,2</sup>		
Tobacco Budworm <sup>1</sup>		
Webworm species		
Aphid species <sup>1</sup>		
Leafminer species <sup>1,3</sup>	0.03	1.92
Spider Mite species <sup>3</sup>		
Whitefly species <sup>1,3</sup>		

<sup>1</sup> See **Resistance** statement under **Use Directions**.

<sup>2</sup> Does not include Western Flower Thrips

<sup>3</sup> Suppression only.

## Application Instructions for CUCURBIT VEGETABLES

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all plant parts. When applying by air, apply in a minimum of 2 gal. total solution per acre. When applying by ground, a minimum of 10 gal. total solution per acre is recommended.
- Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher label rates for longer residual.
- Insects that bore or tunnel into leaves, vines, stems or fruit must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of **Omni Brand Lambda-Cy 250 CS**.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on Cucurbit Vegetables.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.18 lb. a.i. (11.5 fl. oz. or 0.72 pt. of product) per acre per year.
- **DO NOT** apply within 1 day of harvest.

## FRUITING VEGETABLES:

### Eggplant; Ground cherry; Pepino; Peppers (bell and nonbell); Tomatillo; Tomato

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Cabbage Looper		
Cutworm species	0.015 - 0.025	0.96 - 1.60
Hornworm species		
Aphid species <sup>2,3</sup>		
Beet Armyworm <sup>1,3</sup>		
Blister Beetle species		
Colorado Potato Beetle <sup>3</sup>		
Cucumber Beetle species (Adult)		
European Corn Borer <sup>4</sup>		
Fall Armyworm <sup>1</sup>		
Flea Beetle species		
Grasshopper species		
Japanese Beetle (Adult)		
Leafhopper species		
Leafminer species <sup>2</sup>		
Meadow Spittlebug		
Pepper Weevil (Adult)	0.02 - 0.03	1.28 - 1.92
Plant Bug species		
Southern Armyworm <sup>1</sup>		
Spider Mite species <sup>2</sup>		
Stalk Borer <sup>4</sup>		
Stink Bug species		
Thrips <sup>5</sup>		
Tobacco Budworm <sup>3</sup>		
Tomato Fruitworm		
Tomato Pinworm		
Tomato Psyllid <sup>2,3</sup>		
Vegetable Weevil (Adult)		
Whitefly species <sup>2,3</sup>		
Yellowstriped Armyworm <sup>1</sup>		

<sup>1</sup> For control of first and second instar only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

<sup>4</sup> For control before the larva bores into the plant stalk or fruit.

<sup>5</sup> Does not include Western Flower Thrips.

### Application Instructions for FRUITING VEGETABLES

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on Fruiting Vegetables.
- **DO NOT** apply within 5 days of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.36 lb. a.i. (23.04 fl. oz. or 1.44 pt. of product) per acre per year.

### GRASS FORAGE, FODDER AND HAY:

#### Pasture and Rangeland; Grass; Grass Grown for Hay or Silage and Grass Grown for Seed

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Army Cutworm		
Cutworm species		
Essex Skipper	0.015 - 0.025	0.96 - 1.60
Range Caterpillar		
Striped Grass Looper		
Beet Armyworm		
Billbug species <sup>3</sup>		
Bird Cherry-Oat Aphid <sup>1</sup>		
Black Grass Bug		
Black Turfgrass Beetle (adult)		
Blue Stem Midge		
Cereal Leaf Beetle		
Chinch Bug		
Crane Fly species		
Cricket species		
English Grain Aphid <sup>1</sup>		
Fall Armyworm		
Flea Beetle species		
Grass Mealybug		
Grass Sawfly (adult)		
Grasshopper species		
Green June Beetle (adult)	0.02 - 0.03	1.28 - 1.92
Greenbug <sup>1,2</sup>		
Japanese Beetle (adult)		
Katydid species		
Leafhopper species		
Mite species <sup>3</sup>		
Russian Wheat Aphid <sup>1</sup>		
Southern Armyworm		
Spittlebug species		
Stink Bug species		
Sugarcane Aphid		
Thrips species		
Tick species		
True Armyworm		
Webworm species		
Yellowstriped Armyworm		

<sup>1</sup> Best control is obtained before insects begin to roll leaves.

<sup>2</sup> See **Resistance** statement under **Use Directions**.

<sup>3</sup> Suppression only.

## Application Instructions for GRASS FORAGE, FODDER AND HAY

- Apply as required by scouting. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. total solution per acre. When applying by ground, a minimum of 7 gal. total solution per acre is recommended.
- Use higher application volumes and rates when foliage is dense, pest populations are high, larvae are large and/or weather conditions are adverse. Use higher label rates for longer residual.
- For chinch bug control, **Omni Brand Lambda-Cy 250 CS** may only suppress heavy infestations and/or migrations. In this situation, a second application using an alternative chemistry may be needed.
- Greenbug is known to have many biotypes. **Omni Brand Lambda-Cy 250 CS** may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.
- Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application. **DO NOT** cut grass to be dried and harvested for hay until 7 days after the last application.

### Grass grown for seed:

- Straw and mature seed (seed screenings) may be used as feed 7 days after the last application. Regrowth of grass grown for seed may be used for grazing, cut for forage or cut to be dried and harvested for hay.
- **DO NOT** apply more than 0.03 lb. a.i. (1.92 fl. oz. or 0.12 pt. of product) per acre per cutting for pastures, rangeland and grasses grown for seed. A minimum re-treatment interval (RTI) of 30 days is required for pastures and rangeland receiving 0.03 lb. a.i. per acre which have not been cut between applications.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.09 lb. a.i. (5.76 fl. oz. or 0.36 pt. of product) per acre per year.

## LEGUME VEGETABLES (BEANS AND PEAS):

**Edible Podded (Only):** *Canavalia ensiformis* – jackbean; *Canavalia gladiata* - sword bean; *Glycine max* – soybean (immature seed);

**Edible Podded Succulent Shelled or Dried Shelled** :- *Cajanus cajan* - Pigeon Blister; *Phaseolus* species - includes: field, kidney, lima, navy, pinto, runner, snap, tepary and wax beans; *Pisum* species including: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas; *Vigna* species - includes: adzuki, asparagus, moth, mung, rice, urd and yard long beans, black- eye pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea;

**Succulent Shelled or Dried Shelled:** *Vicia faba*. - broadbean (favabean);

**Dried Shelled (Only):** *Cicer arietinum* - chickpea (garbanzo bean), *Cyamopsis tetragonoloba* – guar, *Lablab purpureus* - Lablab bean (hyacinth bean), *Lupinus* species - includes: grain, sweet, white and sweet white lupines, *Lens esculata* - Lentils

Target Pests	Rate	
	lb. a.i./A	fl. oz /A
Cutworm species		
Green Cloverworm		
Imported Cabbageworm		
Mexican Bean Beetle	0.015 - 0:025	0.96 - 1.60
Saltmarsh Caterpillar		
Velvetleaf Caterpillar		
Alfalfa Caterpillar		
Aphid species <sup>4</sup>		
Armyworm <sup>2</sup>		
Bean Leaf Beetle		
Bean Leaf-skeletonizer		
Blister Beetle species		
Corn Earworm	0.02 - 0.03	1.28 - 1.92
Corn Rootworm Beetle species (Adult)		
Cucumber Beetle species (Adult)		
Curculio and Weevil species <sup>1</sup> (foliage and pod feeding adults and larvae)		
European Corn Borer		

Fall Armyworm <sup>2</sup> Flea Beetle species (Adult) Flea Hopper species Grasshopper species Japanese Beetle (Adult) Leafhopper species Leaftier species Looper Species Meadow Spittlebug Painted Lady Butterfly (larvae) Plant Bug species Including Lygus species <sup>4</sup> Stalk Borer <sup>1</sup> Stink Bug species Three-cornered Alfalfa Hopper Thrips species <sup>4,5</sup> Tobacco Budworm Webworm species Western Bean Cutworm <sup>2</sup> Western Yellowstriped Armyworm Yellowstriped Armyworm <sup>2</sup>		
Beet Armyworm <sup>3,4</sup> Leafminer species <sup>3,4</sup> Lesser Cornstalk Borer <sup>3</sup> Soybean Looper <sup>3,4</sup> Spider Mite species <sup>3</sup> Whitefly species <sup>3,4</sup>	0.03	1.92

<sup>1</sup> For control before the larva bores into the plant stalk or pods.

<sup>2</sup> Use higher label rates for large larvae.

<sup>3</sup> For suppression only.

<sup>4</sup> See **Resistance** statement under **Use Directions**.

<sup>5</sup> Does not include Western Flower Thrips.

### Application Instructions for LEGUME VEGETABLES (BEANS AND PEAS)

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre.
- For edible podded and succulent shelled legume, vegetables, **DO NOT** apply within 7 days of harvest.
- For dried shelled legume vegetables, **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on Legume Vegetables (Beans and Peas).
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.
- For succulent and dried shelled peas and beans, **DO NOT** graze livestock in treated areas or harvest vines for forage or hay.

### LEGUME VEGETABLES (SOYBEANS)

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Bean Leaf Beetle Cabbage Looper Corn Earworm Corn Rootworm Beetle (Adult): Mexican Northern Southern Western Cutworm Species Green Cloverworm Mexican Bean Beetle	0.015 - 0.025	0.96 - 1.60

Painted Lady (Thistle) Caterpillar Potato Leafhopper Saltmarsh Caterpillar Soybean Aphid <sup>4</sup> Three-cornered Alfalfa Hopper Thrips species <sup>5</sup> Velvetbean Caterpillar Woolybear Caterpillar		
Armyworm <sup>1</sup> Blister Beetle species European Corn Borer Fall Armyworm <sup>1</sup> Grasshopper species Japanese Beetle (Adult) Plant Bug species Silverspotted Skipper Stink Bug species Tobacco Budworm <sup>3</sup> Webworm species Yellowstriped Armyworm <sup>1</sup>	0.025 - 0.03	1.60 - 1.92
Beet Armyworm <sup>2,3</sup> Lesser Cornstalk Borer <sup>2</sup> Soybean Looper <sup>2,3</sup> Spider Mite species <sup>2</sup>	0.03	1.92

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

<sup>4</sup> Use lower label rates for early season applications and/or lighter populations.

<sup>5</sup> Does not include Western Flower Thrips.

#### Application Instructions for LEGUME VEGETABLES (SOYBEANS)

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- DO NOT** graze or harvest treated soybean forage, straw or hay for livestock feed.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 0.02 lb. a.i. (1.28 fl. oz. of product) per acre.
- DO NOT** apply as foliar broadcast application using a mechanically pressurized handgun.
- DO NOT** apply within 30 days of harvest.
- DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- DO NOT** apply more than 0.06 lb. a.i. (3.84 fl. oz. or 0.24 pt. of product) per acre per year.

#### LETTUCE (HEAD AND LEAF)

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Alfalfa Looper Cabbage Looper Cutworm species Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar	0.015 - 0.025	0.96 - 1.60
Aphid species <sup>2,3</sup> Armyworm Beet Armyworm <sup>1,3</sup> Corn Earworm Diamondback Moth <sup>3</sup> European Corn Borer Fall Armyworm <sup>1</sup>	0.02 - 0.03	1.28 - 1.92

Flea Beetle species Grasshopper species Japanese Beetle (Adult) Leafhopper species Meadow Spittlebug Plant Bug species including <i>Lygus</i> species <sup>3</sup> Southern Armyworm Spider Mite species <sup>2</sup> Stink Bug species Tobacco Budworm <sup>3</sup> Vegetable Weevil (Adult) Whitefly species <sup>2,3</sup>		
---	--	--

<sup>1</sup> For control of first and second instar only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

#### Application Instructions for LETTUCE (HEAD AND LEAF)

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on Legume Vegetables (Soybeans).
- **DO NOT** apply within 1 day of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.3 lb. a.i. (19.2 fl. oz. or 1.2 pt. of product) per acre per year.

#### ONION (BULB) AND GARLIC

Target Pests	Rate	
	Ib. a.i./A	fl. oz. /A
Cutworm species		
Leafminer species (Adult)		
Onion Maggot (Adult)	0.015 - 0.025	0.96 - 1.60
Seedcorn Maggot (Adult)		
Aphid species <sup>2</sup>		
Armyworm species <sup>1</sup>		
Flower Thrips <sup>2,3</sup>		
Onion Thrips <sup>3</sup>		
Plant Bug species	0.02 - 0.03	1.28 - 1.92
Stink Bug species		
Tobacco Thrips <sup>3</sup>		
Western Flower Thrips <sup>3</sup>		

<sup>1</sup> For control of the first and second instar only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

#### Application Instructions for ONION (BULB) AND GARLIC

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Use the higher label rates as thrips population increases and avoid rescue situations.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage.
- **DO NOT** apply as foliar broadcast application using a mechanically pressurized handgun on Onion (bulb) and Garlic.
- **DO NOT** apply within 14 days of harvest.

- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.24 lb. a.i. (15.36 fl. oz. or 0.96 pt. of product) per acre per year.

## PEANUTS

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Cutworm species		
Green Cloverworm		
Potato Leafhopper		
Red-necked Peanut Worm	0.015 - 0.025	0.96 - 1.60
Three-cornered Alfalfa Hopper		
Velvetbean Caterpillar		
Bean Leaf Beetle		
Com Earworm		
Fall Armyworm <sup>1</sup>		
Grasshopper species		
Southern Corn Rootworm (Adult)	0.02 - 0.03	1.28 - 1.92
Stink Bug Species		
Tobacco Thrips		
Vegetable Weevil		
Whitefringed Beetle (Adult)		
Aphid species <sup>2</sup>		
Beet Armyworm <sup>2,3</sup>		
Lesser Cornstalk Borer <sup>2</sup>	0.03	1.92
Soybean Looper <sup>2,3</sup>		
Spider Mite species <sup>2</sup>		

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

## Application Instructions for PEANUTS

- Apply as required by scouting, usually at intervals of 7 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.

## POMEFRUITS:

### Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Aphid		
Apple Maggot (Adult)		
Cherry Fruit Fly species (Adult)		
Codling Moth		
Green Fruitworm		
Japanese Beetle		
Leafhopper species		
Leafroller species	0.02 - 0.04	1.28 - 2.56
Lesser Appleworm		
Omnivorous Leafroller		
Orange Tortrix		
Oriental Fruit Moth		
Pear Psylla <sup>1</sup>		
Pear Sawfly		
Periodical Cicada		

Plant Bug species Plum Curculio Rosy Apple Aphid San Jose Scale (fruit infestations only) Spirea Aphid <sup>1</sup> Stink Bug species Tent Caterpillar species Tentiform Leaf Miner species Tree Borer species Tufted Apple Budworm Webworm species		
---	--	--

<sup>1</sup> Suppression only

### Application Instructions for POME FRUITS

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds and 1PM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gal. of water per acre, but use higher volumes as appropriate for thorough coverage.
- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on orchards.
- **DO NOT** apply as a foliar broadcast application or as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards and vineyards.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.2 lb. a.i. (12.8 fl. oz. or 0.80 pt. of product) per acre per year.
- **DO NOT** apply more than 0.16 lb. a.i. (10.24 fl. oz. or 0.64 pt. of product) per acre per year post bloom.

### STONE FRUITS:

**Apricot, Chickasaw Plum, Damson Plum, Japanese Plum, Nectarine, Peach, Plum, Plumcot, Prune, Sweet and Tart Cherry,**

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
American Plum Borer Apple Maggot (Adult) Black Cherry Aphid Cherry Fruit Fly species (Adult) Codling Moth Green Fruitworm Japanese Beetle June Beetle Leafhopper species Leafroller species Oriental Fruit Moth Peach Twig Borer Peachtree Borer species Pear Sawfly Periodical Cicada Plant Bug species Plum Curculio Rose Chafer Stink Bug species Tent Caterpillar species Thrips species	0.02 - 0.04	1.28 - 2.56

### Application Instructions for STONE FRUITS

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic threshold and 1PM recommendations.

- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 5 gals. of water per acre, but use higher volumes as appropriate for thorough coverage.
- **DO NOT** apply as a foliar broadcast application or as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards and vineyards.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.2 lb. a.i. (12.8 fl. oz. or 0.80 pt. of product) per acre per year.
- **DO NOT** apply more than 0.16 lb. a.i. (10.24 fl. oz. or 0.64 pt. of product) per acre per year post bloom.

## SUGARCANE

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Mexican Rice Borer <sup>1</sup>		
Pygmy Mole Cricket		
Rice Stalk Borer <sup>1</sup>		
Sugarcane Aphid <sup>3</sup>		
Sugarcane Beetle (Adult) <sup>2</sup>	0.025 - 0.04	1.60 - 2.56
Sugarcane Borer <sup>1</sup>		
West Indian Crane fly		
Yellow Sugarcane Aphid <sup>3</sup>		

<sup>1</sup> For control before the larva bores into the plant stalk.

<sup>2</sup> Suppression only of beetles active above ground.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

## Application Instructions for SUGARCANE

- Apply as required by scouting, usually at intervals of 7 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gal. of water per acre.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.16 lb. a.i. (10.24 fl. oz. or 0.64 pt. of product) per acre per year.

## SUNFLOWER

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Cutworm species		
Sunflower Beetle	0.015 - 0.025	0.96 - 1.60
Banded Sunflower Moth		
Fall Armyworm <sup>1</sup>		
Grasshopper species		
Head-Clipper Weevil (Adult)		
Japanese Beetle (Adult)		
Leafhopper species		
Meadow Spittlebug		
Painted Lady (Thistle) Caterpillar	0.02 - 0.03	1.28 - 1.92
Seed Weevil (Adult)		
Spotted Cabbage Looper		
Stem Weevil (Adult)		
Stink Bug species		
Sunflower Maggot (Adult)		
Sunflower Moth		
Woollybear Caterpillar		
Beet Armyworm <sup>2,3</sup>	0.03	1.92
Spider Mite species <sup>2</sup>		

<sup>1</sup> Use higher label rates for large larvae.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

### Application Instructions for SUNFLOWERS

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply within 45 days of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.
- **DO NOT** apply more than 0.09 lb. a.i. (5.76 fl. oz. or 0.36 pt. of product) per acre per year after bloom initiation.
- **DO NOT** apply as an ultra-low volume (ULV) spray.

### TOBACCO

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Armyworm species <sup>1</sup>		
Blister Beetle species		
Cabbage Looper		
Corn Earworm		
Cucumber Beetle species (Adult)		
Cutworm species		
Grasshopper species		
Japanese Beetle (Adult)		
Katydid species		
Plant Bug species <sup>3</sup>		
Potato Tuberworm	0.015 - 0.03	0.96 - 1.92
Salt Marsh Caterpillar		
Stinkbug species		
Tobacco Aphid species <sup>2,3</sup>		
Tobacco Budworm <sup>3</sup>		
Tobacco Flea Beetle (Adult)		
Tobacco Hornworm		
Tobacco Thrips species <sup>2</sup>		
Tomato Hornworm		
Tree Cricket species		
Vegetable Weevil (Adult)		
Webworm species		

<sup>1</sup> For control of first and second instars only.

<sup>2</sup> Suppression only.

<sup>3</sup> See **Resistance** statement under **Use Directions**.

### Application Instructions for TOBACCO

- Apply as required by scouting, usually at intervals of 7 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage. When applying by air, apply in a minimum of 2 gal. of water per acre.
- **DO NOT** apply as foliar broadcast application using a mechanically pressurized handgun on Tobacco.
- **DO NOT** apply within 40 days of harvest.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.09 lb. a.i. (5.76 fl. oz. or 0.36 pt. of product) per acre per year.

**TREE NUTS:**

**Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut (Bush Nut), Pistachio, Walnut, Black Walnut, English (Persian)**

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Chinch Bug		
Codling Moth		
Filbertworm		
Leaffooted Bug		
Leafroller species	0.02 - 0.04	1.28 - 2.56
Navel Orangeworm		
Peach Twig Borer		
Plant Bug species		
Stink Bug species		
Walnut Aphid		
Walnut Husk Fly species (Adult)		

**Pecans**

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Hickory Shuckworm		
Pecan Aphid species		
Pecan Casebearer species		
Pecan Phylloxera species	0.02 - 0.04	1.28 - 2.56
Pecan Spittlebug		
Pecan Weevil		
Stink Bug species		

**Application Instructions for TREE NUTS**

- Apply as required by scouting, usually at intervals of 5 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gal. of water per acre, but use higher label rates as appropriate for thorough coverage.
- **DO NOT** apply as a foliar broadcast application or as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards and vineyards.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.16 lb. a.i. (10.24 fl. oz. or 0.64 pt. of product) per acre per year.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year post bloom.

**TUBEROUS AND CORM VEGETABLES (Potato, Sweet Potato, Yams and Related):**

**Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem only), Canna (edible), Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen, Ginger, Leren, Potato, Sweet Potato, Tanier, Turmeric, Yam (bean and true)**

Target Pests	Rate	
	Ib. a.i./A	fl. oz./A
Cutworm species		
Leafhopper species		
Saltmarsh Caterpillar	0.015 - 0.025	0.96 - 1.60
Sweet Potato Hornworm		
Woollybear Caterpillar species		
Aphid species <sup>1</sup>		
Armyworm species <sup>1</sup>		
Blister Beetle species	0.02 - 0.03	1.28 - 1.92
Colorado Potato Beetle <sup>1</sup>		
Corn Earworm		

Cricket species		
Cucumber Beetle species (adults)		
European Corn Borer		
Flea Beetle species (adults)		
Grasshopper species		
Looper species <sup>1</sup>		
Lygus Bug species <sup>1</sup>		
Plant Bug species		
Potato Psyllid		
Potato Tuberworm		
Stink Bug species		
Sweet Potato Leaf Beetle (adults)		
Sweet Potato Vine Borer		
Thrips species <sup>1,2</sup>		
Tortoise Beetle species		
Webworm species		
Weevil species (adults)		
Leafminer species <sup>1,3</sup>		
Whitefly species <sup>1,3</sup>	0.03	1.92
Spider Mite species <sup>3</sup>		

<sup>1</sup> See **Resistance** statement under **Use Directions**.

<sup>2</sup> Does not include Western Flower Thrips.

<sup>3</sup> Suppression only.

#### Application Instructions for TUBEROUS AND CORM VEGETABLES

- Apply as required by scouting, usually at intervals of 7 or more days. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all above ground plant parts. When applying by air, apply in a minimum of 2 gal. total solution per acre. When applying by ground, a minimum of 10 gal. total solution per acre is recommended.
- Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher label rates for longer residual.
- Insects that bore or tunnel into leaves, vines, stems, tubers or corms must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of Omni Brand Lambda-Cy 250 CS.
- **DO NOT** apply as foliar broadcast application using a mechanically pressurized handgun.
- **DO NOT** apply more than 0.03 lb ai (1.92 fl oz or 0.12 pt of product) per acre per application.
- **DO NOT** apply more than 0.12 lb. a.i. (7.68 fl. oz. or 0.48 pt. of product) per acre per year.
- **DO NOT** apply within 7 days of harvest.

#### CONIFER AND DECIDUOUS TREES:

##### Plantations and Nurseries

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
Bagworm		
Balsam Twig Aphid		
Balsam Wooly Aphid		
Birch Leafminer		
Black Pine Weevil		
Elm Leaf Beetle		
European Elm Bark Beetle	0.02 - 0.04	1.28 - 2.56
Gypsy Moth		
Japanese Beetle		
June Beetle species		
Leaf Beetle species		
Leafroller species		
May Beetle species		
Mealybug species <sup>1</sup>		

Pales Weevil Pine Chafer Pine Colaspis Beetle Pine Conelet Bug Pine Leaf Chermid Pine Needle Scale Pine Sawfly species Pine Tip Moth species Pine Tortoise Scale Pine Weevil species Poplar Aphid species Sawfly species Spittlebug species Spruce Budworm Tent Caterpillar species Tussock Moth species Webworm species		
--	--	--

<sup>1</sup> Suppression only.

#### Application Instructions for CONIFER AND DECIDUOUS TREES

- To control exposed foliage, flower, cone, seed and bark feeding insects, apply as required by scouting. Base timing and frequency of applications upon insect populations reaching locally determined economic thresholds.
- Apply with ground equipment using sufficient water to obtain full coverage of target site. When applying by air, apply a minimum of 2 gal. of water per acre.
- **DO NOT** apply more than 0.04 lb ai (2.56 fl oz or 0.16 pt of product) per acre per application.
- **DO NOT** apply more than 0.24 lb. a.i. (15.36 fl. oz. or 0.96 pt. of product) per acre per year.

#### CONIFER AND DECIDUOUS TREES:

##### Seed Orchards

Target Pests	Rate
Coneworm species	For high volume sprayers, dilute 2.56 fl. oz. per 100 gal. of water and apply 5-10 gal. of finished spray per application.
Seed Bug species	For low volume sprayers, dilute 10 fl. oz. per 100 gal. of water and apply 100 gal. of finished spray per application.
Thrips species	For aerial applications, apply 7.5 fl. oz. per acre in a minimum of 10 gal. finish spray per application.

#### Application Instructions for CONIFER AND DECIDUOUS TREES: Seed Orchards

- **DO NOT** apply as a foliar broadcast application using a mechanically pressurized handgun on orchards.
- **DO NOT** apply as a soil, drench, or ground-directed application using a mechanically pressurized handgun on orchards.
- **DO NOT** apply more than 0.5 lb. a.i. (32 fl. oz. or 2 pt. of product) per acre per year.

## NON-AGRICULTURAL USES

### NON-CROPLAND (Crop Outlets, including turn rows, wind rows, hedge rows, field borders and buffer zones) (EXCLUDING PUBLIC LAND)

Target Pests	Rate	
	lb. a.i./A	fl. oz./A
See Specific Crops on this <b>Omni Brand Lambda-Cy 250 CS</b> label for target pest and rates.	See Specific Crops	See Specific Crops

### Application Instructions for NON-CROPLAND (EXCLUDING PUBLIC LAND)

- Spray non-cropland adjacent to agricultural areas to control migratory insects, which may threaten crops.
- Follow **Use Directions**, rates and spray directions found elsewhere in this label for the adjacent crop outlet and target pests.
- Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages.
- Repeat as necessary to maintain control.
- **DO NOT** exceed 0.2 lb. a.i. (25.6 fl. oz. or 1.6 pt. of product) per acre per year.
- **DO NOT** graze livestock in treated areas.

### STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

#### PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER HANDLING:

**Nonrefillable container.** **DO NOT** reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For packages up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

drip.

**For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For refillable containers:** Refill this container with this product only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For final disposal, offer for recycling or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

#### **CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES**

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and must be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Agri-Enterprises, LLC (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. To the extent consistent with applicable law, the Company makes no other warranties or representations of any kind; express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Agri-Enterprises, LLC's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 20XX.

®/™ All other products are trademarks of their respective companies.