



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505T)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:
 5905-672

Date of Issuance:
 3/31/26

NOTICE OF PESTICIDE:
 Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:
 Unconditional

Name of Pesticide Product:
 Omni Brand Pendi-Cap 3.8

Name and Address of Registrant (include ZIP Code):

Helena Agri-Enterprises, LLC
 225 Schilling Boulevard, Suite 300
 Collierville, TN 38017

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
 Herbicide Branch, Registration Division (7505T)

Date:

3/31/26

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 5905-672."
3. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF:

- Basic CSF dated 07/18/2024

If you have any questions, please contact Jamie Millard at (202) 566-2726 or by email at millard.jamie@epa.gov.

Enclosure

{Note to reviewer: [Text] in brackets denotes optional text. {Text} in braces denotes where in the final label text will appear and notes to reviewer.}

{BOOKLET FRONT PANEL}

PENDIMETHALIN	GROUP	3	HERBICIDE
---------------	-------	----------	-----------

Omni Brand Pendi-Cap 3.8

Herbicide

[For Use in Selected Crops]

[(See Table 1. Crop Uses)]

ACTIVE INGREDIENT:	% BY WT
pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine*	38.7%
OTHER INGREDIENTS:	61.3%
TOTAL:	100.0%

*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	

[See] [inside] [label] [booklet] [side] [panel] [for] [First Aid][,] [additional] [Precautionary Statements][,] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

EPA Reg. No.: 5905-[XX]

EPA Est. No.:


Manufactured for:

Helena Agri-Enterprises, LLC
225 Schilling Boulevard, Suite 300
Collierville, TN 38017

[AD XXXXX]{internal tracking}

[Lot][Batch][No.]:

NET CONTENTS:



{INSIDE BOOKLET}

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwater or rinsate.

NON-TARGET ORGANISM ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Endangered Species Protection

This product may have effects on federally listed threatened or endangered plant species or their critical habitat. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-844-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will be available from the above sources 6 months before their effective dates.

If endangered plant species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave an untreated buffer zone of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASABE fine to medium/coarse nozzles.
- If applied by air, leave an untreated buffer zone of 170 feet. Must use straight-stream nozzles (D-6 or larger); wind can be no more than 8 mph, and release height must be 15 feet or less.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide protection.

Do not enter or allow other people (or pets) to enter the treated area until sprays have dried.

Observe all precautions and restrictions in this label and the labels of products used in combination with **Omni Brand Pendi-Cap 3.8** herbicide. Use of **Omni Brand Pendi-Cap 3.8** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

[Helena Agri-Enterprises, LLC intends that this product may not be used for manufacturing products for application to turf and ornamentals.]

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)

Steps to take if material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

EXCEPTION: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

PRODUCT INFORMATION

Omni Brand Pendi-Cap 3.8 herbicide is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to **Table 1** for crop uses. Refer to **Table 2** for a complete list of controlled weeds. **Omni Brand Pendi-Cap 3.8** will not control established weeds.

Table 1. Crop Uses

African marigold	fig trees, nonbearing	perennial grasses grown for seed
alfalfa	forage grasses (cool-season)	pome fruit trees, bearing and
artichoke	forage grasses (warm-season)	nonbearing
asparagus	fruiting vegetables	pomegranate
berries, low-growing	garlic	potato
Brassica head and stem vegetables	grain sorghum	rice
bushberries, bearing and	grape, bearing and nonbearing	rosemary
nonbearing	vineyards	safflower
caneberries, bearing and	hops	small fruit climbing vines, bearing
nonbearing	leaf lettuce	and nonbearing
carrot	leafy Brassica greens	soybean
celery (and other leaf petiole	leek	stone fruit trees, bearing and
vegetables in Group 22B)	lentil and peas	nonbearing
citrus fruit trees, bearing and	melons	strawberry
nonbearing	mint	sugarcane
corn (field, field seed, fresh sweet,	monarda	sunflower (and other Group 20B
popcorn, popcorn seed)	nut trees, bearing and nonbearing	oilseeds)
cotton	olive trees, bearing and nonbearing	tobacco
date palm trees, nonbearing	onions and shallots (dry bulb,	triticale
edible beans	green)	vegetable soybean (edamame)
fallow	peanut	wheat

Table 2. Weeds Controlled

(see crop sections for additional weeds controlled)

Weeds controlled with Omni Brand Pendi-Cap 3.8 applied up to 4 pts/A.		
GRASS WEEDS		
annual ryegrass*	foxtail, green	oat, wild*
barnyardgrass	foxtail, yellow	panicum, fall
canarygrass*, ¹	goosegrass	panicum, Texas
cheat*, ¹	hairy chess*, ²	sandbur, field
crabgrass	itchgrass*	shattercane*
crowfootgrass	Italian ryegrass*	signalgrass*
cupgrass	Japanese brome*, ²	wild proso millet*
downy brome*	johnsongrass (seedling)	witchgrass
foxtail, giant	jointed goatgrass*, ²	woolly cupgrass*
BROADLEAF WEEDS		
amaranth, Palmer	lady's thumb	pusley, Florida
bittercress ^[1]	lambsquarters, common	redstem, filaree
black nightshade	lambsquarters, slimleaf ¹	shepherdspurse*
bugloss, small ²	London rocket*	smartweed, Pennsylvania*
carpetweed	mustard, black ¹	spurge, annual
chickweed, common*	pigweed species	velvetleaf*
common groundsel	prickly sida	waterhemp species
henbit	prostrate knotweed ^[1]	wild mustard
kochia	purslane	
Weeds controlled with Omni Brand Pendi-Cap 3.8 applied at 4 pts/A or more.		
GRASS WEEDS		
annual bluegrass	junglerice	sprangletop, red
browntop panicum	lovegrass	swollen fingergrass
grass, Guinea ¹	sprangletop, Mexican	

BROADLEAF WEEDS		
dodder [†] fiddleneck morningglory**	prickly lettuce prostrate knotweed puncturevine	redmaids rockpurslane
<p>* Suppression, but controlled when Omni Brand Pendi-Cap 3.8 use rate exceeds 4 pts/A. ** Suppression. [†] For optimum dodder control, use the highest labeled rate of Omni Brand Pendi-Cap 3.8 specified in the specific crop. ¹ Not controlled in California. ² Not suppressed or controlled in California.</p>		

Use **Table 3** and **Table 4** to determine the corresponding amount of active ingredient (pendimethalin) from **Omni Brand Pendi-Cap 3.8** product use rates.

Table 3. Use Rate Equivalency for Omni Brand Pendi-Cap 3.8 in Quarts (3.8 lbs ai/gallon)

Amount of Omni Brand Pendi-Cap 3.8 (qts/A)	Amount of Omni Brand Pendi-Cap 3.8 (fl ozs/A)	Amount of Pendimethalin (lbs ai/A)
1.1	35.2	1.05
2.1	67.2	2.00
2.6	83.2	2.47
3.2	102.4	3.04
3.5	112.0	3.33
4.2	134.4	3.99
6.3	201.6	5.99

Table 4. Use Rate Equivalency for Omni Brand Pendi-Cap 3.8 in Pints (3.8 lbs ai/gallon)

Amount of Omni Brand Pendi-Cap 3.8 (pts/A)	Amount of Omni Brand Pendi-Cap 3.8 (fl ozs/A)	Amount of Pendimethalin (lbs ai/A)
1.0	16.0	0.48
1.1	17.6	0.52
1.5	24.0	0.71
2.0	32.0	0.95
2.1	33.6	1.00
2.4	38.4	1.14
3.0	48.0	1.43
3.1	49.6	1.47
3.2	51.2	1.52
4.0	64.0	1.90
4.2	67.2	2.00
6.4	102.4	3.04
8.0	128.0	3.80
8.2	131.2	3.90
12.5	200.0	5.94
12.6	201.6	5.99

MODE OF ACTION

Pendimethalin, the active ingredient in **Omni Brand Pendi-Cap 3.8**, is a **Group 3 (WSSA)/Group K1 (HRAC)** herbicide belonging to the dinitroaniline chemistry class. **Omni Brand Pendi-Cap 3.8** is a meristematic inhibitor that interferes with meristematic plant cell division or mitosis inhibiting germinating seedling growth.

HERBICIDE RESISTANCE MANAGEMENT

Omni Brand Pendi-Cap 3.8 is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to **Omni Brand Pendi-Cap 3.8** and other Group 3 herbicides. Weed species with resistance to Group 3 may eventually dominate the weed population if Group 3 herbicides are used repeatedly in the same field or in

successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by **Omni Brand Pendi-Cap 3.8** or other Group 3 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of **Omni Brand Pendi-Cap 3.8** or other target site-of-action Group 3 herbicides that have a similar target site of action on the same weed species.
- Using tank mixes or premixes with herbicides from different target-site-of-action groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM (Integrated Pest Management) program including cultural and mechanical methods.
- Monitoring treated weed populations for loss of field efficacy, and control of escapes with effective alternative herbicides or mechanical methods.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed control program needs to consider all of the weeds present.
- Scout fields prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- Scout fields after application to verify the treatment was effective.
- Suspected herbicide-resistance weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species.
- If resistance is suspected, treat weed escapes with an herbicide with a different MOA and/or use non-chemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Report any incidence of non-performance of this product against a particular weed species to your local Helena Agri-Enterprises, LLC representative.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management directions for specific crops and resistant weed biotypes.

APPLICATION RATE

Use rates for **Omni Brand Pendi-Cap 3.8** herbicide when used alone, in tank mix, or for sequential applications are given in **Crop-specific Information**. Use rates of this product vary by soil texture and organic matter. See **Table 5** for soil texture groupings used in this label.

Do not apply more than the maximum labeled rate of **Omni Brand Pendi-Cap 3.8** for any soil type.

Table 5. Soil Texture Groups

Coarse	Medium	Fine
sands loamy sands sandy loams	sandy clay loams* sandy clays loams silt loams silts	silty clay loams* silty clays clay loams clays
* Sometimes considered transitional soils and may be classified as medium-texture or fine-texture soils.		
For peat and muck soils. Omni Brand Pendi-Cap 3.8 may be used on peat and muck soils, but weed control may be inconsistent and/or reduced. Use maximum labeled use rate allowed in the specific crop.		

APPLICATION TIMINGS

Omni Brand Pendi-Cap 3.8 will provide most effective weed control when applied by ground or aerial equipment and subsequently incorporated into soil by rainfall, sprinkler irrigation, or mechanical tillage before weed seedling germination.

Omni Brand Pendi-Cap 3.8 can also be applied through chemigation, including flooded basin irrigation systems. **Omni Brand Pendi-Cap 3.8** may be applied preplant surface, preplant incorporated, surface incorporated,

preemergence, early postemergence, postemergence incorporated (CULTI-SPRAY), or by layby treatment. See **Crop-specific Information** for specific application directions and restrictions by crop.

Preplant Surface Application. For use in minimum tillage or no-tillage production systems, apply **Omni Brand Pendi-Cap 3.8** alone or in tank mixes within 45 days of planting. When making early preplant surface application (15 to 45 days before planting), tank mix **Omni Brand Pendi-Cap 3.8** or follow with a postemergence herbicide application. Rainfall or sprinkler irrigation after application is required to move this product into the upper soil surface where weed seeds germinate.

Preplant Incorporated Application. Apply **Omni Brand Pendi-Cap 3.8** and incorporate into the upper (1 inch to 2 inches) soil surface within 60 days of planting. Use an implement capable of giving uniform incorporation; two-pass incorporation usually results in a more consistent result.

Surface Incorporated Application. Uniformly apply **Omni Brand Pendi-Cap 3.8** as broadcast or banded treatment to soil surface underneath established trees and/or in ground areas between trees rows. Incorporate into upper (1 inch to 2 inches) soil surface using rainfall, sprinkler irrigation, or shallow mechanical incorporation using an implement capable of giving uniform incorporation; two-pass mechanical incorporation usually results in a more consistent result.

Preemergence Surface Application. Broadcast treatment uniformly to the soil surface at planting and up to 2 days after planting (refer to **Crop-specific Information** section for exceptions). Rainfall, sprinkler irrigation, or shallow mechanical incorporation after application is required to move this product into the upper soil surface where weed seeds germinate. If adequate rainfall or irrigation does not occur, or soil crusting or soil compaction has occurred, and weed seedling emergence begins, a shallow cultivation or rotary hoeing or light harrow will improve performance. Make sure that crop seeds are below the tilled soil surface area.

Early Postemergence Application. **Omni Brand Pendi-Cap 3.8** must be applied before weed seedling emergence or in a tank mix with products that control the emerged weeds. Refer to **Crop-specific Information** for specific postemergence application instructions by crop.

Postemergence Incorporated Application (CULTISPRAY). Before application, crop must be cultivated in such a manner as to throw at least 1 inch of soil over the base of the crop plants. This will prevent direct contact of **Omni Brand Pendi-Cap 3.8** and the zone of brace root formation. **Omni Brand Pendi-Cap 3.8** must be applied broadcast with a ground sprayer when crop is at least 4 inches tall up to layby. Use drop nozzles if crop foliage will prevent uniform coverage of the soil surface within the rows. Thoroughly and uniformly incorporate **Omni Brand Pendi-Cap 3.8** treatments into the soil:

1. With a sweep-type or rolling cultivator set to provide thorough incorporation in the top 1 inch of soil, or
2. With adequate overhead irrigation water or rainfall. See **Crop-specific Information (Corn and Grain Sorghum)** for more details on (CULTI-SPRAY) application.

Layby Application. Apply **Omni Brand Pendi-Cap 3.8** herbicide directly to the soil between rows as a directed spray after the last normal cultivation (layby). See **Crop-specific Information** for more details on layby application.

Split Application. **Omni Brand Pendi-Cap 3.8** may be applied preplant incorporated within 60 days of planting and followed by a preemergence application at planting or up to 2 days after planting (refer to **Crop-specific Information** section for exceptions). The total amount of **Omni Brand Pendi-Cap 3.8** applied per acre per year cannot exceed the highest labeled rate for any given soil type. See **Crop-specific Information** for more details on split applications.

Fall Application. **Omni Brand Pendi-Cap 3.8** may be used in fall application programs in certain crops. See **Crop-specific Information** for details on fall application timing.

APPLICATION INSTRUCTIONS

Omni Brand Pendi-Cap 3.8 may be applied using water or sprayable fluid fertilizer (including straight 32-0-0 or 28-0-0) as the spray carrier. Do not apply **Omni Brand Pendi-Cap 3.8** after crop emergence (postemergence) in sprayable liquid fertilizers. Additionally, **Omni Brand Pendi-Cap 3.8** may be impregnated on dry bulk fertilizer. For all applications, avoid overlapping that could result in possible crop injury.

Spray Mix Compatibility Test

If liquid fertilizer and/or herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result. Always predetermine the compatibility of **Omni Brand Pendi-Cap 3.8** alone or with other herbicides based on the following compatibility jar test:

1. Add 1 pint of fertilizer to a quart jar.
2. Add 1 to 4 teaspoon(s) of the dry flowable (DF), wettable powder (WP), aqueous solution (AS), flowable (F), or liquid (L) formulation (depending on mixing ratio required) to the liquid fertilizer. To calculate teaspoons of the formulation to add:

$$\frac{\text{lbs or pts of product/acre}}{\text{gallons of fertilizer/acre}} \times 11.4 = \text{teaspoons of herbicide to add to 1 pint of fertilizer}$$

3. Close the jar and agitate until the herbicide(s) are evenly dispersed in the liquid fertilizer. If the materials do not disperse well, it may be necessary to slurry the chemicals in water before adding to the fertilizer.
4. After dispersing the materials, add appropriate number of teaspoons of **Omni Brand Pendi-Cap 3.8** to the jar and shake well. Add water soluble concentrate herbicides to the mixture last and agitate. Let the mixture stand for 30 minutes; then observe the results. Look for signs of separation: an oily layer or globules, sludge, flakes or other precipitates.
5. Evaluate compatibility.
 - a. If the herbicide(s) and liquid fertilizer mixture does not separate, use this mixture in your spray tank.
 - b. If the mixture separates but mixes readily with shaking, the mixture can be used if good agitation is maintained in the spray tank.
 - c. If separation of the mixture occurs and agitation does not correct this problem, a compatibility agent is needed.
6. If the need for a compatibility agent is demonstrated, Helena Agri-Enterprises, LLC advises the following procedure: Using a clean quart jar, repeat step 1 above and add ½ teaspoon of the compatibility agent to the liquid fertilizer. Mix well and repeat steps 2, 3, and 4. If separation or precipitation occurs with the compatibility agent. Do not use **Omni Brand Pendi-Cap 3.8** with that specific liquid fertilizer.

Aerial Application

Uniformly apply in 5 or more gallons of water per acre.

Ground Application (Broadcast)

Uniformly apply in 10 or more gallons of water per acre or 20 or more gallons of liquid fertilizer per acre. Keep the bypass line on or near the bottom of the tank to minimize foaming. Nozzle and in-line screens must be no finer than 50 mesh.

Ground Application (Band)

Uniformly apply the broadcast equivalent rate and volume per acre. Calculate:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{band rate per acre}$$

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast volume per acre} = \text{band volume per acre}$$

Ground Application (Dry Bulk Fertilizer)

Apply **Omni Brand Pendi-Cap 3.8** herbicide/dry bulk fertilizer mixtures only with ground equipment. See **Crop-specific Information** for crops suitable for dry bulk fertilizer applications. Do not impregnate **Omni Brand Pendi-Cap 3.8** onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with **Omni Brand Pendi-Cap 3.8**. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Calculate the amount of **Omni Brand Pendi-Cap 3.8** impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer to be applied per acre:

$$\frac{2000}{\text{pounds of dry fertilizer per acre}} \times \text{Omni Brand Pendi-Cap 3.8 (rate per acre)} = \text{Omni Brand Pendi-Cap 3.8 per ton of fertilizer}$$

To impregnate **Omni Brand Pendi-Cap 3.8** on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of **Omni Brand Pendi-Cap 3.8** onto the fertilizer during mixing.

Apply the **Omni Brand Pendi-Cap 3.8**/dry bulk fertilizer mixture with a calibrated dry fertilizer spreader. The **Omni Brand Pendi-Cap 3.8**/dry bulk fertilizer mixture must be spread uniformly on the soil surface.

Chemigation Application via Sprinkler Irrigation and Drip Irrigation Systems

Omni Brand Pendi-Cap 3.8 may be applied as a chemigation treatment through sprinkler irrigation and drip irrigation systems. Refer to **Crop-specific Information** sections for individual crops. Do not apply **Omni Brand Pendi-Cap 3.8** via chemigation to crops unless specified in **Crop-specific Information** section.

Apply this product **ONLY** through a sprinkler irrigation system of the following type: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move.

Apply this product **ONLY** through a drip irrigation system that has emitters above the soil surface.

Do not apply this product through any other type of sprinkler irrigation or drip irrigation system.

Uniform distribution of **Omni Brand Pendi-Cap 3.8**-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness, or illegal pesticide residues in the crop. If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.

The system must be calibrated (with water only) to ensure the amount of **Omni Brand Pendi-Cap 3.8** applied corresponds to the specified rate. Apply **Omni Brand Pendi-Cap 3.8** in ½ to ¾ inch of water during the first sprinkler set (use at least 1 inch of water in the states of **New Mexico, Oklahoma, and Texas**). Helena Agri-Enterprises, LLC advises mixing **Omni Brand Pendi-Cap 3.8** with water at a 1:1 ratio in the injection nurse tank to assist with product flowability. Maintain agitation in the injection nurse tank to keep a uniform herbicide suspension during application. When application is complete, flush the system with water.

Chemigation Instructions (for low-volume micro sprinklers)

Output of low-volume sprinkler equals 4 to 50 gallons per hour (gph) per emitter. Point of application **MUST** be above ground.

Irrigation system must run a sufficient amount of time before **Omni Brand Pendi-Cap 3.8** injection to have all emitters functioning properly. After system is operating properly, length of injection must be such that at one period of time during the injection, the first and last emitters in the system contain **Omni Brand Pendi-Cap 3.8**-treated water. Add **Omni Brand Pendi-Cap 3.8** to the supply tank already filled with the volume of water required for the injection period. Maintain agitation in **Omni Brand Pendi-Cap 3.8** injection tank. Mix **Omni Brand Pendi-Cap 3.8** in clean water and inject down-line from filters. After **Omni Brand Pendi-Cap 3.8** injection, flush system for a period of time sufficient to clear the line of **Omni Brand Pendi-Cap 3.8**. (If **Omni Brand Pendi-Cap 3.8** is applied during a normal irrigation cycle, make injection during the last stage.)

Chemigation Calibration (for low-volume micro sprinklers)

Calculation of use rate is based on wetted area around emitters – **NOT** on tree acres. To calculate amount of **Omni Brand Pendi-Cap 3.8**:

1. Treated area per each emitter = A
 $A = 3.14 \times (\text{radius} \times \text{radius})$
2. The area in square feet wet in each acre = B
 $B = \frac{A \times \text{emitters/acre}}{144}$
3. The total area (in square feet) wet by your system = C
 $C = B \times \text{acres covered by system}$
4. Rate per treated acre of **Omni Brand Pendi-Cap 3.8** (based on length of control desired) = R
5. Amount of **Omni Brand Pendi-Cap 3.8** to inject = S
 $S = \frac{C}{43,560} \times R = \text{qts of } \mathbf{Omni Brand Pendi-Cap 3.8}$

Example:

If the average distance from emitter to perimeter of wetted area measured 1 inch below soil surface is 13 inches, then:

$$A = 3.14 \times (13 \text{ inches} \times 13 \text{ inches})$$

$$A = 530.7 \text{ square inches}$$

If there are 300 emitters per acre, then:

$$B = \frac{530.7 \times 300}{144}$$

$$B = 1105.6 \text{ square feet wetted per acre}$$

If the system covers 20 acres, then:

$$C = 1105.6 \text{ square feet per acre} \times 20 \text{ acres}$$

$$C = 22,112 \text{ square feet wetted by system}$$

If the desired application rate per treated acre is 2.0 qts of **Omni Brand Pendi-Cap 3.8** herbicide, then:

$$S = \frac{22,112}{43,560} \times 2.0$$

$$S = 1.0 \text{ qt} = \text{amount of } \mathbf{Omni \text{ Brand Pendi-Cap 3.8}} \text{ to inject into the system}$$

Special Instructions for Chemigation

- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Recirculate and/or contain tail water (runoff water) from chemigation that contains **Omni Brand Pendi-Cap 3.8** in the field in a cistern or holding reservoir from the initial application and/or use only on adjacent, approved crops for which **Omni Brand Pendi-Cap 3.8** is registered for this type of application.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. It must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is automatically or manually shut down.
- The sprinkler chemigation system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. In addition, systems must use a metering pump, including a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- The sprinkler chemigation system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Chemigation Systems Connected to Public Water Systems

- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

- All chemigation systems connected to public water systems must also follow restrictions listed in the preceding sections involving **Chemigation**.

Application via Flood, Flooded Basin, or Gravity Flow Irrigation Systems

Omni Brand Pendi-Cap 3.8 may be applied via flood, flooded basin, or gravity flow irrigation systems, but only to the following crops: alfalfa, bearing and nonbearing fruit and nut trees, bearing and nonbearing bushberries and caneberries, bearing and nonbearing olive trees, bearing and nonbearing vineyards, bearing and nonbearing small fruit climbing vines, nonbearing fig trees, and nonbearing date palm.

Use Instructions and Restrictions for Flood, Flooded Basin, and Gravity Flow Irrigation

- **Omni Brand Pendi-Cap 3.8** may be applied through flood, flooded basin, or gravity flow irrigation systems designed to uniformly distribute irrigation water along the soil surface. Solid set systems using tall riser for overhead application are excluded.
- Follow all label directions for **Omni Brand Pendi-Cap 3.8** regarding rates per acre, timing of application, and crop-specific restrictions and precautions.
- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Helena Agri-Enterprises, LLC advises that **Omni Brand Pendi-Cap 3.8** is mixed with water at a 1:1 ratio in the injection nurse tank to assist with product flowability. Maintain agitation in the injection nurse tank to keep a uniform herbicide suspension during application. When application is complete, flush the system with water.
- Systems using a gravity-flow pesticide dispensing system must meter the pesticide in the water at the head of the field downstream of a hydraulic discontinuity, including a drop structure or weir box, to decrease potential for water source contamination from backflow water.
- Tail water (runoff water) from flood, flooded basin, or gravity flow irrigation that contains **Omni Brand Pendi-Cap 3.8** herbicide must be recirculated and/or contained in the field in a cistern or holding reservoir from the initial application and/or used only on adjacent approved crops for which **Omni Brand Pendi-Cap 3.8** is registered for this type of application.
- Systems using a pressurized water and pesticide injection system must meet the following requirements:
 - The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipe to prevent water source contamination from backflow.
 - The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent flow of fluids back toward the injection pump.
 - The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is automatically or manually shut down.
 - The system must contain a functional interlocking control to automatically shut off the pesticide injection pump when the water pump stops.
 - The irrigation pipe or water pump must include a functional pressure switch, which will stop the pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 - Systems must use a metering pump, including a positive displacement injection pump (e.g. diaphragm pump) of effective design and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
 - Any alternative to the above safety devices must conform to the list of EPA-approved alternative devices.
- Regularly measure the flow in the field to ensure the correct amount of **Omni Brand Pendi-Cap 3.8** is metered into irrigation water and also regularly monitor to ensure treated water is uniformly distributed across the field. Flow rates through metering devices and distribution of **Omni Brand Pendi-Cap 3.8** can vary with water temperature and speed of water flow across the field.
- Uniform distribution of **Omni Brand Pendi-Cap 3.8**-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness, or illegal pesticide residues in the crop.
- For questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

SPRAY DRIFT

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must not exceed 65% of the wingspan for fixed wing aircraft or 75% of the rotor diameter for helicopters. Otherwise, the boom length must not exceed 75% of the wingspan for fixed wing aircraft or 90% of the rotor diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Applicators must only apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

ADDITIVES

Spray adjuvants have little or no influence on performance of **Omni Brand Pendi-Cap 3.8** herbicide when applications are made before weed emergence. However, several tank mixes with **Omni Brand Pendi-Cap 3.8** require adjuvants to improve burndown of emerged weeds. Therefore, surfactants, liquid fertilizer (28%, 30%, or 32% UAN [urea ammonium nitrate] or AMS [ammonium sulfate]), or crop oil concentrate (COC) may be used with **Omni Brand Pendi-Cap 3.8** tank mixes applied preplant, preemergence, or early postemergence to the crop. Follow the adjuvant directions on the tank mix partner's label. The adjuvants must contain ingredients accepted by the Environmental Protection Agency.

When an adjuvant is to be used with this product, Helena Agri-Enterprises, LLC advises the use of a Chemical Producers and Distributors Association certified adjuvant.

TANK MIXING INFORMATION

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Omni Brand Pendi-Cap 3.8 may be tank mixed or applied sequentially with other herbicide products registered for use in any labeled crop found in this label for a broader spectrum of residual weed control and/or control of emerged weeds. Refer to the tank mix product labels to confirm that the respective tank mix products are registered for use on the labeled crop. Read and follow tank mix product labels for application instructions, use restrictions and precautions, and rotational cropping guidance. The most restrictive labeling applies to tank mixes. Always perform a mixing test to check the compatibility of **Omni Brand Pendi-Cap 3.8** with all potential tank mix partners.

Mixing Instructions

1. Fill tank $\frac{1}{2}$ to $\frac{3}{4}$ full with clean water or liquid fertilizer and agitate. Before mixing **Omni Brand Pendi-Cap 3.8** or **Omni Brand Pendi-Cap 3.8** tank mixes in liquid fertilizer, refer to appropriate label sections for specified uses in liquid fertilizer, application instructions, and compatibility determinations.

NOTE: Omni Brand Pendi-Cap 3.8 will **NOT** mix in high salt formulation fertilizers, including 10-34-0. When using high salt formulation fertilizers as the spray carrier, use **one** of the following:

- a. Pre-slurry **Omni Brand Pendi-Cap 3.8** in water before adding to tank; use 1:1 ratio of water to **Omni Brand Pendi-Cap 3.8**.

- b. Add water to fertilizer solution before adding **Omni Brand Pendi-Cap 3.8**. The amount of water must be equal to or more than the amount of **Omni Brand Pendi-Cap 3.8** to be used.
2. **Omni Brand Pendi-Cap 3.8 Alone**
When using **Omni Brand Pendi-Cap 3.8** alone, add **Omni Brand Pendi-Cap 3.8** to the partially filled tank while agitating; then fill the remainder of the tank with water or liquid fertilizer.
3. **Omni Brand Pendi-Cap 3.8 Tank Mixes**
Add the tank mix ingredients in the following order before adding **Omni Brand Pendi-Cap 3.8**:
 - a. **Wettable Powder (WP) formulations.** Make a slurry of the WP in water (1:2 ratio). Add the slurry slowly into the partially filled tank while agitating.
 - b. **Dry Flowable (DF)/Water-dispersible Granule (WDG) formulations.** Add the granules to the partially filled tank while agitating. Make a slurry of the granules in water before adding to liquid fertilizer.
 - c. **Flowable (F) formulations.** Add the F formulation to the partially filled tank while agitating.
 - d. Add **Omni Brand Pendi-Cap 3.8** to the partially filled tank while agitating.
 - e. **Water-soluble Concentrate (WSC) formulations.** Add the WSC formulation to the partially filled tank while agitating.
 - f. **Emulsifiable Concentrate (EC) formulations.** Add the EC formulation to the partially filled tank while agitating.

Fill the remainder of the tank with water or liquid fertilizer while agitating.
4. Thorough and continuous sprayer-tank agitation **MUST** be maintained during mixing and spraying of **Omni Brand Pendi-Cap 3.8**. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial spray cleaner according to the manufacturer's directions; then triple rinse the equipment before and after applying this product.

USE PRECAUTIONS

- **Omni Brand Pendi-Cap 3.8** will not control established weeds. Destroy emerged weeds before application.
- **Omni Brand Pendi-Cap 3.8** is most effective in controlling weeds when mechanically incorporated or when incorporated into the weed germination zone by adequate rainfall or overhead irrigation after application.
- In the event of a crop loss because of adverse weather conditions or other reasons, any crop registered for a preplant incorporated application of **Omni Brand Pendi-Cap 3.8** can be replanted without adverse effects the same year (see **Crop-specific Information** for exceptions). If replanting is necessary, do not work the soil deeper than the treated zone.
- Refer to **Crop-specific Information** for crop-specific preharvest intervals and feeding and grazing restrictions.

Use Area



ROTATIONAL CROP RESTRICTIONS

- Use of **Omni Brand Pendi-Cap 3.8** herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors,

including arid conditions, make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. Soil characteristics and environmental conditions which may contribute to crop stress that may be accentuated by the use of **Omni Brand Pendi-Cap 3.8** include: coarse soils, compaction, high salinity, eroded knolls/hilltops, cold and/or wet soils, drought, and heavy rainfall soon after application.

- When **Omni Brand Pendi-Cap 3.8** is used in tank mix or sequential combinations, refer to label of other herbicides for additional rotational crop restrictions.
- After harvest of furrow-irrigated crops, thoroughly mix the soil by plowing or deep disking to minimize the potential for herbicide carryover to the following crop.
- Refer to **Crop-specific Information** for specific rotational restrictions when **Omni Brand Pendi-Cap 3.8** is applied to specific crops.
- **Restrictions for rotational cropping after the use of Omni Brand Pendi-Cap 3.8 depend on the application use rate of Omni Brand Pendi-Cap 3.8 in the primary crop. The user must thoroughly read the following restrictions to determine the rotational crops for the specific situation, according to application use rate. For field and row crops, see the following table.**

Orchard, Grove, and Vineyard Crops

In the growing season after application of **Omni Brand Pendi-Cap 3.8** to bearing fruit and nut trees, bushberries, caneberries, or grapes, plant only those crops for which **Omni Brand Pendi-Cap 3.8** is labeled for preplant incorporated treatment or crop injury may occur. Do not rotate to other crops (except for fruit and nut trees, bushberries, caneberries, or grapes) for 24 months after **Omni Brand Pendi-Cap 3.8** application.

Table 6. Field and Row Crops

Rotational Crops	States	Omni Brand Pendi-Cap 3.8 Rate (pts/A)	Rainfall + Irrigation Amount (inches) between Omni Brand Pendi-Cap 3.8 application and rotational crop planting	Rotational Planting Interval (months) after Omni Brand Pendi-Cap 3.8 application		
				Spring	Fall	
All crops labeled for preplant incorporated application	All	> 4.0	—	the next growing season		
All other crops			—	24		
Cotton, Edible beans, Fruiting vegetables, Lentil, Peas, Peanut, Safflower, Soybean, Sunflower (and other Group 20B Oilseeds), Vegetable soybean	All	≤ 4.0	—	0		
Alfalfa stand establishment	All	≤ 4.0	> 12	6		
Wheat*, Barley*	Colorado, Idaho, Kansas, Montana, Nebraska, Nevada, Oregon, Utah, Washington, Wyoming	≤ 3.2	—	4		
		> 3.2 but ≤ 4.0	> 12	4		
			< 12	12	14	
	All other states	≤ 4.0	> 12	4		
			< 12	12	14	
Proso millet, Grain sorghum, Annual or perennial grass crops or mixtures	Minnesota, North Dakota, South Dakota	≤ 4.0	—	18	20	
		All other states	≤ 4.0	> 20	10	12
				< 20	18	20
Red beet**, Spinach**	All	≤ 4.0	> 12	12	14	
			< 12	18	20	
Sugar beet**	Nebraska, and counties Goshen, Laramie, Platte in Wyoming	≤ 2.6	> 12 and only if cropland is under center pivot irrigation	10	14	
		> 2.6 but ≤ 4.0		12	14	
		≤ 4.0		18	20	
	All other states and other counties in Wyoming	≤ 4.0		> 12	12	14
				< 12	18	20
All other crops	All	≤ 4.0	> 12	12	12	
			< 12	18	20	

* In dryland areas and/or areas where irrigation is necessary to produce the crop, Do not plant winter wheat or barley as a follow crop if crop failure/destruction occurs and land is fallowed during the summer.

** To ensure thorough mixing of soil before planting sugar beet, red beet, and spinach, plow land using a moldboard plow to a depth of 12 inches.

CROP-SPECIFIC INFORMATION

Crop Injury. **Omni Brand Pendi-Cap 3.8** herbicide use may result in crop injury, loss or damage to certain crops under a number of conditions, including but not limited to agronomic, cultural, mechanical, and environmental. Numerous risks of loss or damage to certain crops may be associated with the use of **Omni Brand Pendi-Cap 3.8** even when directions for use are followed completely. The user or grower needs to take all such risks into consideration before deciding to apply the product. **Helena Agri-Enterprises, LLC advises testing on a small portion of the target crop to determine if damage is likely to occur.** Each grower who is considering the product for such use is advised to test **Omni Brand Pendi-Cap 3.8** to determine its suitability. A grower may use **Omni Brand Pendi-Cap 3.8** only to the extent that, in his sole opinion, the benefit of **Omni Brand Pendi-Cap 3.8** use outweighs the potential injury to the grower's crop.

In addition, many factors can affect crop growth and/or yield, including but not limited to insects, diseases, weed competition, poor seed quality, improper planting depth, mechanical cultivation, poor weather (including freezing or excessive wind, rain, heat, or cold), lack of or excessive moisture, crusting, fertility, or hardpans. Risk of loss or damage to crops may be associated with the use of **Omni Brand Pendi-Cap 3.8** and contribute to poor stands because of failure of crop to emerge, swelling of roots or other belowground plant parts, less vigorous plant growth and development, and reduction in yield potential. **Omni Brand Pendi-Cap 3.8** may also cause injury to sensitive rotational crops.

AFRICAN MARIGOLD

Omni Brand Pendi-Cap 3.8 is only registered for use in African marigold grown for the purpose of lutein and zeaxanthin extraction.

Omni Brand Pendi-Cap 3.8 may be applied by ground, chemigation, or on dry bulk fertilizer.

Plant African marigold ¼-inch to ¾-inch deep and completely cover with soil. Make sure that crop seeds are below the tilled soil surface area.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied in a single application or sequentially with an interval of 30 days or more.

Uniformly apply **Omni Brand Pendi-Cap 3.8** up to 4.2 pints per acre as a broadcast treatment before target-weed germination.

Preplant Incorporated. Apply **Omni Brand Pendi-Cap 3.8** within 60 days of planting and incorporate.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting, but before crop emerges. Apply to a seedbed that is firm and free of clods.

Postemergence. Apply **Omni Brand Pendi-Cap 3.8** over-the-top postemergence after crop emergence for residual weed control.

Postemergence-directed. Apply **Omni Brand Pendi-Cap 3.8** as a postemergence-directed spray on the soil at the base of established plants, beneath plants, and between rows for residual weed control. Avoid direct contact with foliage or stems.

Postemergence and postemergence-directed treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received after application. If sufficient rainfall or irrigation does not occur to activate the herbicide, mechanically incorporate. Apply **Omni Brand Pendi-Cap 3.8** before weeds germinate or after clean cultivation to remove existing weeds because **Omni Brand Pendi-Cap 3.8** will not control emerged weeds. Apply a postemergence herbicide to control emerged weeds.

Restrictions

- Do not apply more than 4.2 pints of **Omni Brand Pendi-Cap 3.8** (2.0 lbs of pendimethalin) per acre in a single application.
- Do not apply more than a maximum cumulative total of 8.4 pints of **Omni Brand Pendi-Cap 3.8** (4.0 lbs of pendimethalin) per acre per year.
- Do not graze livestock in treated fields or feed treated plant materials to animals.

Precautions

- Avoid overlapping spray patterns because crop injury can occur.
- Avoid heavy residue on soil surface from previous year's crops.
- Avoid applying **Omni Brand Pendi-Cap 3.8** to open furrows with exposed seed.

ALFALFA (GROWN FOR FORAGE, HAY, OR SEED)

Omni Brand Pendi-Cap 3.8 may be applied by ground; air; chemigation; flood, flooded basin, and gravity flow irrigation systems; or on dry bulk fertilizer.

Use Method, Rate, and Timing

Established Alfalfa for Forage/Hay and Seed Production. Apply to established alfalfa grown for forage or hay or seed production (defined as alfalfa planted in the fall or spring which has gone through a first cutting/ mowing). Apply in a single application or in sequential applications. Uniformly apply **Omni Brand Pendi-Cap 3.8** at a broadcast rate of 1.1 to 4.2 quarts per acre before weed germination. Application can be made in the fall after the last cutting/ mowing, during winter dormancy, in the spring, or between cuttings. Apply before alfalfa reaches 6 inches in regrowth.

Seedling Alfalfa. Apply to seedling alfalfa grown for forage or hay or seed production (defined as alfalfa planted in the fall or spring which has **NOT** gone through a cutting/ mowing/seed harvest). Uniformly apply **Omni Brand Pendi-Cap 3.8** at a broadcast rate of 1.1 to 2.1 pints per acre before weed germination. Application can be made when seedling alfalfa has reached the second trifoliolate growth stage. Apply before alfalfa reaches 6 inches in growth.

Alfalfa Stand Establishment. Apply **Omni Brand Pendi-Cap 3.8** at a broadcast rate of 1.0 to 1.5 pints per acre as a preplant incorporated or preemergence treatment in direct-seeded alfalfa. Some crop stand reduction and stunting may occur with this use of **Omni Brand Pendi-Cap 3.8**; however, reduced weed competition will allow establishment of a quality stand. Use lower rates on coarse-texture soil or in lower rainfall areas (receiving less than 20 inches of rainfall and irrigation a year).

Preplant Incorporated. Uniformly incorporate **Omni Brand Pendi-Cap 3.8** herbicide into the top 2 to 3 inches of the final seedbed before planting.

Preemergence. Apply directly after drill seeding alfalfa. Plant alfalfa into a seedbed that is firm and free of clods.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Flood, Flooded Basin, and Gravity Flow Irrigation Systems. **Omni Brand Pendi-Cap 3.8** may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all directions, special instructions, and restrictions about flood, flooded basin, and gravity flow irrigation systems in the Application Instructions section of this label.

Restrictions

- Do not apply more than 4.2 quarts of **Omni Brand Pendi-Cap 3.8** (4.0 lbs of pendimethalin) per acre between spring green-up and the last cutting/ mowing of the year.
- Do not apply more than 4.2 quarts of **Omni Brand Pendi-Cap 3.8** (4.0 lbs of pendimethalin) per acre in a single application.
- For multiple applications, do not apply more than a maximum cumulative total of 6.3 quarts of **Omni Brand Pendi-Cap 3.8** (6.0 lbs of pendimethalin) per acre in any one year.
- {alternate text;}[Maximum cumulative total of 4.2 quarts of **Omni Brand Pendi-Cap 3.8** (4.0 lbs of pendimethalin) per acre in any one year in California.]
- Retreatment interval for sequential applications between the last cutting/mowing of the year and spring green-up of the following year – 28 days.
- Preharvest Interval (PHI) for the maximum dormant period application of 4.2 quarts per acre – 50 days.
- Preharvest Interval (PHI) for alfalfa forage and hay – 14 days.
- Preharvest Interval (PHI) for alfalfa seed – 90 days.

Precautions

- Some stunting and chlorosis of the alfalfa may occur with postemergence applications.

- Application made after alfalfa exceeds 6 inches in height may result in poor weed control because of possible reduced spray coverage to the soil.

ARTICHOKE

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 must be applied pre-transplant, at least 1 to 2 days before transplanting artichoke. For a single application, uniformly apply **Omni Brand Pendi-Cap 3.8** up to 3.0 pints per acre as a broadcast spray to the soil surface at least 60 days before harvest, or uniformly apply 3.1 to 8.2 pints per acre as a broadcast spray to the soil surface at least 200 days before harvest.

Restrictions

- Do not apply more than 3.0 pints (1.425 lbs of pendimethalin) per acre per year when using the 60-day preharvest interval.
- If more than 3.0 pints per acre (up to 8.2 pints per acre) of **Omni Brand Pendi-Cap 3.8** is applied, do not harvest artichoke until 200 days after application.
- Do not apply more than 8.2 pints (3.895 lbs of pendimethalin) per acre per year.
- Do not feed forage or graze livestock in treated fields.

Precautions

- Do not apply postemergence over the top of or to foliage of artichoke because severe injury may occur.

ASPARAGUS

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Apply **Omni Brand Pendi-Cap 3.8** only to established asparagus or to newly planted crown asparagus. Do not apply to newly seeded asparagus. When applying to newly planted crown asparagus, assure crowns are fully covered with 2 to 4 inches of soil.

With a single application, uniformly apply **Omni Brand Pendi-Cap 3.8** to asparagus up to 8.2 pints per acre as a broadcast spray to the soil surface at least 14 days before the first spear harvest or after seasonal harvest is complete. Application must be made before spear emergence or remove emerged spears before making the application. If asparagus is grown on sandy soils, Do not apply **Omni Brand Pendi-Cap 3.8** at more than 2.4 pts/A.

Restrictions

- Do not apply more than 8.2 pints (3.895 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 14 days.
- Do not feed forage or graze livestock in treated fields.
- Do not apply by chemigation methods.

Precautions

- Do not apply postemergence over the top of emerged spears or severe injury may occur.

BEARING AND NONBEARING FRUIT AND NUT TREES

Omni Brand Pendi-Cap 3.8 may be applied in the following individual crops within the fruit tree and tree nut crop groups:

Citrus Fruit Crop Group

Australian desert lime	Japanese summer	Orange, sour	Tangelo
Australian finger lime	grapefruit	Orange, sweet	Tangerine (mandarin)
Australian round lime	Kumquat	Pummelo	Tangor
Brown River finger lime	Lemon	Russell River lime	Trifoliate orange
Calamondin	Lime	Satsuma mandarin	Uniq fruit
Citron	Mediterranean mandarin	Sweet lime	
Citrus hybrids	Mount White lime	Tachibana orange	
Grapefruit	New Guinea wild lime	Tahiti lime	

Tree Nuts Crop Group

African nut-tree	Cashew	Hickory nut	Pequi
Almond	Chestnut	Japanese horse-chestnut	Pili nut
Beech nut	Chinquapin	Macadamia nut	Pine nut
Brazil nut	Coconut	Mongongo nut	Pistachio
Brazilian pine	Coquito nut	Monkey-pot	Sapucaia nut
Bunya	Dika nut	Monkey puzzle nut	Tropical almond
Bur oak	Ginkgo	Okari nut	Walnut, black
Butternut	Guiana chestnut	Pachira nut	Walnut, English
Cajou nut	Hazelnut (filbert)	Peach palm nut	Yellowhorn
Candlenut	Heartnut	Pecan	

Pome Fruits Crop Group

Apple	Loquat	Pear	Quince, Chinese
Azarole	Mayhaw	Pear, Asian	Quince, Japanese
Crabapple	Medlar	Quince	Tejocote

Stone Fruits Crop Group

Apricot	Cherry, tart	Plum, beach	Plum, Klamath
Apricot, Japanese	Jujube, Chinese	Plum, Canada	Plum, prune
Capulin	Nectarine	Plum, cherry	Plumcot
Cherry, black	Peach	Plum, Chickasaw	Sloe
Cherry, Nanking	Plum	Plum, Damson	
Cherry, sweet	Plum, American	Plum, Japanese	

Other Fruit Trees

Date palm ^[1]	Fig* (nonbearing only)	Olive	Pomegranate
--------------------------	------------------------	-------	-------------

[* Not for use in California.]

Omni Brand Pendi-Cap 3.8 herbicide may only be applied by ground; chemigation; or flood, flooded basin, and gravity flow irrigation systems.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied in a single application or sequentially with an interval of 30 days or more. Apply **Omni Brand Pendi-Cap 3.8** at 2.1 to 6.3 quarts per acre per application depending on the grower's weed control program, level of weed infestation, and desired use strategy, but not more than a total of 4.2 quarts/A per year in nonbearing fig, olive, pome, pomegranate, and stone fruit trees, and not more than a total of 6.3 quarts/A per year in citrus and nut trees, nonbearing date palm and nonbearing fig.

Ground Application (Bearing)

Omni Brand Pendi-Cap 3.8 may be applied surface incorporated or (surface) preemergence.

Apply **Omni Brand Pendi-Cap 3.8** broadcast or banded using ground equipment before weed germination. Apply spray directly to the ground beneath trees and/or in areas between rows. Do not apply over the top of trees with leaves, buds, or fruit. Contact by the spray mixture with leaves, shoots, or buds may cause injury or result in illegal pesticide residues on fruit.

Ground Application (Nonbearing)

Omni Brand Pendi-Cap 3.8 may be applied for preplant incorporated, preplant surface, surface incorporated, or preemergence weed control in several nonbearing fruit and nut tree crops. **Omni Brand Pendi-Cap 3.8** may be used before or after transplanting nonbearing crops.

Preplant Surface. Before transplanting, uniformly apply with ground equipment. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preplant Incorporated. Uniformly apply **Omni Brand Pendi-Cap 3.8** before transplanting but before weeds germinate. Incorporate **Omni Brand Pendi-Cap 3.8** to a depth of 1 to 2 inches. Application and incorporation must be made before transplanting to avoid mechanical injury to the crop. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preemergence. Application may be in a band or broadcast.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation and drip irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not apply **Omni Brand Pendi-Cap 3.8**-treated irrigation water over the top of trees with leaves, buds, or fruit. Contact with leaves, shoots, or buds by spray mixture may cause injury or result in illegal pesticide residues on fruit.

Flood, Flooded Basin, and Gravity Flow Irrigation Systems. **Omni Brand Pendi-Cap 3.8** may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all directions, special instructions, and restrictions about flood, flooded basin, and gravity flow irrigation systems in the Application Instructions section of this label.

Restrictions

- Do not apply more than 4.2 quarts of **Omni Brand Pendi-Cap 3.8** (4.0 lbs of pendimethalin) per acre per year in olive, pome, pomegranate, and stone fruit trees.
- Do not apply more than 6.3 quarts of **Omni Brand Pendi-Cap 3.8** (6.0 lbs of pendimethalin) per acre per year in citrus and nut trees, and nonbearing date palm, and nonbearing fig trees.
- Do not apply by air.
- Do not feed forage or graze livestock in treated groves or orchards.
- Preharvest Interval (PHI) for citrus fruit – 1 day.
- Preharvest Interval (PHI) for olive, pome, pomegranate, stone fruit, and tree nuts – 60 days.
- Do not apply to newly seeded nursery stock.

BEARING AND NONBEARING BUSHBERRIES AND CANEBERRIES

Omni Brand Pendi-Cap 3.8 herbicide may be applied in the following individual crops within the bushberry crop group:

Aronia berry	Currant, black	Huckleberry	Salal
Blueberry, highbush	Currant, red	Jostaberry	Sea buckthorn
Blueberry, lowbush	Elderberry	Juneberry (Saskatoon berry)	
Buffalo currant	European barberry	Lingonberry	
Chilean guava	Gooseberry	Native currant	
Cranberry, highbush	Honeysuckle, edible		

Omni Brand Pendi-Cap 3.8 may be applied in the following individual crops within the caneberry crop group:

Blackberry	Raspberry, black	Raspberry, wild
Loganberry	Raspberry, red	

Omni Brand Pendi-Cap 3.8 may only be applied by ground; chemigation; or flood, flooded basin, and gravity flow irrigation systems.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied in a single application or sequentially with an interval of 30 days or more. Apply **Omni Brand Pendi-Cap 3.8** at 2.1 to 6.3 quarts per acre depending on the grower's weed control program, level of weed infestation, and desired use strategy. For newly (less than 1 year) transplanted bushberry and caneberry plants, apply **Omni Brand Pendi-Cap 3.8** at between 2.1 and 3.2 quarts per acre after

transplanting. For established (at least 1 year after transplanting) plants, apply **Omni Brand Pendi-Cap 3.8** at between 2.1 and 6.3 quarts per acre.

Ground Application (Bearing)

Omni Brand Pendi-Cap 3.8 may be applied surface incorporated or (surface) preemergence.

Apply **Omni Brand Pendi-Cap 3.8** as a broadcast or banded treatment using ground equipment before weed germination. Apply spray directly to the ground beneath the bushes or canes and/or in areas between rows or trellised rows. Do not apply over the top of bushes, canes, or primocanes with leaves, buds, or fruit. Contact by the spray mixture with leaves, shoots, or buds may cause injury.

Omni Brand Pendi-Cap 3.8 may also be broadcast applied to (wild) lowbush blueberry after pruning in fall or spring but before new growth/emergence in spring, or broadcast applied over the top of dormant bushes before new growth/ emergence in spring. Do not apply broadcast over the top of (wild) lowbush blueberry if new spring growth/ emergence is imminent.

Ground Application (Nonbearing)

Omni Brand Pendi-Cap 3.8 may be applied for preplant incorporated, preplant surface, surface incorporated, or preemergence weed control in nonbearing bushberry and caneberry crops. **Omni Brand Pendi-Cap 3.8** may be applied before transplanting or applied after transplanting and the establishment of the nonbearing crops. **Omni Brand Pendi-Cap 3.8** may be used after transplanting the nonbearing crops and after the ground is settled and no soil cracking is present at 2.1 to 3.2 quarts per acre.

Preplant Surface. Before transplanting, uniformly apply with ground equipment. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preplant Incorporated. Uniformly apply **Omni Brand Pendi-Cap 3.8** before transplanting but before weeds germinate. Incorporate **Omni Brand Pendi-Cap 3.8** to a depth of 1 to 2 inches. Application and incorporation must be made before transplanting to avoid mechanical injury to the crop. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preemergence. Application may be in a band or broadcast.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation and drip irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not apply **Omni Brand Pendi-Cap 3.8**-treated irrigation water over the top of bushes, canes, or primocanes with leaves, buds, or fruit. Contact with leaves, shoots, or buds by spray mixture may cause injury.

Flood, Flooded Basin, and Gravity Flow Irrigation Systems Application. **Omni Brand Pendi-Cap 3.8** may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all directions, special instructions, and restrictions about flood, flooded basin, and gravity flow irrigation systems in the Application Instructions section of this label.

Restrictions

- Do not apply more than 6.3 quarts of **Omni Brand Pendi-Cap 3.8** (6.0 lbs of pendimethalin) per acre per year.
- Do not apply by air.
- Do not apply to newly (less than 1 year) transplanted bushberries and caneberries until ground is settled and no soil cracking is present.
- Do not apply more than 3.2 quarts of **Omni Brand Pendi-Cap 3.8** (3.0 lbs of pendimethalin) per acre per year in newly (less than 1 year) transplanted bushberries and caneberries.
- Do not feed forage or graze livestock in treated field or planting.
- Preharvest Interval (PHI) – 30 days.
- Do not apply to newly seeded nursery stock.

Precautions

- **Omni Brand Pendi-Cap 3.8** inhibits root growth and development. Use is not advised if bushberry and caneberry roots are shallow or if root growth is inhibited or stressed due to pathogens, insects, or environmental factors.

BEARING AND NONBEARING GRAPE

Omni Brand Pendi-Cap 3.8 herbicide may be only applied by ground; chemigation; or flood, flooded basin, and gravity flow irrigation systems.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied in a single application or sequentially with an interval of 30 days or more. Uniformly apply **Omni Brand Pendi-Cap 3.8** in grape vineyards at 2.1 to 6.3 quarts per acre depending on the grower's weed control program, level of weed infestation, and desired use strategy.

Omni Brand Pendi-Cap 3.8 may be applied anytime after fall harvest, during winter dormancy, in spring, and in season before harvest.

Ground Application (Bearing)

Omni Brand Pendi-Cap 3.8 may be applied surface incorporated or (surface) preemergence.

Apply **Omni Brand Pendi-Cap 3.8** broadcast or banded using ground equipment before weed germination. Apply spray directly to the ground beneath grape vines and/or in areas between rows. Do not apply over the top of grape vines with leaves, buds, or fruit. Contact with leaves, shoots, or buds by the spray mixture may cause injury or result in illegal pesticide residues on fruit.

Ground Application (Nonbearing)

Omni Brand Pendi-Cap 3.8 may be applied for preplant incorporated, preplant surface, surface incorporated, or preemergence weed control in nonbearing vineyards. **Omni Brand Pendi-Cap 3.8** may be used before or after transplanting.

Preplant Surface. Before transplanting, uniformly apply with ground equipment. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preplant Incorporated. Uniformly apply **Omni Brand Pendi-Cap 3.8** before transplanting but before weeds germinate. Incorporate **Omni Brand Pendi-Cap 3.8** to a depth of 1 to 2 inches. Application and incorporation must be made before transplanting to avoid mechanical injury to the crop. Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preemergence. Application may be in a band or broadcast.

Nonbearing Grape

Newly Transplanted and One-year-old Grapevines:

- Do not allow spray to contact buds or leaves or leaf distortion may occur.
- Do not apply to newly transplanted vines until ground has settled and no cracks are present.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation and drip irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not apply **Omni Brand Pendi-Cap 3.8**-treated irrigation water over the top of grape vines with leaves, buds, or fruit. Contact with leaves, shoots, or buds by spray mixture may cause injury or result in illegal pesticide residues on fruit.

Flood, Flooded Basin, and Gravity Flow Irrigation Systems. **Omni Brand Pendi-Cap 3.8** may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all directions, special instructions, and restrictions about flood, flooded basin, and gravity flow irrigation systems in the Application Instructions section of this label.

Restrictions

- Do not apply over the top of grape vines with leaves, buds, or fruit.
- Do not apply by air.
- Do not apply more than 6.3 quarts (6.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 21 days.
- **Omni Brand Pendi-Cap 3.8**-treated vineyards (post-harvest) may be grazed by livestock.

BEARING AND NONBEARING SMALL FRUIT CLIMBING VINES

Omni Brand Pendi-Cap 3.8 may be applied in the following small fruit climbing vines: Amur river grape, gooseberry, fuzzy kiwifruit, hardy kiwifruit, maypop, and Schisandra berry.

Omni Brand Pendi-Cap 3.8 may be only applied by ground; chemigation; flood, flooded basin, and gravity flow irrigation systems.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied in a single application or sequentially with an interval of 30 days or more. Uniformly apply **Omni Brand Pendi-Cap 3.8** in small fruit climbing vines at 2.1 to 4.2 quarts per acre depending on the grower's weed control program, level of weed infestation, and desired use strategy.

Omni Brand Pendi-Cap 3.8 may be applied anytime after fall harvest, during winter dormancy, and in spring.

Ground Application

Omni Brand Pendi-Cap 3.8 may be applied surface incorporated or (surface) preemergence.

Apply **Omni Brand Pendi-Cap 3.8** broadcast or banded using ground equipment before weed germination. Apply spray directly to the ground beneath small fruit climbing vines and/or in areas between rows. Do not apply over the top of small fruit climbing vines with leaves, buds, or fruit. Contact with leaves, buds, or fruit by the spray mixture may cause injury or result in illegal pesticide residues on fruit.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation and drip irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not apply **Omni Brand Pendi-Cap 3.8**-treated irrigation water over the top of small fruit climbing vines with leaves, buds, or fruit. Contact with leaves, buds, or fruit by the spray mixture may cause injury or result in illegal pesticide residues on fruit.

Flood, Flooded Basin, and Gravity Flow Irrigation Systems. **Omni Brand Pendi-Cap 3.8** herbicide may be applied in flood, flooded basin, and gravity flow irrigation systems. Follow all directions, special instructions, and restrictions about flood, flooded basin, and gravity flow irrigation systems in the Application Instructions section of this label.

Restrictions

- Do not apply over the top of small fruit climbing vines with leaves, buds, or fruit.
- Do not apply by air.
- Do not apply more than 4.2 quarts (4.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 60 days.
- Do not feed forage or graze livestock in treated vines.
- Do not apply **Omni Brand Pendi-Cap 3.8** when impregnated onto dry bulk fertilizer in small fruit climbing vines.

BRASSICA HEAD AND STEM VEGETABLES

Omni Brand Pendi-Cap 3.8 may only be applied to the following Brassica head and stem vegetables:

Broccoli	Chinese broccoli	Cauliflower
Brussels sprouts	Chinese cabbage (napa)	Cavalo broccolo
Cabbage	Chinese mustard cabbage	Kohlrabi

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Uniformly apply **Omni Brand Pendi-Cap 3.8** to Brassica head and stem vegetables by ground or air as a preplant surface application before transplanting or as a postemergence application.

Uniformly apply **Omni Brand Pendi-Cap 3.8** only by ground as a postemergence-directed application to transplanted or established direct-seeded Brassica head and stem vegetables.

Do not apply before direct-seeded Brassica head and stem vegetables.

With a single application, apply up to 2.1 pints per acre of **Omni Brand Pendi-Cap 3.8** to Brassica head and stem vegetables as a broadcast or banded spray to the soil surface at pretransplant time, as a broadcast postemergence foliar spray, or as a postemergence-directed spray between vegetable rows. Apply postemergence or postemergence-directed to 2-leaf to 4-leaf vegetable transplants at 1 to 3 days after transplanting, or to the 2-leaf to 4-leaf stage of direct seeded vegetable plants.

Apply **Omni Brand Pendi-Cap 3.8** as a postemergence-directed spray on the soil, beneath plants, and between vegetative rows.

Do not spray foliage or stems because crop injury will occur. Roots of transplants must be established. Following the postemergence-directed application if sufficient rainfall or irrigation does not occur, mechanically incorporate to activate the herbicide. Apply **Omni Brand Pendi-Cap 3.8** before weed germination. Emerged weeds will not be controlled by this treatment.

Use Rate

Pretransplant, Postemergence, Postemergence-directed

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.0 to 1.5
Medium	1.5 to 2.1
Fine	

Restrictions

- Do not apply more than 2.1 pints (1.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) for broccoli – 60 days.
- Preharvest Interval (PHI) for cabbage and other Brassica head and stem vegetables – 70 days.
- Do not feed forage or graze livestock in treated fields.
- Do not apply via chemigation methods.
- Not for use in California.

Precautions

- Avoid root contact with **Omni Brand Pendi-Cap 3.8**-treated soil when placing transplants into furrow or hole, or crop injury may occur.
- Avoid overlapping spray patterns because crop injury can occur.

CARROT

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or chemigation.

Use Method, Rate, and Timing

Preemergence. Make a single broadcast application by ground, air, or chemigation at 2.0 pints per acre of

Omni Brand Pendi-Cap 3.8 as a postplant treatment before emergence of the crop and before weed germination. Apply as a preemergence treatment within 2 days after planting.

Layby. **Omni Brand Pendi-Cap 3.8** may be applied only by ground equipment at layby (last mechanical cultivation) at 2.0 pints per acre as a directed spray to the soil between rows. Apply **Omni Brand Pendi-Cap 3.8** before weed germination. Emerged weeds will not be controlled by this treatment. Do not allow the spray to contact carrot plants or injury may occur.

Do not apply layby applications by chemigation or air.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not allow **Omni Brand Pendi-Cap 3.8**-treated irrigation water to contact carrot plants.

Do not apply tank mixes through any type of irrigation system unless the label instructions on chemigation of all products are followed.

Restrictions

- Do not apply more than 2.0 pints (1.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 60 days.
- Do not feed forage or graze livestock in treated fields.
- Do not apply as a broadcast spray over top of carrots or crop injury may result.
- Do not apply layby applications by chemigation or air.

CARROT GROWN FOR SEED PRODUCTION

Omni Brand Pendi-Cap 3.8 herbicide may be applied only by layby with ground equipment.

Use Method, Rate, and Timing

Last Cultivation (Layby). Apply **Omni Brand Pendi-Cap 3.8** after the last normal mechanical cultivation (layby) at a rate of 1.0 to 4.0 pints per acre (on a broadcast basis). Uniformly apply as a directed spray to the soil between rows. Do not allow the spray to contact carrot plants or injury may occur. Use protective shields to avoid contact with carrot foliage. Use calibrated nozzles and equipment.

Layby application can be made to carrots previously treated with herbicides registered in/on carrots. Consult the labels of those herbicides for suggested treatments, rates, and precautions or restrictions for use in carrots and for rotational crop restrictions.

Restrictions

- Do not apply as a broadcast spray over top of carrots or crop injury may result.
- Do not apply layby applications by chemigation or air.
- Do not apply within 60 days before carrot seed harvest.
- Do not feed forage or graze livestock in treated fields.
- Do not harvest carrots for food or feed use.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/nonfood use. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, shall provide a copy of this pesticide label to the seed conditioner.

Consequently, no portion of this carrot seed crop, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, foliage, and seed screenings, may be used or distributed for food or feed purposes.

Processed carrot seed from a field treated with this product must bear a specific tag or conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading, with the following statement: "Not for human consumption or animal feed." All seed screenings from seed processing shall be disposed of in such a manner that the screenings cannot be distributed or used for human food or animal feed purposes.

The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter, or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.

CELERY (AND OTHER LEAF PETIOLE VEGETABLES IN GROUP 22B)

Omni Brand Pendi-Cap 3.8 may be applied to celery and the following other leaf petiole vegetables of subgroup 22B:

Cardoon	Fuki	Udo
Chinese celery	Rhubarb	Zuiki

Omni Brand Pendi-Cap 3.8 may only be applied by ground or air.

Use Method, Rate, and Timing

Pre-transplant. Uniformly apply **Omni Brand Pendi-Cap 3.8** as a broadcast spray to celery and other leaf petiole vegetables as a preplant surface application before transplanting. Do not incorporate **Omni Brand Pendi-Cap 3.8** into soil before transplanting.

Do not apply before direct-seeded celery and other leaf petiole vegetables.

Post-transplant. Uniformly apply **Omni Brand Pendi-Cap 3.8** as a postemergence broadcast spray to celery and other leaf petiole vegetables 1 or more days after transplanting until no less than 60 days before harvest.

Sequential applications. **Omni Brand Pendi-Cap 3.8** may be applied in sequential applications with pretransplant followed by post-transplant. Separate applications by at least 21 days and Do not exceed the maximum cumulative rate for soil type per year.

Use Rate

Pretransplant, Postransplant

Soil Texture	Broadcast Rate (pts/A)
Coarse (mineral)	2.1 to 3.2
Medium (mineral)	2.1 to 3.2
Fine (mineral)	2.1 to 3.2
Muck	4.0

Restrictions – Mineral Soils

- Do not apply more than 3.2 pts (1.5 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre in a single application.
- Do not apply more than 6.4 pts (3.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) – 60 days.

Restrictions – Muck Soils

- Do not apply more than 4.0 pts (1.9 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** herbicide per acre in a single application.
- Do not apply more than 8.0 pts (3.8 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) – 60 days.

Precautions

- Avoid root contact with **Omni Brand Pendi-Cap 3.8**-treated soil when placing transplants into furrow or hole, or crop injury may occur.
- Before applying **Omni Brand Pendi-Cap 3.8** to transplanted celery and other leaf petiole vegetables, roots of plants must be established and plants must have regained full turgidity.

CORN (FIELD, FIELD SEED, FRESH SWEET, POPCORN, AND POPCORN SEED)

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer. **Omni Brand Pendi-Cap 3.8** may be applied in conventional tillage, minimum tillage, or no-till as a preemergence, postemergence, or postemergence incorporated (CULTI-SPRAY) application in field corn.

Omni Brand Pendi-Cap 3.8 may be applied in conventional tillage as a preemergence or postemergence application in field seed corn, popcorn, popcorn seed corn, and fresh sweet corn.

Regardless of tillage system, plant corn at least 1½ inches deep and completely cover with soil.

In conventional-tillage systems, plant into a seedbed that is firm and free of clods and trash. Use only where tillage provides good soil coverage of corn seed.

In no-till systems, use a no-till planter capable of planting through crop residue. Use of no-till planters under conditions that do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if **Omni Brand Pendi-Cap 3.8** contacts the germinating corn seed. Check equipment to ensure good seed coverage.

Additional Weeds Controlled. In addition to weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** controls the following weeds in corn with CULTI-SPRAY application: wild proso millet and shattercane.

Use Method, Rate, and Timing

Preemergence. Apply after planting but before weeds germinate and crop emerges.

Postemergence. Apply postemergence until field corn is 30-inches tall (20-inches to 24-inches tall for popcorn, popcorn seed, field seed, and fresh sweet corn) or in the V8 growth stage, whichever is more restrictive. If the corn canopy prevents applications from reaching the soil, use drop nozzles and apply as a directed spray.

CULTI-SPRAY. Apply **Omni Brand Pendi-Cap 3.8** alone or **Omni Brand Pendi-Cap 3.8** plus atrazine when field corn is at least 4-inches tall until last cultivation (layby). **Omni Brand Pendi-Cap 3.8** plus atrazine must be applied before the field corn reaches 12 inches in height.

See specific directions for (CULTI-SPRAY) application under Application Timings.

Do not apply more than 1.2 lbs ai per acre of atrazine, as specified on the atrazine label. Under situations of low rainfall or soil moisture, when deep germinating weeds like shattercane or field sandbur are anticipated, mechanical incorporation provides the best results. If cultivation is needed after application and incorporation of **Omni Brand Pendi-Cap 3.8**, ensure the depth of cut is no deeper than the depth of cut used to incorporate.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Use Rate

Preemergence, Postemergence

Soil Texture	Organic Matter		
	< 1.5% (pts/A)	1.5% to 3.0% (pts/A)	> 3.0% (pts/A)
Coarse	2.0	3.0	3.0
Medium	3.0	3.0	4.0
Fine	3.0	4.0	4.0

CULTI-SPRAY (Field Corn ONLY)

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹ (pts/A)
Coarse	1.5	2.0
Medium	2.0	3.0
Fine	3.0	3.0

¹ See Use Precautions for map of specific states.

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** in reduced-tillage, minimum-tillage, or no-till fresh sweet corn, seed corn, or popcorn.
- Do not apply **Omni Brand Pendi-Cap 3.8** in no-till in California.
- Do not apply preplant incorporated.
- Do not apply postemergence in liquid fertilizer.
- Livestock can graze or be fed forage from treated corn 21 days or more after application.
- **Omni Brand Pendi-Cap 3.8** may be applied sequentially in a single year as long as the total use rate applied in the year does not exceed the highest rate per acre for any given soil type.

COTTON

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer to cotton (including those varieties with a herbicide-resistant trait) grown under conventional tillage, minimum-tillage, or no-till systems, or on stale seedbeds.

Additional Weeds Suppressed. In addition to weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** herbicide will suppress Russian thistle in the state of Arizona.

Use Method, Rate, and Timing

Fall Application. **Omni Brand Pendi-Cap 3.8** may be applied for weed control in cotton in the fall, after October 15 (up to 140 days before planting cotton) in Arizona, California, Louisiana, Mississippi, New Mexico, Oklahoma, and Texas. Apply **Omni Brand Pendi-Cap 3.8** at the broadcast rate of 2.0 pints per acre on coarse or medium soils and 3.0 pints per acre on fine soils.

Preplant Surface. Apply **Omni Brand Pendi-Cap 3.8** within 15 days of planting.

Preplant Incorporated. Apply **Omni Brand Pendi-Cap 3.8** within 60 days of planting and incorporate.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting. Apply to a seedbed that is firm and free of clods.

Preplant Incorporated followed by Preemergence. Apply **Omni Brand Pendi-Cap 3.8** within 60 days of planting and incorporate. Apply overlay application of **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting. Total amount of **Omni Brand Pendi-Cap 3.8** applied per acre cannot exceed the highest labeled rate for a given soil type. Preplant incorporated and preemergence applications of **Omni Brand Pendi-Cap 3.8** may be applied with the labeled tank mix herbicide(s).

Layby Application (at last cultivation). Apply **Omni Brand Pendi-Cap 3.8** directly to the soil between rows as a directed spray after the last normal cultivation (layby). Layby applications can be applied in cotton previously treated with **Omni Brand Pendi-Cap 3.8** or any herbicide(s) registered for use in cotton. The total amount of **Omni Brand Pendi-Cap 3.8** applied per acre per year cannot exceed the highest labeled rate for a given soil type. **Omni Brand Pendi-Cap 3.8** may also be applied at layby in cotton with herbicide-resistant traits.

Postemergence. **Omni Brand Pendi-Cap 3.8** may be applied by ground or air as a broadcast over-the-top postemergence application in cotton, including those varieties with a herbicide-resistant trait.

Postemergence treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received after application. Apply **Omni Brand Pendi-Cap 3.8** before weeds germinate or after clean cultivation to remove existing weeds because **Omni Brand Pendi-Cap 3.8** will not control emerged weeds. Apply a postemergence herbicide to control emerged weeds.

Apply **Omni Brand Pendi-Cap 3.8** broadcast postemergence (in water) over the top of cotton after cotton reaches the 4-leaf growth stage, but not after the 8-leaf growth stage. Over the-top applications made before the 4-leaf growth stage or after the 8-leaf growth stage may result in crop injury and/or yield loss.

Omni Brand Pendi-Cap 3.8 may also be tank mixed with herbicide products registered for use on cotton varieties with a herbicide-resistant trait. When tank mixing **Omni Brand Pendi-Cap 3.8** with another herbicide product, always follow the most restrictive labeling pertaining to application rates, methods, timings and precautions. Do not tank mix and apply over-the-top postemergence with fluometuron, glyphosate plus s-metolachlor, prometryn, or pyriithiobac-sodium, or s-metolachlor.

Dry ammonium sulfate (AMS) at 17 lbs/100 gallons of spray solution must be used when tank mixing **Omni Brand Pendi-Cap 3.8** with glyphosate-containing products. Liquid AMS may also be used, but must be used at an equivalent rate to 17 lbs of dry weight AMS/100 gallons of spray solution. A nitrogen replacement may not be used with this tank mix unless specified as acceptable from Helena Agri-Enterprises, LLC in writing. An appropriate mixing order is as follows: fill tank to at least ½ full with water; then add in order: AMS, **Omni Brand Pendi-Cap 3.8**, glyphosate; then fill the tank to capacity with water.

Over-the-top postemergence application of **Omni Brand Pendi-Cap 3.8** can be applied in cotton previously treated with at-planting soil applications of **Omni Brand Pendi-Cap 3.8** or any other soil-applied herbicide(s) registered for use in cotton. Consult the labels of those herbicides for suggested treatments, rates, precautions, or restrictions for use in cotton, and for rotational crop restrictions. Follow the most restrictive label instructions when using products in combination with soil-applied **Omni Brand Pendi-Cap 3.8**.

Precautions

- Postemergence application of **Omni Brand Pendi-Cap 3.8** may cause temporary growth reduction and/or leaf discoloration or malformation of cotton after application.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Use Rate

Preplant, Preemergence, Layby

Soil Texture	Conventional or Minimal Tillage (pts/A)	No-till** (pts/A)
Coarse	1.0 to 2.0*	2.0
Medium	2.0	3.0

Fine	3.0	4.0
* Do not exceed 1.6 pts/A on coarse-texture soils in California.		
** Do not use on soils with more than 3% organic matter.		

Postemergence

Omni Brand Pendi-Cap 3.8 herbicide Alone or in Tank Mix

Soil Texture	Conventional, Minimum, or No Tillage (pts/A)
Coarse	1.0 to 2.0*
Medium	1.5 to 2.0
Fine	2.0
* Do not exceed 1.6 pts/A on coarse-texture soils in California.	

Restrictions

- Do not apply postemergence in cotton in California.
- Do not apply **Omni Brand Pendi-Cap 3.8** in no-till in California.
- Do not apply more than the highest yearly rate per acre for any given soil type.
- Do not apply more than 2.0 pts/A of **Omni Brand Pendi-Cap 3.8** (0.95 lb active ingredient/A) when applied postemergence to cotton for any given soil type.
- Preharvest Interval (PHI) – 60 days.
- In treated cotton fields, forage may be fed to or grazed by livestock.
- Do not apply more than the maximum cumulative yearly rate of 4.2 pts of **Omni Brand Pendi-Cap 3.8** per acre (2 lbs ai/A) for combined preplant/preemergence and postemergence applications.
- Do not apply over the top in fluid fertilizer.
- Do not apply in tank mix with any adjuvant, surfactant, oil, or other pesticide (except for cotton insecticides).
- Do not apply in any manner except as described in this label or crop injury and/or yield reduction may occur.
- Do not apply if cotton is under stress (including stress related to previous pesticide treatments, poor fertilization, environmental conditions, and/or pest damage) at time of application. If cotton is under stress (including stress related to previous pesticide treatments, poor fertilization, environmental conditions, and/or pest damage) at time of application, **Omni Brand Pendi-Cap 3.8** may retard cotton recovery and/or adversely affect yield.

DRY BULBS (GARLIC, ONIONS, SHALLOTS)

Omni Brand Pendi-Cap 3.8 may be applied to the following dry bulb crops:

Daylily, bulb	Garlic, serpent, bulb	Onion, pearl
Fritillaria, bulb	Lily, bulb	Onion, potato, bulb
Garlic, bulb	Onion, bulb	Shallot, bulb
Garlic, great-headed, bulb	Onion, Chinese, bulb	

Omni Brand Pendi-Cap 3.8 may be applied to direct-seeded and transplanted dry bulb onions and dry bulb shallots.

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or chemigation.

Use Method, Rate, and Timing Mineral Soils

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5
Medium	2.0
Fine	3.2

Preemergence. After garlic planting but before crop and weeds emerge.

Postemergence. When garlic is in the 1st to 5th true-leaf growth stage.

Split Application. In garlic at both preemergence and postemergence timings.

In all states except California, apply **Omni Brand Pendi-Cap 3.8** as a broadcast treatment when dry bulb onions or dry bulb shallots have 2 to 9 true leaves.

In California, apply **Omni Brand Pendi-Cap 3.8** only as a single application when dry bulb onions or dry bulb shallots have 2 to 6 true leaves.

Additional State-specific Instructions

Additional Use in Colorado, Kansas, and Nebraska

- **Omni Brand Pendi-Cap 3.8** may be applied sequentially in seeded dry bulb onions. Make first application of **Omni Brand Pendi-Cap 3.8** at loop stage. Make sequential application of **Omni Brand Pendi-Cap 3.8** early postemergence (2nd to 9th true-leaf stage). Do not apply more than the maximum labeled rate for a given soil texture. Do not apply **Omni Brand Pendi-Cap 3.8** at loop stage through the 9th true-leaf stage if heavy rains are expected or severe crop injury may result.

Additional Use in Colorado and the High Plains of Texas

- For transplanted dry bulb onions only, apply and shallow incorporate (less than 2-inches deep) **Omni Brand Pendi-Cap 3.8** into preformed beds before transplanting.

Additional Use in Idaho, Oregon, and Washington

- Apply **Omni Brand Pendi-Cap 3.8** as a broadcast treatment when dry bulb onions or dry bulb shallots are between the flag leaf to 9th true-leaf stage.
- **Omni Brand Pendi-Cap 3.8** may be used at 3.0 to 4.0 pints per acre for dodder control on medium-texture and fine-texture soils.
- Do not apply **Omni Brand Pendi-Cap 3.8** using chemigation at the dodder control rate.
- **Omni Brand Pendi-Cap 3.8** may be applied in the fall or spring to the furrow area of land bedded in the fall in preparation for planting seed of dry bulb onions the following spring.
- Apply **Omni Brand Pendi-Cap 3.8** as a banded application at rates based on appropriate soil texture. Band width is ½ the width of the row spacing. Keep **Omni Brand Pendi-Cap 3.8** away from the area where dry bulb onion seed will be planted.
- Harrow off tops of beds after **Omni Brand Pendi-Cap 3.8** furrow application before planting dry bulb onions.
- For selective weed control in the onion row, apply **Omni Brand Pendi-Cap 3.8** herbicide as a banded postemergence application to flag-leaf dry bulb onions at the labeled rates based on soil texture. Apply **Omni Brand Pendi-Cap 3.8** only once to the furrow area and once to the dry bulb onion row as a postemergence application.

Additional Use in Michigan

- For mineral soils containing more than 10% organic matter, follow the directions for muck soils (see following).

Restrictions (Mineral Soils)

- Do not mechanically incorporate except as specified for use on dry bulb onions in Colorado and the Texas High Plains.
- Do not apply more than 3.2 pints (1.5 lbs of pendimethalin) per acre per year (except as described above for Idaho, Oregon, and Washington).
- Preharvest Interval (PHI) – 60 days in California; 45 days in all other states.
- Do not feed or graze these crops.
- Do not apply **Omni Brand Pendi-Cap 3.8** preemergence through the loop stage if heavy rains are expected or severe crop injury may result. If irrigating immediately after **Omni Brand Pendi-Cap 3.8** application at the preemergence through loop stage, do not irrigate more than ½ inch of water.

Use Method, Rate, and Timing Muck Soils

Omni Brand Pendi-Cap 3.8 may be applied sequentially in dry bulb onions or dry bulb shallots on muck soils, only once preemergence and only twice postemergence, as follows:

Application Timing and Growth Stage	Rate (pts/A)
--	-------------------------

Preemergence through Loop Stage	4.0
Early Postemergence (2nd to 6th true-leaf stage)	4.0
Late Postemergence (6th to 9th true-leaf stage)	4.0

Restrictions (Muck Soils)

- Do not apply to muck soils in California.
- Preharvest Interval (PHI) – 45 days.
- Do not feed or graze these crops.
- Do not apply more than 12.6 pints per acre per year on muck soils. To maximize crop safety, ensure good soil coverage during planting or transplanting and delay preemergence applications to the loop stage, if possible.
- Do not apply **Omni Brand Pendi-Cap 3.8** preemergence through the loop stage if heavy rains are expected or severe crop injury may result. If irrigating immediately after **Omni Brand Pendi-Cap 3.8** application at the preemergence through loop stage, do not irrigate more than ½ inch of water.
- Do not plant sugar beets, red beets, spinach, winter wheat, or winter barley as rotational crops on muck soils for 12 months from the time of last application if more than 3.2 pints per acre of **Omni Brand Pendi-Cap 3.8** is applied to the onion crop.
- If loss of onion crop occurs, do not replant any crop other than onions in muck soil during the same cropping year and do not work the soil deeper than 2 inches.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Do not irrigate more than ½ inch of water. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

EDIBLE BEANS

Omni Brand Pendi-Cap 3.8 may be applied to the following edible beans: dry beans [navy, great northern, red kidney, black, black turtle, cranberry, small white type, guar, adzuki, broad, faba, fava, field, lima (dry), pinto], lima bean, snap bean, chickpea (garbanzo bean), southern pea (cowpea), and sweet lupins.

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Omni Brand Pendi-Cap 3.8 may only be applied (fall) preplant surface or preplant incorporated in dry beans, lima bean, snap bean, and southern pea (cowpea). **Omni Brand Pendi-Cap 3.8** may be applied (fall) preplant surface or preplant incorporated or (spring) preplant surface in chickpea (garbanzo bean). **Omni Brand Pendi-Cap 3.8** may be applied (fall) preplant surface or preplant incorporated or preemergence in sweet lupins.

Use Method, Rate, and Timing Fall Application

For use only in Idaho, Minnesota, Montana, North Dakota, Oregon, South Dakota, Washington, and Wyoming). Apply **Omni Brand Pendi-Cap 3.8** preplant surface or preplant incorporated (rainfall, irrigation, or mechanically) in late fall before planting edible beans [chickpea (garbanzo bean), dry beans, lima bean, snap bean, southern pea (cowpea), and sweet lupins] the following spring. Apply **Omni Brand Pendi-Cap 3.8** in the late fall when soil temperatures are 45°F or below but before the ground freezes.

Do not apply when the air temperature is below 45°F.

Rainfall or irrigation is required for incorporation and activation. Unpredictable weed control can be expected because factors like length of time between application and planting as well as uncontrollable weather factors will determine herbicide activity and longevity.

Use Rate (Fall)¹

Soil Texture	Broadcast Rate < 3% Organic Matter (pts/A)	Broadcast Rate > 3% Organic Matter (pts/A)
Coarse	2.0	2.0
Medium	2.5	3.0
Fine	3.0	3.0

¹ Use limited to certain states. Follow state-specific instructions and/or restrictions.

Preplant Incorporated. Apply within 60 days of planting and incorporate.

Preemergence. Apply only to sweet lupins at planting or up to 2 days after planting. Apply to a seedbed that is firm and free of clods.

Use Rate

Preplant Incorporated, Preemergence

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹	
		< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	2.0	2.0
Medium	2.0	2.5	3.0
Fine	3.0	3.0	3.0

¹ See Use Precautions for map of specific states.

State-specific Instructions

Idaho, Montana, North Dakota, Oregon, and Washington

Omni Brand Pendi-Cap 3.8 herbicide may be applied to chickpeas grown in no-tillage and/or minimum tillage systems in Idaho, Montana, North Dakota, Oregon, and Washington. Preplant surface applications must be made within 30 days of planting. Do not apply more than 1.5, 2.0, and 3.0 pts/A of **Omni Brand Pendi-Cap 3.8** in coarse, medium, and fine texture soils, respectively. When planting, ensure the seed furrow is fully closed because conditions that allow the seed furrow to inadequately close and/or allow **Omni Brand Pendi-Cap 3.8** to contact the seed may result in crop injury. Certain unfavorable environmental conditions, including cool temperatures, excessive moisture after application, and wet and/or compacted soil conditions, may result in delayed emergence and stunting with **Omni Brand Pendi-Cap 3.8** use in chickpea. Adequate rainfall or irrigation after application before weed seedling germination provides the most effective weed control. Herbicide performance from surface application may be decreased compared to soil incorporated application.

Idaho, Oregon, and Washington

Omni Brand Pendi-Cap 3.8 may be applied postplant preemergence only to chickpea grown in conventional tillage systems in Idaho, Oregon, and Washington. Application must be made within 2 days of planting. Apply up to but not more than 1.5 pts/A. Apply to a firm seedbed free of clods. Soil conditions that cause poor seed furrow closure and coverage may result in delayed emergence and stunting of the crop. Under certain environmental conditions, including cool temperatures, excessive moisture after application and wet soil conditions may result in delayed emergence and stunting with **Omni Brand Pendi-Cap 3.8** use in chickpea. Adequate rainfall or irrigation after application before weed seedling germination provides the most effective weed control.

Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Oregon, Washington, and Wyoming

Apply **Omni Brand Pendi-Cap 3.8** by ground postplant preemergence to dry beans grown under sprinkler irrigation in Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Oregon, Washington, and Wyoming. Dry beans must have a minimum planting depth of 2 inches. Before applying **Omni Brand Pendi-Cap 3.8** to dry beans, growers are advised to check with their local seed company or seed supplier for sensitive varieties and to verify the selectivity of **Omni Brand Pendi-Cap 3.8** on the grower's specific dry bean variety. **Omni Brand Pendi-Cap 3.8** application made postplant preemergence to dry beans must be immediately followed by 0.50 to 0.75 inch of water from overhead irrigation/rainfall. Apply **Omni Brand Pendi-Cap 3.8** within 1 to 4 days of planting and up to but not more than 2.0 pts/A. Apply to a firm seedbed free of clods. Soil conditions that cause poor seed furrow closure and coverage may result in delayed emergence and stunting of the crop. Do not apply as a chemigation application. Do not apply **Omni Brand Pendi-Cap 3.8** in tank mix with halosulfuron or flumioxazin as a preemergence application to dry beans because of unacceptable crop response. Under certain environmental conditions, including cool temperatures, excessive moisture after application and wet soil conditions may result in crop injury, delayed emergence, and/or stunting with **Omni Brand Pendi-Cap 3.8** use in dry beans. Adequate rainfall or irrigation after application before weed seedling germination provides the most effective weed control.

Restrictions

- Do not feed lupin hay and forage or graze livestock in treated lupin fields.
- Do not apply **Omni Brand Pendi-Cap 3.8** more than once per year.
- Do not apply in any type of irrigation system.

FALLOW

Omni Brand Pendi-Cap 3.8 may be applied to fallow ground following crop harvest as a planned residual treatment to control labeled broadleaf and grass weeds as they germinate.

Omni Brand Pendi-Cap 3.8 may be applied to fallow ground by ground, air, or chemigation.

Use Method, Rate, and Timing

Apply as a broadcast spray at rates up to but not more than 3.0 pts/A of **Omni Brand Pendi-Cap 3.8**. Emerged weeds will not be controlled by this treatment. **Omni Brand Pendi-Cap 3.8** must be applied with a tank mix partner (i.e. glyphosate) for control of emerged weeds.

Do not make more than one application of **Omni Brand Pendi-Cap 3.8** during a single fallow period.

Do not apply **Omni Brand Pendi-Cap 3.8** to fallow ground after July 1 if treated fields are to be planted the following spring to crops not labeled for preplant or preplant incorporated applications of **Omni Brand Pendi-Cap 3.8**.

There must be at least a 4-month interval between a **Omni Brand Pendi-Cap 3.8** fallow application and the rotational planting of any fall-seeded cereal crop. Otherwise, specific rotational crop intervals must be adhered to between a fallow application of **Omni Brand Pendi-Cap 3.8** and the planting of the following crop (see Rotational Crop Restrictions section of this label).

State-specific Instructions

In **Colorado, Idaho, Kansas, Montana, Nebraska, Nevada, Oregon, Utah, Washington, and Wyoming**, apply as a broadcast spray at rates up to, but not more than, 3.2 pts/A of **Omni Brand Pendi-Cap 3.8** herbicide.

FARMSTEAD

Apply **Omni Brand Pendi-Cap 3.8** as a broadcast spray at 2.1 qts/A for short-term (2 to 4 months) or at 4.2 qts/A for long-term (6 to 8 months) preemergence control of labeled broadleaf and grass weeds as they germinate on farmstead nonagricultural areas including barnyards, lanes, driveways, machinery or implement yards, windbreaks, and nonagricultural fencerows or ditchbanks.

FORAGE GRASSES (COOL-SEASON)

Omni Brand Pendi-Cap 3.8 may be applied by ground or chemigation, by air, or on dry bulk fertilizer.

Use Method, Rate and Timing

Established Perennial Cool-season Forage Grasses. Apply **Omni Brand Pendi-Cap 3.8** to solid stands of perennial cool-season forage grasses [including bentgrass, bluegrass (Kentucky), brome grass, fescue (fine, tall), orchardgrass, perennial ryegrass, timothy, wheatgrass, and others] grown for forage, green chop, silage, hay production, and/or grown in pastures, rangeland, or Federal Conservation Reserve Program (CRP) land for livestock grazing.

Apply **Omni Brand Pendi-Cap 3.8** only to established (grass with 6 or more tillers per plant) perennial cool-season forage grasses.

Apply **Omni Brand Pendi-Cap 3.8** before target-weed germination in fall after the last cutting/mowing/grazing, in winter, in spring, or in-season between cuttings.

Uniformly apply at a broadcast rate of 1.1 to 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre in a single application or sequential applications made 30 or more days apart.

Do not apply more than a maximum cumulative total of 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre per year.

Apply the higher rate of **Omni Brand Pendi-Cap 3.8** when higher weed pressure is anticipated or when a longer duration of residual weed control is desired.

Omni Brand Pendi-Cap 3.8 may cause temporary injury to cool-season forage grass stands. Application made in periods of cold temperatures that temporarily limit normal crop growth or in extended cold temperature periods that initiate winter dormancy in grass crops may result in crop injury. Disease, extremely cold weather, drought,

extensive frost heaving, salinity, low pH, or high pH may weaken stands and make the crop more susceptible to herbicide injury.

Mixed Stands of Established Cool-season Forage Grasses and Alfalfa. Apply **Omni Brand Pendi-Cap 3.8** to mixed stands of established cool-season forage grasses and alfalfa (established alfalfa is defined as alfalfa planted in fall or spring which has gone through a first cutting/mowing). Apply **Omni Brand Pendi-Cap 3.8** in fall after the last cutting/mowing/ grazing, in winter, in spring before weed germination, or in-season between cuttings.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Tank Mixes

Omni Brand Pendi-Cap 3.8 may be applied in a sequential use program with other herbicides labeled for use in cool-season forage grasses or applied as a tank mix with other registered herbicides that control emerged weeds. Application of postemergence herbicides may cause crop injury. Helena Agri-Enterprises, LLC advises testing **Omni Brand Pendi-Cap 3.8** tank mixes on a small portion of the target crop to determine if damage is likely to occur. Consult your local Helena Agri-Enterprises, LLC dealer for local tank mix options. Always perform a mixing test to check the compatibility of **Omni Brand Pendi-Cap 3.8** with all potential tank mix partners and fertilizers. Follow all precautions and restrictions on the labels of all products applied in combination with **Omni Brand Pendi-Cap 3.8**. Always follow the most restrictive label.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Omni Brand Pendi-Cap 3.8** with other pesticides, additives, or fertilizers.

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** if surface water is present in the field.
- Do not apply more than a maximum cumulative total of 4.2 quarts (4.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- There is no preharvest or pre-grazing interval for **Omni Brand Pendi-Cap 3.8**-treated grass forage, green chop, silage, hay, pasture, or rangeland.
- Grazing and Preharvest Interval (PHI) for forage and hay of mixed stand alfalfa/cool-season forage grasses – 14 days.
- Do not apply **Omni Brand Pendi-Cap 3.8** to mixed stands of cool-season forage grasses with other forage legumes besides alfalfa.
- Not for use in California unless otherwise directed.

FORAGE GRASSES (WARM-SEASON)

Omni Brand Pendi-Cap 3.8 may be applied by ground or chemigation, by air, or on dry bulk fertilizer.

Use Method, Rate, and Timing

Established Bermudagrass and other Perennial Warm-season Forage Grasses. Apply **Omni Brand Pendi-Cap 3.8** in fields of Bermudagrass (and other perennial warm-season grasses including Bahiagrass, buffalograss, switchgrass, and others) grown for forage or hay production, and/or grown in pastures, rangeland, or Federal Conservation Reserve Program (CRP) land for livestock grazing.

Apply **Omni Brand Pendi-Cap 3.8** herbicide only to established (defined as planted in fall or spring which has gone through a first cutting/mowing) Bermudagrass and other perennial warm-season forage grasses.

Apply **Omni Brand Pendi-Cap 3.8** before target-weed germination in the dormant season [i.e., when grasses are not actively growing in fall (postharvest), during winter dormancy period, or in early spring before greenup] or in-season between cuttings. Do not apply to Bermudagrass and other perennial warm-season grasses after greenup in spring before the first cutting.

Uniformly apply at a broadcast rate of 1.1 to 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre in a single application.

Omni Brand Pendi-Cap 3.8 may also be applied as a sequential or split program when the initial application(s) is made in the dormant season and the subsequent application(s) is made in-season between cuttings. Do not apply more than a maximum cumulative total of 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre per year.

Apply the higher rate of **Omni Brand Pendi-Cap 3.8** when higher weed pressure is anticipated or when a longer duration of residual weed control is desired.

Omni Brand Pendi-Cap 3.8 may be applied in a sequential use program or as a tank mix with other registered herbicides that control emerged weeds.

Omni Brand Pendi-Cap 3.8 may cause temporary injury to Bermudagrass and other perennial warm-season grass stands. Disease, extremely cold weather, drought, extensive frost heaving, salinity, low pH, or high pH may weaken stands and make the crop more susceptible to herbicide injury.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Tank Mixes

Omni Brand Pendi-Cap 3.8 may be tank mixed with other herbicides labeled for use in Bermudagrass and other perennial warm-season grass fields. Helena Agri-Enterprises, LLC advises testing **Omni Brand Pendi-Cap 3.8** tank mixes on a small portion of the target crop to determine if damage is likely to occur.

Application of postemergence herbicides may cause crop injury, including stunting or chlorosis of Bermudagrass and other perennial warm-season grasses. Consult your local Helena Agri-Enterprises, LLC dealer for local tank mix options. Always perform a mixing test to check the compatibility of **Omni Brand Pendi-Cap 3.8** with all potential tank mix partners and fertilizers. Follow all precautions and restrictions on the labels of all products applied in combination with **Omni Brand Pendi-Cap 3.8**. Always follow the most restrictive label.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Omni Brand Pendi-Cap 3.8** with other pesticides, additives, or fertilizers.

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** if surface water is present in the field.
- Do not apply more than a maximum cumulative total of 4.2 quarts (4.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- There is no preharvest interval for grass forage or hay, or for livestock grazing after application of **Omni Brand Pendi-Cap 3.8**.
- Not for use in California unless otherwise directed.

FRUITING VEGETABLES

Omni Brand Pendi-Cap 3.8 may be applied to the following fruiting vegetables:

African eggplant	Garden huckleberry	Okra	Tomatillo
Bell pepper	Goji berry	Pea eggplant	Tomato
Bush tomato	Groundcherry	Pepino	Tree tomato
Cocona	Martynia	Roselle	
Currant tomato	Naranjilla	Scarlet eggplant	
Eggplant	Nonbell pepper	Sunberry	

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Uniformly apply **Omni Brand Pendi-Cap 3.8** by ground or air as a broadcast preplant incorporated application or as a broadcast preplant surface application before transplanting fruiting vegetables.

Uniformly apply **Omni Brand Pendi-Cap 3.8** only by ground as a post-directed application to transplanted or established direct-seeded fruiting vegetables.

Do not apply before direct-seeded fruiting vegetables.

Do not apply postemergence over the top of or to foliage of fruiting vegetables because severe injury may occur. **Omni Brand Pendi-Cap 3.8** can be applied as a post-directed spray on the soil at the base of the plant, beneath plants, and between rows. Avoid direct contact with foliage or stems. Roots of transplants must be established. After the post-directed spray and when sufficient rainfall or irrigation does not occur to activate the herbicide,

mechanically incorporate at the time of blocking and thinning or at layby. Apply **Omni Brand Pendi-Cap 3.8** before weed germination. Emerged weeds will not be controlled by this treatment.

Omni Brand Pendi-Cap 3.8 may also be applied in fruiting vegetables transplanted to raised beds. Before transplanting, apply **Omni Brand Pendi-Cap 3.8** preplant nonincorporated in a band to the top of the pressed bed just before laying plastic. After transplanting, **Omni Brand Pendi-Cap 3.8** may also be applied in a band to the previously untreated row middles between the transplanted beds. For banded application to the bed or row middles, Do not overlap spray and apply more than the maximum broadcast use rate of **Omni Brand Pendi-Cap 3.8** on a per acre basis for the given soil texture.

Omni Brand Pendi-Cap 3.8 herbicide applied at 2.0 to 3.0 pts/A may aid in control or suppression of the following weeds when used as part of a comprehensive weed management program: black nightshade, hairy nightshade.

Use Rate

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.0 to 1.5
Medium	1.5 to 2.0
Fine	1.5 to 3.0

Restrictions

- Do not apply more than 3.0 pints (1.425 lbs of pendimethalin) **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) for tomatoes – 21 days.
- Preharvest Interval (PHI) for all other fruiting vegetables – 70 days.
- Do not plant lettuce within 6 months after a **Omni Brand Pendi-Cap 3.8** application if the rows were covered with plastic.

Precautions

- Avoid root contact with **Omni Brand Pendi-Cap 3.8**-treated soil when placing transplants into furrow or hole or injury may occur.

GRAIN SORGHUM

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Omni Brand Pendi-Cap 3.8 may be applied postemergence incorporated (CULTI-SPRAY) in grain sorghum grown in all states.

In addition, **Omni Brand Pendi-Cap 3.8** may be applied early postemergence in grain sorghum grown in states east of the Mississippi River and in Arizona, Arkansas, California, eastern Texas, Louisiana, and the Missouri bootheel.

Additional Weeds Controlled. In addition to weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** as a CULTI-SPRAY application controls the following weeds in grain sorghum: wild proso millet and shattercane.

Use Method, Rate, and Timing

CULTI-SPRAY. **Omni Brand Pendi-Cap 3.8** treatments can be applied from the 4-inch growth stage to as late as the last cultivation (layby) of grain sorghum. See specific directions for (CULTI-SPRAY) application under Application Timing.

Early Postemergence. For use only in states east of the Mississippi River and in Arizona, Arkansas, California, eastern Texas, Louisiana, and the Missouri bootheel.

Ensure the seedbed is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good seed coverage. Plant grain sorghum at least 1½ inches deep to ensure good seed coverage.

Use Rate

CULTI-SPRAY

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹ (pts/A)
Coarse	1.5	2.0
Medium	2.0	3.0
Fine	3.0	3.0

¹ See Use Precautions for map of specific states.

Early Postemergence

Soil Texture	Omni Brand Pendi-Cap 3.8 (pts/A)
Coarse	DO NOT USE
Medium, Fine	2.0

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** in grain sorghum preplant incorporated or preemergence because serious crop injury can result.
- Do not apply **Omni Brand Pendi-Cap 3.8** in grain sorghum more than once per year.
- Do not apply **Omni Brand Pendi-Cap 3.8** as a CULTI-SPRAY treatment in grain sorghum planted in double-row beds.
- Do not replant grain sorghum if crop loss occurs.
- Do not apply in liquid fertilizer.
- Livestock can graze or be fed forage from treated grain sorghum 21 days or more after application.

GREEN ONIONS

Omni Brand Pendi-Cap 3.8 may be applied to the following individual crops in the green onion crop subgroup:

Chive, fresh leaves	Fritillaria, leaves	Leek, wild	Onion, macrostem
Chive, Chinese, fresh leaves	Kurrat	Onion, Beltsville bunching	Onion, tree, tops
Elegans hosta	Lady's leek	Onion, fresh	Onion, Welsh, tops
	Leek	Onion, green	Shallot, fresh leaves

Omni Brand Pendi-Cap 3.8 may be applied preemergence, postemergence, or split application by ground, air, or chemigation.

Use Method, Rate, and Timing

Uniformly apply 2.0 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast spray to the soil surface as preemergence spray or as a postemergence spray to the crop at the 2 to 3 trueleaf stage at least 30 days before harvest. If **Omni Brand Pendi-Cap 3.8** is to be applied sequentially as both a preemergence and postemergence spray, the preemergence spray must be applied 30 days before the postemergence spray. Onion seed must be fully covered by soil at planting. Injury may occur if onion seed is exposed to **Omni Brand Pendi-Cap 3.8**.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** herbicide may be applied through sprinkler irrigation systems. Apply at 2 to 3 true-leaf stage at least 30 days before harvest. Do not irrigate more than of 0.5 inch of water. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Restrictions

- Do not apply more than 2.0 pints (1.0 lbs of pendimethalin) per acre per application.
- Do not apply more than 4.0 pints (2.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 30 days.
- Do not feed forage or graze livestock in treated fields.
- Not for use in California.

Precautions

- Only apply **Omni Brand Pendi-Cap 3.8** preemergence to green onions grown on muck soils or on mineral soils with more than 3% organic matter.

HOPS

Omni Brand Pendi-Cap 3.8 may be applied only by ground.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be used as part of a weed management program in baby (first year planting) and established hops. Apply **Omni Brand Pendi-Cap 3.8** before target-weed germination when hops are in the dormant or vegetative growth stages.

Apply **Omni Brand Pendi-Cap 3.8** as a broadcast or banded treatment (including postemergence-directed) using ground equipment. Apply the spray directly to the ground beneath the vines and/or in areas between rows. Do not apply over the top of vines with leaves or cones. Contact with hop foliage or cones by spray mixture or drift may cause injury.

Uniformly apply at a broadcast rate of 1.1 to 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre in a single application or sequential applications with an interval of 30 days or more.

Do not apply more than a maximum cumulative total of 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre per year.

Omni Brand Pendi-Cap 3.8 may be applied in a sequential use program or as a tank mix with other registered herbicides that control emerged weeds.

Tank Mixes

Omni Brand Pendi-Cap 3.8 may be tank mixed with other herbicides labeled for use in hops. Helena Agri-Enterprises, LLC advises testing **Omni Brand Pendi-Cap 3.8** tank mixes on a small portion of the target crop to determine if damage is likely to occur.

Follow all precautions and restrictions on the labels of all products applied in combination with **Omni Brand Pendi-Cap 3.8**. Always follow the most restrictive label.

Restrictions

- Do not apply to hops by air or through any type of irrigation system.
- Do not apply more than a maximum cumulative total of 4.2 quarts (4.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) for hop cones - 90 days.
- Not for use in California.

LEAF LETTUCE

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

With a single application, uniformly apply up to 2.1 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast foliar spray to direct-seeded or transplanted leaf lettuce from the 3-leaf stage until 20 days before harvest.

Restrictions

- Do not apply preplant, pretransplant, preemergence (direct-seeded) to leaf lettuce because severe injury may occur.
- Do not foliar apply to leaf lettuce before the 3-leaf growth stage because severe injury may occur.
- Do not apply more than 2.1 pints (1.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) - 20 days.

LEAFY BRASSICA GREENS

Omni Brand Pendi-Cap 3.8 may be applied in the following leafy Brassica greens: mustard greens, broccoli raab, Chinese cabbage (bok choy), collards, kale, mizuna, mustard spinach, rape greens, and turnip greens (cultivars or varieties grown for leaves only).

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

With a single application, uniformly apply up to 2.1 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast foliar spray to direct-seeded or transplanted leafy Brassica greens at the 4-leaf to 5-leaf stage.

Restrictions

- Do not apply preplant, pretransplant, or preemergence (direct-seeded) to leafy Brassica greens because severe injury may occur.
- Do not foliar apply to leafy Brassica greens before the 4-leaf growth stage because severe injury may occur.
- Do not apply to turnip greens varieties grown for roots or to dual-purpose varieties grown for roots and tops.
- Do not use **Omni Brand Pendi-Cap 3.8**-treated turnip greens roots for any feed or food purpose.
- Do not apply more than 2.1 pints (1.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 21 days.

LENTIL AND PEAS

Omni Brand Pendi-Cap 3.8 may be applied to lentil and the following peas: dry, dwarf, edible-podded, English, garden, green, and pigeon.

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied preplant surface or preplant incorporated in lentil and peas.

Preplant Surface and Preplant Incorporated (Fall Application in Idaho, Minnesota, Montana, North Dakota, Oregon, South Dakota, Washington, and Wyoming). Apply **Omni Brand Pendi-Cap 3.8** herbicide and incorporate (rainfall, irrigation, or mechanically) in late fall before planting lentils or peas the following spring. Apply **Omni Brand Pendi-Cap 3.8** in the late fall when soil temperatures are 45°F or below but before the ground freezes.

Do not apply when the air temperature is below 45°F.

Rainfall or irrigation is required for incorporation and activation. Unpredictable weed control can be expected because factors like length of time between application and planting as well as uncontrollable weather factors determine herbicide activity and longevity.

Preplant Incorporated. **Omni Brand Pendi-Cap 3.8** may be applied within 60 days of planting. After application, rotary hoeing and shallow cultivation/tillage can be practiced without reducing weed control. Avoid tillage that will bring untreated soil to the surface.

Use Rate

Preplant Surface¹, Preplant Incorporated¹

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5
Medium	2.0
Fine	3.0

¹ Follow annual restrictions and/or state-specific instructions.

State-specific Instructions

Idaho, Montana, North Dakota, Oregon, and Washington

Omni Brand Pendi-Cap 3.8 may be applied to lentil or peas (dry peas only) grown in no-tillage and/or minimum-tillage systems in Idaho, Montana, North Dakota, Oregon, and Washington. Preplant surface application must be made within 30 days of planting. When planting, ensure the seed furrow is fully closed because conditions that allow the seed furrow to inadequately close and/or allow **Omni Brand Pendi-Cap 3.8** to contact the seed may result in crop injury. Certain unfavorable environmental conditions, including cool temperatures, excessive moisture after application, and wet and/or compacted soil conditions, may result in delayed emergence and stunting with **Omni Brand Pendi-Cap 3.8** use in lentil or peas. Adequate rainfall or irrigation after application

before weed seedling germination provides the most effective weed control. Herbicide performance from surface application may be decreased compared to soil incorporated application.

Idaho, Oregon, and Washington

Omni Brand Pendi-Cap 3.8 may be applied postplant preemergence only to lentil or all peas grown in conventional-tillage systems in Idaho, Oregon, and Washington. Application must be made within 2 days of planting. Apply up to but not more than 1.5 pts/A. Apply to a firm seedbed free of clods. Soil conditions that cause poor seed furrow closure and coverage may result in delayed emergence and stunting of the crop. Certain environmental conditions including cool temperatures, excessive moisture after application, and wet soil conditions may result in delayed emergence and stunting with **Omni Brand Pendi-Cap 3.8** use in lentil or peas. Adequate rainfall or irrigation after application before weed seedling germination provides the most effective weed control.

Restrictions

- Do not use in California.
- Do not apply **Omni Brand Pendi-Cap 3.8** preemergence in peas unless otherwise noted in state-specific instructions.
- Do not apply **Omni Brand Pendi-Cap 3.8** more than once per year.
- Do not apply to lentil or peas, lentil or pea forage, pea silage, pea hay, or pea straw grown for livestock feed.
- Do not apply in any type of irrigation system.

Precautions

- Any crop registered for a preplant incorporated application of **Omni Brand Pendi-Cap 3.8** can be double cropped after peas.

MELONS

Omni Brand Pendi-Cap 3.8 may be applied in the following melons: cantaloupe, citron melon, muskmelon, and watermelon.

Omni Brand Pendi-Cap 3.8 may be applied only by ground.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied sequentially in melon production. Initially apply up to 2.1 pints per acre of **Omni Brand Pendi-Cap 3.8** as a shielded application between rows with 6 inches on either side of row middles (before melon transplanting or before a seeded crop has emerged) or between rows covered with plastic mulch (before holes are punched in plastic for melon planting). Make a second shielded application at up to 2.1 pints per acre of **Omni Brand Pendi-Cap 3.8** between rows with a minimum of 6 inches on either side of stem or vines or between plastic mulch before melon vine running. The interval between the sequential **Omni Brand Pendi-Cap 3.8** applications must be at least 21 days. Avoid spray contact with melon foliage or running vines because crop injury will occur.

Restrictions

- Do not apply more than 2.1 pints (1.0 lbs of pendimethalin) per acre in a single application or more than 4.2 pints per acre per year.
- Preharvest Interval (PHI) – 35 days.
- Do not feed forage or graze livestock in treated fields.
- Not for use in California.

MINT (PEPPERMINT AND SPEARMINT)

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Make a single broadcast preemergence application of **Omni Brand Pendi-Cap 3.8** at 1.5 pints to 4.0 pints per acre, depending on soil texture (see chart following), to dormant established mint before weed germination. After **Omni Brand Pendi-Cap 3.8** application, temporary crop injury may be observed early in the growing season as mint breaks dormancy and begins to grow.

Omni Brand Pendi-Cap 3.8 herbicide will not cause crop injury when applied according to the label under normal growing conditions. Nonuniform application may result in injury to crops, poor stands, or soil residues; conversely, uneven application may reduce weed control. Diseases, cold weather, excessive moisture, deep planting, low or high pH, salinity, or drought may weaken seedlings and plants and make them more susceptible to herbicide damage.

Use Rate

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5 to 2.0
Medium	2.0 to 4.0
Fine	2.0 to 4.0

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** to mint in the first year of growth and establishment.
- Do not apply to mint that has broken dormancy or crop injury may result. Application to mint near dormancy break can result in crop injury. Risk of crop injury increases the closer application is to mint dormancy break.
- Do not apply to mint stands that have been weakened by age, disease, cold weather, excessive moisture, or other factors that reduce crop vigor. Mint growing under stress is more susceptible to herbicide damage.
- Do not apply more than 4.0 pints (2.0 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 90 days.
- Do not allow livestock to graze on treated spent hay or feed treated spent hay to livestock.
- Do not apply this product on mint through any type of irrigation system.

MONARDA

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer.

Use Method, Rate, and Timing

Before weed germination, make a broadcast application of **Omni Brand Pendi-Cap 3.8** at 1.5 pints to 4.0 pints per acre, depending on soil texture (see chart following), to established monarda in dormancy, after emergence in the spring, or after the first harvest. After **Omni Brand Pendi-Cap 3.8** herbicide application, temporary crop injury may be observed during the growing season as monarda breaks dormancy and begins to grow, or re-grows after harvest.

Omni Brand Pendi-Cap 3.8 will not cause crop injury when applied according to the label under normal growing conditions. Nonuniform application may result in injury to crops, poor stands, or soil residues; conversely, uneven application may reduce weed control. Diseases, cold weather, excessive moisture, deep planting, low or high pH, salinity, or drought may weaken seedlings and plants and make them more susceptible to herbicide damage.

Use Rate

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5 to 2.0
Medium	2.0 to 4.0
Fine	2.0 to 4.0

Chemigation Applications. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and precautions about chemigation in the Application Instructions section of this label.

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** to monarda in the first year of growth until transplants are well established.
- Do not apply to monarda stand that have been weakened by age, disease, cold weather, excessive moisture, or other factors that reduce crop vigor, as monarda growing under stress is more susceptible to herbicide damage.
- Do not apply more than 4.0 pts (1.9 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.

- Preharvest Interval (PHI) – 90 days.
- Do not allow livestock to graze on treated spent hay or feed treated spent hay to livestock.

Precautions

- Avoid overlapping spray patterns because crop injury can occur.
- Avoid heavy residue on soil surface from previous year’s crops.

PEANUT

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Do not use in California.

Use Method, Rate, and Timing

Preplant Incorporated. Apply **Omni Brand Pendi-Cap 3.8** up to 60 days before planting and incorporate.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting and before crop emergence. For peanuts grown under overhead irrigation or to prevent decreased crop pegging, adequate incorporation must be achieved by applying a minimum of 0.75 inch of overhead irrigation or rainfall within 48 hours after **Omni Brand Pendi-Cap 3.8** application.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Use Rate

Region	Rate (pts/A)
New Mexico, Oklahoma, and Texas	1.0 to 2.0
Other peanut growing states*	2.0
* For heavy weed infestations, especially of Texas panicum, up to 3.2 pts/A of Omni Brand Pendi-Cap 3.8 herbicide can be used in Alabama, Florida, or Georgia.	

PERENNIAL GRASSES GROWN FOR SEED PRODUCTION

Omni Brand Pendi-Cap 3.8 may be applied to established (defined as planted in the fall or spring which has gone through a first cutting/mowing) warm-season perennial grasses (including Bermudagrass, switchgrass, and others) and to established (6 or more tillers per plant) cool-season perennial grasses (including Kentucky bluegrass, tall fescue, orchardgrass, perennial ryegrass, fine fescue, and others).

Omni Brand Pendi-Cap 3.8 may be applied by ground, chemigation, air, or on dry bulk fertilizer.

Use Method, Rate, and Timing

In warm-season perennial grasses, apply **Omni Brand Pendi-Cap 3.8** to postharvest grass during the fall or during winter dormancy or after the first seed harvest/cutting. Do not apply to warm-season perennial grasses after greenup in the spring before the first seed harvest/cutting. In cool-season perennial grasses, apply **Omni Brand Pendi-Cap 3.8** to postharvest grass during regrowth at the beginning of significant fall rains or in spring.

Apply **Omni Brand Pendi-Cap 3.8** before target-weed germination. Uniformly apply at a broadcast rate of 2.1 to 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre in a single application. **Omni Brand Pendi-Cap 3.8** may also be applied in two split applications, with ½ the annual application rate applied in fall or winter followed by the remaining ½ of the annual application rate applied in spring. Do not apply more than a maximum cumulative total of 4.2 quarts of **Omni Brand Pendi-Cap 3.8** per acre in any one year.

In warm-season and cool-season perennial grasses, use the high application rate of **Omni Brand Pendi-Cap 3.8** where more dense infestations of targeted annual grasses, annual broadleaf weeds, or volunteer grass seedlings are anticipated, or when a longer duration of residual weed control is desired. Excess grass straw and crop residue from the previous harvest must be evenly spread or removed by such methods as crew cutting, propane

flaming, or open field burning (when local regulations allow) before **Omni Brand Pendi-Cap 3.8** application, or reduced weed control may result.

Omni Brand Pendi-Cap 3.8 may be applied in a sequential use program or as a tank mix with other registered herbicides that control emerged weeds.

Omni Brand Pendi-Cap 3.8 may cause temporary injury to perennial grass stands. Application made in periods of cold temperatures that temporarily limit normal crop growth or in extended cold temperature periods that initiate winter dormancy in grass crops may result in crop injury. Diseases, extremely cold weather, drought, extensive frost heaving, low or high pH, or salinity may weaken stands and make them more susceptible to herbicide damage.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in Application Instructions section of this label.

Additional Weeds Controlled

Omni Brand Pendi-Cap 3.8 applied before weed germination controls annual bluegrass, volunteer fescue, and volunteer ryegrass in addition to weeds listed in **Table 2**.

Tank Mixes

Omni Brand Pendi-Cap 3.8 may be tank mixed with dimethenamid-p or with other herbicides labeled for use in perennial grasses grown for seed. Helena Agri-Enterprises, LLC advises testing **Omni Brand Pendi-Cap 3.8** tank mixes on a small portion of the target crop to determine if damage is likely to occur.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Omni Brand Pendi-Cap 3.8** with other pesticides, additives, or fertilizers.

Application of postemergence herbicides may cause crop injury. Consult your local Helena Agri-Enterprises, LLC dealer regarding local tank mix options. Always perform a mixing test to check the compatibility of **Omni Brand Pendi-Cap 3.8** with all potential tank mix partners. Follow all precautions and restrictions on the labels of all products applied in combination with **Omni Brand Pendi-Cap 3.8**. Always follow the most restrictive label.

Restrictions

- Do not apply if surface water is present in the field.
- Do not apply more than a maximum cumulative total of 4.2 quarts (4.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre in any one year.
- There is no preharvest interval for grass forage or hay, or for livestock grazing after application of **Omni Brand Pendi-Cap 3.8**.
- Preharvest Interval (PHI) for seed of warm-season and cool-season perennial grasses – 90 days.
- Not for use in California except as otherwise directed.

Precautions

- Some stunting and chlorosis of perennial grasses may occur with postemergence application.
- Application made after perennial grasses exceed 6 inches in height may result in poor weed control because of possible reduced spray coverage to the soil.
- Grass straw remaining after seed harvest of warm-season and cool-season perennial grasses may be used as livestock bedding, and/or grazed by or fed to livestock. The grower must notify the seed processor that there is no pesticide tolerance on grass seed screenings; therefore, it cannot be used in livestock feed.

POTATO

Omni Brand Pendi-Cap 3.8 herbicide may be applied by ground, air, chemigation, or on dry bulk fertilizer (preemergence incorporated only).

Additional Weeds Controlled. In addition to the weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** controls stinging nettle in potatoes.

Use Method, Rate, and Timing

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** after planting, but before potatoes and weeds emerge, or after dragoff.

Preemergence Incorporated. Apply **Omni Brand Pendi-Cap 3.8** and incorporate after planting but before potatoes and weeds emerge. Where dragoff is practiced, apply **Omni Brand Pendi-Cap 3.8** and incorporate before, at, or after dragoff, but before potatoes and weeds emerge. Take care incorporation equipment does not damage seed pieces or elongating sprouts.

Early Postemergence. Apply **Omni Brand Pendi-Cap 3.8** from crop emergence to the 6-inch growth stage. Do not apply **Omni Brand Pendi-Cap 3.8** postemergence if potatoes are under stress from cold/wet or hot/dry conditions or crop injury may occur.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Apply **Omni Brand Pendi-Cap 3.8** preemergence after planting, after dragoff, or early postemergence through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label.

Use Rate

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	1.5
Medium	2.0	3.0
Fine	3.0	3.0

Restrictions

- Do not apply to sweet potatoes or yams.
- Do not apply preplant.
- Do not make more than one application of **Omni Brand Pendi-Cap 3.8** per year.

Precautions

- Application of **Omni Brand Pendi-Cap 3.8** on White Rose variety potatoes during or followed by cool and/or wet weather conditions may result in crop injury.

RICE

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or on dry bulk fertilizer (delayed preemergence and early postemergence applications only) to rice grown under conventional-tillage, reduced-tillage, minimum-tillage, or no-till systems, or on stale seedbeds.

Additional Weeds Controlled. In addition to weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** controls the following weeds in rice: junglerice and sprangletop.

Use Method, Rate, and Timing

Delayed Preemergence. Apply **Omni Brand Pendi-Cap 3.8** alone or with tank mix partner for delayed preemergence weed control in grain-drilled, dry-seeded rice. Apply **Omni Brand Pendi-Cap 3.8** alone or in tank mix to levees after the levees are pulled and planted. The seedbed needs to be firm and free of clods and must be prepared to allow for good seed coverage. Use of a planter under conditions that do not allow good soil coverage of the rice seed can result in reduced stand or stunting if **Omni Brand Pendi-Cap 3.8** contacts germinating rice seed. Exposed seeds that come in contact with **Omni Brand Pendi-Cap 3.8** may be injured. Apply only when growing conditions favor vigorous rice growth. Ensure the seedbed has adequate moisture for seed germination. Not for use in grain-drilled, dry-seeded rice in California.

Uniformly apply the specified rate of **Omni Brand Pendi-Cap 3.8** after rice planting and before rice emergence (spiking) and weed germination. Apply after rice seed has absorbed water and germinated and after the soil has been previously sealed over the seed by at least 1 inch of rainfall or by irrigation (flush). If the soil has not been sealed by rain or flush, apply when 80 percent of germinated seeds have a primary root (radicle) or shoot at least ½-inch long. If there is insufficient moisture, Helena Agri-Enterprises, LLC advises flushing before **Omni Brand Pendi-Cap 3.8** application to supply moisture for root (radicle) initiation and for vigorous rice and weed growth.

If applied to soil before these conditions, or to cracked soil, stand reduction or stunting of rice may occur. Under some conditions, use of gibberellic acid-treated seed, heavy rainfall after application, or flushing after application may result in herbicide injury to rice. Rice can overcome moderate injury with appropriate cultural practices.

Because of the residual activity of **Omni Brand Pendi-Cap 3.8**, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of **Omni Brand Pendi-Cap 3.8**.

Early Postemergence. Apply **Omni Brand Pendi-Cap 3.8** as a tank mix partner in dry-seeded rice. Base applications on weed and crop size guidelines of the tank mix partner. Do not apply to fields with standing water. If necessary, fields may be flushed before treatment to produce vigorous rice and weed growth. Because soil and weeds must be completely exposed to spray coverage, ensure no flood water is on the field at the time of application. Cloddy soil, standing water (puddles) at the time of application, or cracks in the soil that form after application may result in reduced weed control. Because of residual activity of **Omni Brand Pendi-Cap 3.8** herbicide, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of **Omni Brand Pendi-Cap 3.8**.

Postemergence (California water-seeded rice only). As a component of a comprehensive weed management program, apply **Omni Brand Pendi-Cap 3.8** alone or tank mixed with a postemergence herbicide after water-seeded rice has reached the 4-leaf to 6-leaf stage (spike plus 3 to 5 true leaves). Applications made before the 4-leaf rice stage may result in crop injury.

Water-seeded rice must also be well-rooted/pegged (i.e. standing erect after the flood is removed) before application. Do not apply to rice that is leaning over and/or laying flat to the ground after flood removal since this is characteristic of a poorly established root system. Rice roots must be below the **Omni Brand Pendi-Cap 3.8**-treated soil zone.

Injury, stunting, and/or stand reduction can occur if **Omni Brand Pendi-Cap 3.8** contacts the rice roots.

Fields must be completely drained and free of standing water (moist/saturated soil) before application. If soil is saturated at the time of application, allow the soil surface to dry before restoring the permanent flood. **Omni Brand Pendi-Cap 3.8** requires alternate wetting/drying cycles to be activated. Weed control will be reduced if the soil surface is not allowed to dry out before restoration of the permanent flood. Resume normal water management practices after permanent flood restoration.

Omni Brand Pendi-Cap 3.8 does not control weeds postemergence; therefore, **Omni Brand Pendi-Cap 3.8** must be tank mixed with a postemergence herbicide to control emerged weeds at the time of application.

Omni Brand Pendi-Cap 3.8 aids in control or suppression of the following weeds when used as part of a comprehensive weed management program:

Barnyardgrass, early and late watergrass (including biotypes resistant to other herbicide modes of action, e.g. rice mimic), sprangletop, smallflower umbrella sedge*, redstem*

* Suppression only

In California water-seeded rice, **Omni Brand Pendi-Cap 3.8** may be applied with aerial or ground application equipment. For aerial application, apply the specified rate of **Omni Brand Pendi-Cap 3.8** in 5 gallons to 10 gallons of water per acre. If applied as a tank mix with another herbicide, make sure proper gallonage per acre per label directions (i.e. 10 to 15 with propanil) is used to ensure thorough coverage. To minimize drift, Do not apply during periods of wind more than 10 mph, or when wind conditions favor drifting, or if there is a temperature inversion. Helena Agri-Enterprises, LLC advises that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury.

For ground equipment, apply the specified rate of **Omni Brand Pendi-Cap 3.8** in 10 gallons to 20 gallons of water per acre. If **Omni Brand Pendi-Cap 3.8** is applied as a tank mix with another herbicide, make sure proper gallonage per acre per label directions (i.e. 20 to 30 for propanil) is used to ensure thorough coverage. Use a calibrated low-pressure (20 PSI to 40 PSI) sprayer equipped with appropriate nozzles for uniform spray distribution and minimize drift. Keep the bypass line on or near the bottom of the tank to minimize foaming. Nozzle screens must be no finer than 50 mesh. Do not apply **Omni Brand Pendi-Cap 3.8** during periods of gusty winds or when wind velocity is more than 20 mph.

Postemergence Tank Mixes: To control emerged weeds at application, **Omni Brand Pendi-Cap 3.8** may be tank mixed with one of the following postemergence herbicides:

bispyribac-sodium
cyhalofop

orthosulfamuron
penoxsulam

propanil
triclopyr

When using tank mixes with **Omni Brand Pendi-Cap 3.8**, always read the companion product label(s) and follow all precautions and restrictions. Always follow the most restrictive label.

Observe all restrictions regarding propanil-restricted zones.

Restrictions (for water-seeded rice)

- Do not apply **Omni Brand Pendi-Cap 3.8** before the 4-leaf rice stage (spike plus 3 true leaves) or to rice that is not well-rooted/ pegged. The rice must be standing erect after the flood is removed and before application.
- Do not apply to fields with standing water.
- Do not apply **Omni Brand Pendi-Cap 3.8** through any type of irrigation system.
- Do not apply in liquid fertilizer.
- Do not spray target crop within 60 feet of sensitive crops (crops not listed on the **Omni Brand Pendi-Cap 3.8** label).
- Do not spray target crop within 60 feet of crops labeled for **Omni Brand Pendi-Cap 3.8** application where the method of application, rate, or timing of spray application is prohibited.
- Do not apply more than the maximum rate for any soil type in one year.
- Do not use water containing **Omni Brand Pendi-Cap 3.8** residues from rice cultivation to irrigate food or feed crops not registered for use with **Omni Brand Pendi-Cap 3.8**.
- In case of a crop failure because of weather conditions or disease after treatment with **Omni Brand Pendi-Cap 3.8** alone or in tank mix, only drilled dry-seeded rice may be immediately replanted; however, the grower assumes all risks and consequences associated with replanting of rice because there is the potential for stand reduction or stunting. Helena Agri-Enterprises, LLC advises a 10% increase in seeding rate. Replant seed below the herbicide layer because reduced stand or stunting may occur if **Omni Brand Pendi-Cap 3.8** contacts germinating rice seed. Do not replant gibberellic acid-treated seed. Do not reapply **Omni Brand Pendi-Cap 3.8** alone or in tank mix.
- Do not apply to stressed rice. Stress factors include cold or hot temperature extremes, excessive moisture or drought, problem soils, poor field drainage, or deep water after application.

Use Rate

Delayed Preemergence

Soil Texture	Rate (pts/A)
Sands, loamy sands	DO NOT USE
Sandy loams	1.5
Loams, silt loams, silts, sandy clay loams	2.0
Silty clay loams, clay loams, sandy clays, silty clays, clays	2.0

Early Postemergence

Soil Texture	Rate (pts/A)
Coarse	1.5
Medium	2.0
Fine	2.0

Postemergence in California Water-seeded Rice

Soil Texture	Rate (pts/A)
Coarse	1.5
Medium	2.0
Fine	2.0

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** herbicide through any type of irrigation system.
- Do not apply in liquid fertilizer.

- Do not use on water-seeded rice unless otherwise specified.
- Do not apply to rice fields if fields are used for fish production, especially catfish farming.
- **Omni Brand Pendi-Cap 3.8** may be applied to rice fields used for crayfish production.
- Do not use water containing **Omni Brand Pendi-Cap 3.8** residue from rice cultivation to irrigate food or feed crops not registered for use with **Omni Brand Pendi-Cap 3.8**.
- In case of crop failure because of weather conditions or disease after treatment with **Omni Brand Pendi-Cap 3.8** alone or in tank mix, only drilled dry-seeded rice may be immediately replanted; however, the grower assumes all risks and consequences associated with replanting of rice because there is the potential for stand reduction or stunting. Helena Agri-Enterprises, LLC advises a 10% increase in seeding rate. Replant seed below the herbicide layer because reduced stand or stunting may occur if **Omni Brand Pendi-Cap 3.8** contacts germinating rice seed. Do not replant with gibberellic acid-treated seed. Do not reapply **Omni Brand Pendi-Cap 3.8** alone or in tank mix.
- Do not apply **Omni Brand Pendi-Cap 3.8** and then flush for germination.
- Do not apply to stressed rice. Stress factors include cold or hot temperature extremes, excessive moisture or drought, problem soils, poor field drainage, or deep water after application.
- Do not apply early preemergence or preplant incorporated because severe rice injury is possible.

ROSEMARY

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer.

Use Method, Rate, and Timing

Before weed germination, make a broadcast application of **Omni Brand Pendi-Cap 3.8** at 1.5 pints to 4.0 pints per acre, depending on soil texture (see chart following), to established rosemary prior to onset of rapid growth in early spring or after the first harvest. After **Omni Brand Pendi-Cap 3.8** herbicide application, temporary crop injury may be observed during the growing season as rosemary grows or re-grows after harvest.

Omni Brand Pendi-Cap 3.8 will not cause crop injury when applied according to the label under normal growing conditions. Nonuniform application may result in injury to crops, poor stands, or soil residues; conversely, uneven application may reduce weed control. Diseases, cold weather, excessive moisture, deep planting, low or high pH, salinity, or drought may weaken seedlings and plants and make them more susceptible to herbicide damage.

Use Rate

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5 to 2.0
Medium	2.0 to 4.0
Fine	2.0 to 4.0

Chemigation Applications. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and precautions about chemigation in the Application Instructions section of this label.

Restrictions

- Do not apply **Omni Brand Pendi-Cap 3.8** to rosemary in the first year of growth and establishment.
- Do not apply to rosemary stand that have been weakened by age, disease, cold weather, excessive moisture, or other factors that reduce crop vigor, as rosemary growing under stress is more susceptible to herbicide damage.
- Do not apply more than 4.0 pts (1.9 lbs pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) – 60 days.
- Do not allow livestock to graze on treated spent hay or feed treated spent hay to livestock.

Precautions

- Avoid overlapping spray patterns because crop injury can occur.
- Avoid heavy residue on soil surface from previous year's crops.

SAFFLOWER

Omni Brand Pendi-Cap 3.8 herbicide may be applied to safflower by ground, air, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Plant safflower 1½ inches to 2-inches deep and completely cover with soil. In California, plant safflower deep enough to completely cover with soil.

Use Method, Rate, and Timing

Preplant Incorporated. In all states, apply within 60 days of planting and incorporate.

Preplant Incorporated

Fall Application in Minnesota, Montana, North Dakota, and South Dakota. Apply **Omni Brand Pendi-Cap 3.8** and immediately incorporate in late fall before planting safflower the following spring. Apply **Omni Brand Pendi-Cap 3.8** in the late fall when soil temperatures are 45°F or below but before the ground freezes. Do not apply when air temperature is below 45° F. Before safflower planting in the spring, fields treated with **Omni Brand Pendi-Cap 3.8** must receive at least one shallow additional incorporation. Spring incorporation must be at an angle to the last tillage operation.

Fall Application in California. Apply **Omni Brand Pendi-Cap 3.8** and immediately incorporate during tillage operations in the fall to target winter annual weeds before planting safflower the following spring. Before safflower planting in the spring, fields treated with **Omni Brand Pendi-Cap 3.8** must receive at least one additional incorporation.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting.

Preemergence application of **Omni Brand Pendi-Cap 3.8** to safflower may increase the likelihood of crop injury, especially when crops are grown in stress situations, including compacted soils. Decreased herbicide performance compared to preplant incorporated application may also result from a preemergence application. If dry conditions with limited precipitation exist or unseasonably cool temperatures after planting are forecast, apply **Omni Brand Pendi-Cap 3.8** before planting and mechanically incorporate with tillage. **Omni Brand Pendi-Cap 3.8** may be applied preemergence in conventional-tillage safflower.

Preemergence application of **Omni Brand Pendi-Cap 3.8** to safflower grown in California must be followed with irrigation or rainfall to establish a crop stand.

No-till Safflower. **Omni Brand Pendi-Cap 3.8** may be applied at 3.0 pts/A up to 30 days before planting (preplant) to immediately after planting (preemergence). Do not use in California.

Use Rate

Preplant Incorporated, Preemergence

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹	
		< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	2.0	2.0
Medium	2.0	2.5	3.0
Fine	3.0	3.0	3.0

¹ See Use Precautions for map of specific states.

Preplant Incorporated (Fall)¹

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	2.5	2.5
Medium	3.0	3.5
Fine	3.5	3.5

¹ For use in California, Minnesota, Montana, North Dakota, and South Dakota only.

Restrictions (All Tillage Types)

- Do not apply **Omni Brand Pendi-Cap 3.8** postemergence.
- Do not apply more than the highest rate per acre for any soil type.
- Do not feed forage or graze livestock in treated safflower fields.

SOYBEAN

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Omni Brand Pendi-Cap 3.8 may be applied to soybean grown under conventional-tillage, minimum-tillage, or no-till systems.

Additional Weeds Controlled. In addition to weeds listed in **Table 2**, **Omni Brand Pendi-Cap 3.8** controls or suppresses the following weeds in soybean: red rice and itchgrass. For specific rates for itchgrass and red rice management, see table at end of this section.

Use Method, Rate, and Timing

Fall Applied. **Omni Brand Pendi-Cap 3.8** may be surface applied or incorporated in the fall, after fall harvest and before ground freeze in states north of I-80 and the entire states of Illinois, Indiana, Iowa, Kansas, Kentucky, Missouri, Nebraska, Ohio, Oklahoma, and Texas. Fall application of **Omni Brand Pendi-Cap 3.8** will not provide season-long weed control.

Preplant Surface. Apply **Omni Brand Pendi-Cap 3.8** within 15 days of planting. **Omni Brand Pendi-Cap 3.8** may be applied within 45 days of planting when used in tank mix or applied sequentially with imazethapyr- or imazamox-containing herbicides labeled with the registered crop use.

Preplant Incorporated. Apply **Omni Brand Pendi-Cap 3.8** within 60 days of planting and incorporate.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** herbicide at planting or within 2 days after planting. Apply to a firm seedbed free of clods. Do not apply **Omni Brand Pendi-Cap 3.8** preemergence north of Interstate 80, except in the states of Indiana, Michigan, and Ohio.

Use Rate

Fall Surface, Fall Incorporated, Preplant Surface, Preplant Incorporated

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	2.0
Medium	2.5*	3.0
Fine**	3.0	3.0

* Do not apply more than 2.1 pts for Southern states; see Use Precautions for map of specific states.
 ** For heavy clay soils, apply **Omni Brand Pendi-Cap 3.8** at the broadcast rate of 3.2 pints per acre.

Preemergence

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	1.5
Medium	2.0	2.0
Fine	2.0	2.5

Preplant Incorporated

Red Rice Control and Itchgrass Suppression

Soil Texture	Up to 3% Organic Matter ¹ (pts/A)
Coarse	3.0
Medium	3.0
Fine	4.0

¹ Do not use on soils with more than 3% organic matter.

Restrictions

- Do not use **Omni Brand Pendi-Cap 3.8** in soybean in California.
- Livestock can graze or be fed forage from treated soybean fields.
- Preharvest Interval (PHI) – 85 days.

- Do not apply more than one application per year at the highest rate per acre for any given soil type and application method.

STRAWBERRY (AND OTHER LOW-GROWING BERRIES)

Omni Brand Pendi-Cap 3.8 may be applied in the following low-growing berries:

Bearberry	Cloudberry	Muntries
Bilberry	Cranberry	Partridgeberry
Blueberry, lowbush	Lingonberry	Strawberry

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or chemigation.

Use Method, Rate, and Timing

Stunting, reduced growth, or reduction in daughter plants may occur with use of **Omni Brand Pendi-Cap 3.8** in strawberries and other low-growing berries.

Uniformly apply 1.5 to 3.0 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast spray to the soil surface at pretransplant time or post-transplant time (must be within 7 days of transplanting of rootstock in the Pacific Northwest). However, in areas where irrigation is used daily (frequently) after transplanting, apply **Omni Brand Pendi-Cap 3.8** just before the end of the watering regime to maximize weed control benefits of **Omni Brand Pendi-Cap 3.8**. Extended periods of irrigation may reduce residual control provided by **Omni Brand Pendi-Cap 3.8**.

However, **Omni Brand Pendi-Cap 3.8** applications to row middles between the beds are allowed. Do not apply posttransplant if new foliage from rootstock is exposed to spray area. A second application of 1.5 to 3.0 pints per acre of **Omni Brand Pendi-Cap 3.8** may be applied in a band to the soil between crop rows (or between the plastic beds) 35 days before harvest, but Do not CONCENTRATE THE RATE per acre into the treated area, and Do not allow spray to contact strawberry or other low-growing berry plants. The second application rate is based on per unit of treated area.

Omni Brand Pendi-Cap 3.8 may also be applied to strawberries or other low-growing berries in fall or winter dormancy. Uniformly apply 1.5 to 3.0 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast spray to the soil surface before onset of new seasonal growth from strawberry or other low-growing berry crowns. Do not apply if new seasonal growth (leaves) has emerged or is exposed.

Omni Brand Pendi-Cap 3.8 may also be applied to perennial strawberries or other low-growing berries after renovation. Uniformly apply 1.5 to 3.0 pints per acre of **Omni Brand Pendi-Cap 3.8** as a broadcast spray to the soil surface after renovation (mowing or other defoliation operation) when no foliage is exposed but before onset of new seasonal growth from strawberry or other low-growing berry crowns. Do not apply if new seasonal growth (leaves) has emerged or leaves are exposed.

Chemigation Application. **Omni Brand Pendi-Cap 3.8** may be applied through sprinkler irrigation systems. Follow all directions, special instructions, and restrictions about chemigation in the Application Instructions section of this label. Do not allow **Omni Brand Pendi-Cap 3.8**-treated irrigation water to contact strawberry or other low-growing berry plants.

Use Rate

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5
Medium	2.0 to 2.5
Fine	2.5 to 3.0

Restrictions

- Do not apply more than 3.0 pints (1.425 lbs of pendimethalin) per acre per application.
- Do not apply more than 6.0 pints (2.85 lbs of pendimethalin) per acre per year.
- Preharvest Interval (PHI) – 35 days.
- Do not feed forage or graze livestock in treated fields.
- Do not plant lettuce within 6 months after a **Omni Brand Pendi-Cap 3.8** herbicide application if strawberry beds were covered with plastic.

Additional Use in Oregon and Washington in First Year Nonbearing Strawberries

Uniformly broadcast apply **Omni Brand Pendi-Cap 3.8** preemergence before transplanting strawberries. Do not harvest for food or feed any portion of the strawberry plant within 1 year (365 days) of **Omni Brand Pendi-Cap 3.8** application. DO NOT apply **Omni Brand Pendi-Cap 3.8** through any type of irrigation system or by air.

Broadcast Use Rate

First Year Nonbearing Strawberries

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5 to 2.0	2.0 to 3.0
Medium	2.0 to 2.5	2.0 to 3.0
Fine	2.0 to 3.0	2.5 to 3.5

SUGARCANE

Omni Brand Pendi-Cap 3.8 may be applied by ground or air.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied preemergence through layby to plant or ratoon sugarcane. Although there may be adequate crop tolerance for postemergence application at layby, the spray must be directed under the sugarcane canopy to obtain effective weed control.

Use Rate

Use Area	Broadcast Rate ¹ (pts/A)
All states, except Hawaii	4.2 to 6.2
Muck soils (Florida only)	4.2 to 8.4
Hawaii	4.2 to 8.4

¹ Use the high rate if: heavy clay soils; no mechanical incorporation is planned; heavy weed populations are anticipated; itchgrass infestation is anticipated; no shaving is planned.

Additional Use as Fallow Ground Application Only in Louisiana. Apply **Omni Brand Pendi-Cap 3.8** before weed germination for control of annual grasses including itchgrass (Raoulgrass), seedling Johnsongrass, and Panicum spp. in preplant fallow ground sugarcane. If necessary, control emerged weeds before application of **Omni Brand Pendi-Cap 3.8** with postemergence herbicides and/or mechanical cultivation.

After cultivation and forming the beds in the spring, apply **Omni Brand Pendi-Cap 3.8** at 2.6 qts/A using ground equipment. Ensure sugarcane beds are free of trash or clods at the time of application. If sufficient rainfall ($\frac{1}{2}$ to $\frac{3}{4}$ inch) has not occurred within 7 days of application, perform a shallow incorporation (1 to 2 inches) with an additional pass of a Lilliston-type Lely Roterra™ cultivator set to cut 2-inches or 3-inches deep. A minimum interval of 60 days between **Omni Brand Pendi-Cap 3.8** application and planting of sugarcane is required or crop injury may occur. After planting, apply **Omni Brand Pendi-Cap 3.8** to sugarcane preemergence through layby, but Do not apply more than 12.5 pts/A of **Omni Brand Pendi-Cap 3.8** during one year.

Noncropped Water Drainage Areas Application Only in Louisiana. Apply **Omni Brand Pendi-Cap 3.8** before weed germination to nonirrigated, noncropped water drainage areas (ditchbanks) adjacent to sugarcane fields. If necessary, control emerged weeds before application of **Omni Brand Pendi-Cap 3.8** with postemergence herbicides and/or mechanical cultivation.

Apply **Omni Brand Pendi-Cap 3.8** at 2.6 to 3.5 qts/A using ground equipment. Do not apply **Omni Brand Pendi-Cap 3.8** below the high water mark or when water is present in the drainage area (ditchbank). Do not apply more than 12.5 pts/A of **Omni Brand Pendi-Cap 3.8** during one year.

Areas in Hawaii Subject to High Winds. For wind speeds between 10 to 20 mph, Do not apply in a manner that allows spray to drift from the application target site. Use drift-mitigating measures, including lowering the spray boom; use coarse spray according to ASABE 572 definition for standard nozzles; use hooded or shielded sprayers; use spray drift retardants; or use any other measures known to control drift.

Restrictions

- Do not apply more than 12.5 pints (6.0 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre in one year.
- Do not use less than 11 gallons of water as a carrier when applying **Omni Brand Pendi-Cap 3.8** for weed control.
- Do not make aerial application at close-in because complete and uniform coverage cannot be obtained.
- Do not apply through any type of irrigation system.
- Preharvest Interval (PHI) – 90 days.
- Do not graze treated fields or feed treated forage or fodder to livestock.

Precautions

- Ratoon sugarcane must be lightly shaved in early spring to remove the old stubble before incorporation over the line of sugarcane is possible. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots.

SUNFLOWER (AND OTHER GROUP 20B OILSEEDS)

Omni Brand Pendi-Cap 3.8 herbicide may be applied to sunflower and other Group 20B oilseed crops as follows:

Calendula	Evening primrose	Stokes aster	Vernonia
Castor oil plant	Jojoba	Sunflower	
Chinese tallowtree	Niger seed	Tallowwood	
Euphorbia	Rose hip	Tea oil plant	

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, or on dry bulk fertilizer (only fall and preplant incorporated applications).

Plant sunflower and other Group 20B oilseeds 1½-inches to 2-inches deep and completely cover with soil.

Use Method, Rate, and Timing

Preplant Incorporated (Spring). In all states, apply within 60 days of planting and incorporate.

Preplant Incorporated (Fall Applications Only in Minnesota, North Dakota, and South Dakota). Apply **Omni Brand Pendi-Cap 3.8** and immediately incorporate in late fall before planting sunflower or oilseeds the following spring. Apply **Omni Brand Pendi-Cap 3.8** in late fall when soil temperatures are 45°F or below but before the ground freezes. Do not apply when air temperature is below 45° F. Before sunflower or other Group 20B oilseeds planting in the spring, fields treated with **Omni Brand Pendi-Cap 3.8** must receive at least one shallow additional incorporation. Spring incorporation must be at an angle to the last tillage operation.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting. Preemergence application of **Omni Brand Pendi-Cap 3.8** to sunflower or other Group 20B oilseeds may increase the likelihood of crop injury, especially when crops are grown in stress situations, including compacted soils. Decreased herbicide performance compared to preplant incorporated application may also result from a preemergence application. If dry conditions with limited precipitation exist or unseasonably cool temperatures after planting are forecast, apply **Omni Brand Pendi-Cap 3.8** before planting and mechanically incorporate with tillage. **Omni Brand Pendi-Cap 3.8** may be applied preemergence in conventional tillage sunflower or other Group 20B oilseeds, except in the state of California.

No-till Sunflower or other Group 20B Oilseeds. **Omni Brand Pendi-Cap 3.8** may be applied at 3.0 pts/A up to 30 days before planting (preplant) to immediately after planting (preemergence). Do not use in California.

Use Rate

Preplant Incorporated (Spring), Preemergence (Conventional Tillage)

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹	
		< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	2.0	2.0
Medium	2.0	2.5	3.0
Fine	3.0	3.0	3.0

¹ See Use Precautions for map of specific states.

Preplant Incorporated (Fall)¹

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	2.5	2.5
Medium	3.0	3.5
Fine	3.5	3.5

¹ For use in Minnesota, North Dakota, and South Dakota only.

Restrictions (All Tillage Types)

- Do not apply **Omni Brand Pendi-Cap 3.8** postemergence.
- Do not feed forage or graze livestock in treated sunflower or other Group 20B oilseeds fields.

TOBACCO

Omni Brand Pendi-Cap 3.8 may be applied by ground only.

Use Method, Rate, and Timing

Preplant Incorporated. Apply and incorporate within 60 days of transplanting tobacco.

Applied according to directions and under normal growing conditions, **Omni Brand Pendi-Cap 3.8** will not harm transplanted tobacco. Under stress conditions for plant growth, including cold/wet or hot/dry weather, **Omni Brand Pendi-Cap 3.8** can produce a temporary retardation of tobacco development.

Layby. **Omni Brand Pendi-Cap 3.8** may be applied as a directed spray after the last normal cultivation (layby), usually 4 to 6 weeks after transplanting tobacco. Apply **Omni Brand Pendi-Cap 3.8** in a 16-inch to 24-inch band between the crop rows. Do not contact tobacco plants with spray.

Use Rate

Preplant Incorporated

Use Area	Soil Texture	Rate (pts/A)
Florida, Georgia, Maryland, North Carolina, South Carolina, Virginia	Coarse	2.0
	Medium	
	sandy clay loams, loams	2.0
	silt loams, silts	2.5
	Fine	2.5
Other states	Coarse	2.0
	Medium	3.0
	Fine	3.0

Layby

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.5
Medium	2.0
Fine	2.0

Precautions

- Do not apply as a broadcast spray or contact may cause malformed tobacco leaves.

VEGETABLE SOYBEAN (EDAMAME)

Omni Brand Pendi-Cap 3.8 herbicide may only be applied by ground.

Use Method, Rate, and Timing

Omni Brand Pendi-Cap 3.8 may be applied to edamame grown under conventional-tillage, minimum-tillage, or no-till systems.

Preplant Surface. Apply **Omni Brand Pendi-Cap 3.8** within 15 days of planting. **Omni Brand Pendi-Cap 3.8** may be applied within 45 days of planting when used in tank mix or applied sequentially with postemergence-applied herbicides registered for use in edamame.

Preplant Incorporated. Apply **Omni Brand Pendi-Cap 3.8** within 60 days of planting and incorporate.

Preemergence. Apply **Omni Brand Pendi-Cap 3.8** at planting or up to 2 days after planting. Apply to a firm seedbed, free of clods. Do not apply **Omni Brand Pendi-Cap 3.8** preemergence north of Interstate 80, except in the states of Indiana, Michigan, and Ohio.

Use Rate

Preplant Surface, Preplant Incorporated

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	2.0
Medium	2.5*	3.0
Fine**	3.0	3.0

* Do not apply more than 2.1 pts for Southern states; see Use Precautions for map of specific states.
 ** For heavy clay soils, apply **Omni Brand Pendi-Cap 3.8** at the broadcast rate of 3.2 pints per acre.

Preemergence

Soil Texture	< 3% Organic Matter (pts/A)	> 3% Organic Matter (pts/A)
Coarse	1.5	1.5
Medium	2.0	2.0
Fine	2.0	2.5

Restrictions

- Livestock can graze or be fed forage from treated vegetable soybean (edamame) fields.
- Preharvest Interval (PHI) – 85 days.
- Do not apply more than one application per year at the highest rate per acre for any given soil type and application method.
- Do not use in California.

WHEAT AND TRITICALE

Omni Brand Pendi-Cap 3.8 may be applied by ground, air, chemigation, or on dry bulk fertilizer (only delayed preemergence and postemergence applications).

Omni Brand Pendi-Cap 3.8 may be applied preemergence, delayed preemergence, or postemergence for weed control in fall-seeded, winter-seeded, or spring-seeded wheat or triticale.

Use Method, Rate, and Timing

Apply to a seedbed which is firm and free of clods and trash. The seedbed **MUST** be prepared to ensure thorough seed coverage by the soil and seed-to-soil contact. Use high quality seed. When application of **Omni Brand Pendi-Cap 3.8** is intended to be made preemergence or delayed preemergence, plant seed at least 1-inch deep to avoid possible crop injury, but not too deep for proper germination. When application of **Omni Brand Pendi-Cap 3.8** is intended to be made postemergence, plant seed at least ½-inch to 1-inch deep to avoid crop injury.

Uniformly apply **Omni Brand Pendi-Cap 3.8** preemergence, delayed preemergence (after wheat or triticale seed has germinated), or postemergence from the 1st-leaf stage of wheat or triticale until before the flag leaf is visible/emerged for weed control. Apply **Omni Brand Pendi-Cap 3.8** before weed germination. Emerged weeds will not be controlled by this treatment.

For control of established weeds, **Omni Brand Pendi-Cap 3.8** herbicide may be tank mixed with any postemergence herbicide registered for use in wheat or triticale. **Omni Brand Pendi-Cap 3.8** provides residual control of weeds listed in this label. Always perform a mixing test to check the compatibility of **Omni Brand Pendi-Cap 3.8** with all potential tank mix partners.

Use Rate

Soil Texture	Southern States ¹ (pts/A)	Northern States ¹ (pts/A)
Coarse	1.5 to 2.0	1.5
Medium	1.5 to 3.0	1.5 to 2.5
Fine	2.0 to 3.0	2.0 to 3.0

¹ See Use Area map in Use Precautions.

In wheat stubble, **Omni Brand Pendi-Cap 3.8** may be applied in the fall, spring, or early summer during the fallow period after wheat harvest as a planned residual treatment to control labeled broadleaf and grass weeds. **Omni Brand Pendi-Cap 3.8** must be applied with a tank mix partner (i.e. glyphosate) for control of emerged weeds. There must be at least a 4-month interval between a **Omni Brand Pendi-Cap 3.8** fallow application and the rotational planting of any fall-seeded cereal crop. Apply up to, but Do not apply more than, 3 pints/A of **Omni Brand Pendi-Cap 3.8** in any fallow application. Do not make more than one application of **Omni Brand Pendi-Cap 3.8** during a single fallow period before rotational planting of any fall-seeded cereal crops. Follow rotational crop restrictions when planting a rotational crop after a fallow application of **Omni Brand Pendi-Cap 3.8**.

Restrictions

- Do not apply more than 3.0 pints (1.425 lbs of pendimethalin) of **Omni Brand Pendi-Cap 3.8** per acre per year.
- Preharvest Interval (PHI) for grain and straw – 60 days.
- Preharvest Interval (PHI) for hay – 28 days.
- Preharvest Interval (PHI) for forage – 11 days.

Precautions

- **NOTE:** If loss of grain crop occurs, any crop registered for **Omni Brand Pendi-Cap 3.8** preplant incorporated use may be replanted the same year without adverse effects. Do not replant wheat or triticale.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Omni Brand Pendi-Cap 3.8 freezes around 15°F and is stable under conditions of freezing and thawing. Product that has been frozen must be thawed and recirculated prior to use.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. Open dumping is prohibited. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

[Non-Refillable Plastic Container ≤5 gal.] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state or local authorities.

[Non-Refillable Plastic Container >5 gal.] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple Rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold

container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state or local authorities.

[Non-Refillable Plastic Containers [(e.g., Intermediate Bulk Containers [IBC]) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down)]] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state or local authorities.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and must be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Agri-Enterprises, LLC (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. To the extent consistent with applicable law, The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Agri-Enterprises, LLC's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

Uses with Other Products (Tank Mixes)

To the extent consistent with applicable law, if this product is used in combination with any other product except as specifically advised in writing by Helena Agri-Enterprises, LLC, then Helena Agri-Enterprises, LLC shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically advised. To the extent consistent with applicable law, if used in combination advised by Helena Agri-Enterprises, LLC, the liability of Helena Agri-Enterprises, LLC shall in no manner extend to any damage, loss, or injury not directly caused by the inclusion of the Helena Agri-Enterprises, LLC product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

{Referenced Herbicides (tradename, EPA Reg. No., ai):}

{Beyond[®] Xtra herbicide, 241-441, imazamox}

{Outlook[®] herbicide, 7969-156, dimethenamid}

{Pursuit[®] herbicide, 241-310, imazethapyr}

[Beyond, Outlook, and Pursuit are registered trademarks of BASF.]

[Roterra is a trademark of Lely USA, Inc.]

[© 20[XX] Helena Agri-Enterprises, LLC]

[EPA APPROVAL DATE]

{BASE LABEL}

PENDIMETHALIN GROUP 3 HERBICIDE

Omni Brand Pendi-Cap 3.8

Herbicide [For Use in Selected Crops] [(See Table 1. Crop Uses)]

ACTIVE INGREDIENT:	% BY WT
pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine*	38.7%
OTHER INGREDIENTS:	61.3%
TOTAL:	100.0%

*1 gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed or absorbed through skin. Cause moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwater or rinsate.

NON-TARGET ORGANISM ADVISORY STATEMENT

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Endangered Species Protection

This product may have effects on federally listed threatened or endangered plant species or their critical habitat. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the county or parish in which you are applying the pesticide. To determine whether your county or parish has a Bulletin, and to obtain that Bulletin, consult <http://www.epa.gov/espp/>, or call 1-844-447-3813 no more than 6 months before using this product. Applicators must use Bulletins that are in effect in the month in which the pesticide will be applied. New Bulletins will be available from the above sources 6 months before their effective dates.

If endangered plant species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave an untreated buffer zone of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASABE fine to medium/ coarse nozzles.
- If applied by air, leave an untreated buffer zone of 170 feet. Must use straight-stream nozzles (D-6 or larger); wind can be no more than 8 mph, and release height must be 15 feet or less.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Omni Brand Pendi-Cap 3.8 freezes around 15°F and is stable under conditions of freezing and thawing. Product that has been frozen must be thawed and recirculated prior to use.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. Open dumping is prohibited. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

[Non-Refillable Plastic Container ≤5 gal.] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state or local authorities.

[Non-Refillable Plastic Container >5 gal.] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple Rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state or local authorities.

[Non-Refillable Plastic Containers [(e.g., Intermediate Bulk Containers [IBC]) (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down)]] Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into formulation equipment and before final disposal using the following pressure rinsing procedure: Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure spray duration and/or spray volume. If the manufacturer's instructions are not available pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state or local authorities.

[See] [inside] [label] [booklet] [side] [panel] [for] [First Aid][.] [additional] [Precautionary Statements][.] [and] [Directions for Use] [including] [Storage and Disposal] [instructions][.]

EPA Reg. No.: 5905-[XX]

[AD XXXXX][internal tracking]

EPA Est. No.:

[Lot][Batch][No.]:]

Manufactured for:

NET CONTENTS:

Helena Agri-Enterprises, LLC

225 Schilling Boulevard, Suite 300 Collierville, TN 38017