

**FORMULA 202** NET WT. 4 LBS.

# GARDEN DUST

Contains...

**5%**  
**SEVIN\***

For control of insect pests

EPA Reg. No. 5927-7

Active Ingredient:

1-Naphthyl Methylcarbamate ..... 5.00%

Inert Ingredients ..... 95.00%

Trademark Union Carbide Corp.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See other panel for additional cautions

**FORM**

**GARDI**

Co

**5% S**

DIR

**BEANS:** To control Mexican Be per 50 feet of row, 1½ ounce and 2 ounces per 50 feet of r

**POTATOES:** To control Colora ounces per 50 feet of row, 1 Beetle, 2 ounces for Lygus B

**TOMATOES:** To control Tomat Flea Beetles, apply at rate of 2 Tomato Hornworm and Stink 50 feet of row.

**CUCUMBERS AND SUMMER** Spotted Cucumber Beetles, Worm and Melon Worm, apply Repeat at 7 to 10 day interv apply to Egg Plant, Peppers c In dusting, be certain that the and lower sides. Do not use w very warm weather use in th down.

**ANIMALS: DOGS, CATS:** To co liberally to animal and rub tho feet, between the toes and on quarters and other areas wh weekly intervals if needed. Do 4 weeks old.

**CAUTION:** Harmful if swallow

# SEVIN\*

For control of insect pests

EPA Reg. No. 5927-7

Active Ingredient:

1-Naphthyl Methylcarbamate ..... 5.00%

Inert Ingredients ..... 95.00%

Trademark Union Carbide Corp.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

See other panel for additional cautions

**NET WT. 4 LBS.**

Manufactured by

**DACUS, INC.**

TUPELO, MISS. 38801

NET WT. 4 LBS.

**GARDEN DUST**

**FORMULA-202**

**5% SEVIN\***

ounces per 50 feet of row  
Beetle, 2 ounces for Lyg

**TOMATOES:** To control  
Flea Beetles, apply at rate  
Tomato Hornworm and S  
50 feet of row.

**CUCUMBERS AND SUM**  
Spotted Cucumber Beet  
Worm and Melon Worm, a  
Repeat at 7 to 10 day in  
apply to Egg Plant, Pepp  
In dusting, be certain tha  
and lower sides. Do not u  
very warm weather use  
down.

**ANIMALS: DOGS, CATS:**  
liberally to animal and rub  
feet, between the toes and  
quarters and other areas  
weekly intervals if needed  
4 weeks old.

**CAUTION:** Harmful if swa  
avoid breathing dust. Avoi  
handling. Change contami  
of Feed and Food Stuffs.

Do not feed plants treat

Do not reuse empty conta  
burning. Stay away from

ACCEPTED

**FORMULA-202**

# GARDEN DUST

Contains...

**5% SEVIN\***

## DIRECTIONS

**BEANS:** To control Mexican Bean Beetles apply at rate of  $\frac{3}{4}$  ounces per 50 feet of row,  $1\frac{1}{2}$  ounces per 50 feet of row for Leafhopper and 2 ounces per 50 feet of row for Lygus and Stink Bugs.

**POTATOES:** To control Colorado Potato Beetle apply at rate of  $\frac{3}{4}$  ounces per 50 feet of row,  $1\frac{1}{2}$  ounces for Leafhopper and Flea Beetle, 2 ounces for Lygus Bug.

**TOMATOES:** To control Tomato Fruit Worm, Fall Army Worm and Flea Beetles, apply at rate of 2 ounces per 50 feet of row. To control Tomato Hornworm and Stink Bugs, apply at rate of 4 ounces per 50 feet of row.

**CUCUMBERS AND SUMMER SQUASH:** To control Striped and Spotted Cucumber Beetles, Squash Bugs, Flea Beetles, Pickle Worm and Melon Worm, apply at rate of 1 ounce per 50 feet of row. Repeat at 7 to 10 day intervals as necessary for control. Do not apply to Egg Plant, Peppers or Tomatoes within 1 day of harvest. In dusting, be certain that the foliage is covered on both the upper and lower sides. Do not use when temperature is 85° F. or over. In very warm weather use in the morning when the temperature is down.

**ANIMALS: DOGS, CATS:** To control Brown Dog Ticks and Fleas apply liberally to animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Repeat at weekly intervals if needed. Do not treat kittens or puppies less than 4 weeks old.

NET WT.  
4 LBS.

DUST



—Chase Bag Co—5852-325—

5.007  
95.007%



ounces per 50 feet of row, 1½ ounces for Leafhopper and Flea Beetle, 2 ounces for Lygus Bug.

**TOMATOES:** To control Tomato Fruit Worm, Fall Army Worm and Flea Beetles, apply at rate of 2 ounces per 50 feet of row. To control Tomato Hornworm and Stink Bugs, apply at rate of 4 ounces per 50 feet of row.

**CUCUMBERS AND SUMMER SQUASH:** To control Striped and Spotted Cucumber Beetles, Squash Bugs, Flea Beetles, Pickle Worm and Melon Worm, apply at rate of 1 ounce per 50 feet of row. Repeat at 7 to 10 day intervals as necessary for control. Do not apply to Egg Plant, Peppers or Tomatoes within 1 day of harvest. In dusting, be certain that the foliage is covered on both the upper and lower sides. Do not use when temperature is 85° F. or over. In very warm weather use in the morning when the temperature is down.

**ANIMALS: DOGS, CATS:** To control Brown Dog Ticks and Fleas apply liberally to animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Repeat at weekly intervals if needed. Do not treat kittens or puppies less than 4 weeks old.

**CAUTION:** Harmful if swallowed or inhaled. Do not take internally, avoid breathing dust. Avoid contact with skin wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of Feed and Food Stuffs.

Do not feed plants treated with Sevin to meat or dairy animals.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

—Chase Bag Co—5852-325—



95.55%

DREN

ACCEPTED

1957-7