

FORMULA 202

**NET WT.
4 LBS.**

GARDEN DUST

Contains...

**10%
SEVIN***

ACCEPTED
MAY 15 1975
Under the Federal Insecticide, Fungicide, and Herbicide Act, as amended, for the pesticide registered under EPA Reg. No.

For control of insect pests

EPA Reg. No. 5927-12

Active Ingredient:

Carbaryl (1-Naphthyl Methylcarbamate) . . . 10% by wt.
Inert Ingredients . . . 90% by wt.

Active Ingredient:
Carbaryl (1-Naphthyl Methylcarbamate) . . 10% by wt.
Inert Ingredients 90% by wt.
*Trademark of Union Carbide Chemical Company

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other panel for additional cautions

NET WT. 4 LBS.

Manufactured by
DACUS, INC.
TUPELO, MISS. 38801

EPA EST. 5927-MS-1

FORMULA-202

GARDEN DUST

Contains...

10% SEVIN*

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of food, feed, and water supplies. Dispose of container when empty; never reuse. Destroy by burning. Stay away from smoke or fumes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

VEGETABLES: Use the amount of dust necessary to provide adequate coverage. Begin applications when insects or damage appear and continue 7 to 10 day intervals or as necessary for good control.

BEANS: To control Bean Leaf Beetle, Cucumber beetle, Western bean cutworms, Flea Beetle and Japanese beetle, Corn earworm, Armyworms and Velvet bean caterpillar, Lima bean pod borer, Lygus bug and Stink bug, apply the dust to thoroughly cover upper and lower leaf surfaces.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHL-RABI: To control — Flea beetle, Harlequin bugs, Corn earworm, armyworm, imported cabbage worm and small cabbage loopers, apply the dust to thoroughly cover upper and lower surfaces. **CAUTION:** Allow 3 days between last application and harvest.

CORN: To control — Corn earworm, European corn borer, Fall armyworm, Sap beetles and Japanese beetle, apply the dust to thoroughly cover upper and lower leaf surfaces. For corn earworm begin when first silks appear and direct to the zone of ear production. Repeat applications at 2 to 3 day intervals until the silks begin to dry. Three or more applications may be required depending upon the severity of infestation. Allow 7 days between last application and harvest if corn is to be used as forage or fodder to be fed dairy or meat animals. **CAUTION:** Application of this product to the tassel region of the corn during the pollen shed period will seriously reduce bee population.

TOMATO, EGGPLANT, PEPPER: To control — Tomato fruitworm, Fall armyworm, and Flea beetles, apply the dust to thoroughly cover upper and lower leaf surfaces.

Chas Bag Co. 7853-341





tion. Repeat applications at 2 to 3 day intervals until the silks begin to dry. Three or more applications may be required depending upon the severity of infestation. Allow 7 days between last application and harvest if corn is to be used as forage or fodder to be fed dairy or meat animals. **CAUTION:** Application of this product to the tassel region of the corn during the pollen shed period will seriously reduce bee population.

TOMATO, EGGPLANT, PEPPER: To control — Tomato fruitworm, Fall armyworm, and Flea beetles, apply the dust to thoroughly cover upper and lower leaf surfaces.

CUCUMBER, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control — Pickleworm, Melonworm, Cucumber beetles, Squash bug, Flea beetles, apply the dust to thoroughly cover upper and lower leaf surfaces.

LETTUCE: To control — Flea beetles, Harlequin bug, Leafhopper, Corn earworm, Armyworm, imported cabbage worms, spittlebugs, Lygus and Stink bugs, apply the dust to thoroughly cover upper and lower leaf surfaces. On head lettuce do not apply within 3 days of harvest. On leaf lettuce do not apply within 14 days of harvest.

CARROTS: To control — Flea beetles, Leafhoppers, Spittlebugs and Stink bugs, apply the dust to thoroughly cover upper and lower leaf surfaces.

OKRA: To control — Corn earworm and Stink bugs, apply the dust to thoroughly cover upper and lower leaf surfaces.

POTATOES: To control — Colorado potato beetle, Leafhopper, Flea beetles, and Lygus bugs, apply the dust to thoroughly cover upper and lower leaf surfaces.

STRAWBERRIES: To control — Meadow Spittlebug and Strawberry leaf roller, apply the dust to thoroughly cover upper and lower leaf surfaces. Do not dust 1 day before harvest.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

10% SEVIN

FORMULA-202

GARDEN DUST

NET WT. 4 LBS.