

ACCEPTED

2 JUN 1975

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 5967-37

# OXYCOP

ACTIVE INGREDIENT	By Weight
Total Copper Expressed as Metallic Copper .....	50.0%
Copper present as Copper Oxochloride Sulfate	
INERT INGREDIENTS .....	50.0%
TOTAL.....	100.0%

USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

BURN BAG WHEN EMPTY. STAY OUT OF SMOKE.

**CAUTION:** Harmful if swallowed. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Should not be used to spray young, tender foliage as that of Peaches, Almonds and Japanese Plums, nor on varieties of Pears subject to Copper damage such as Comice and D'Anjou and Seckel.

**GROUND APPLICATION** — Vegetable and Field Crops—30 to 250 gallons. Deciduous Fruit Crops—60 to 800 gallons. (On young fruit plantings use a minimum of 15 gallons water per acre.) Where recommended to **AIR APPLICATION** — Field Crops—3 to 20 gallons. Orchard Crops—10 to 20 gallons.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

**CAUTION:**  
**Keep Out of Reach of Children**

EPA REGISTRATION NO. 5967-37 AA

EPA EST. 5967 CA-1

75

MANUFACTURED BY

**MOYER CHEMICAL COMPANY**

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA



## OXYCOP

### DIRECTION FOR USE

**ALMONDS:** For control of brown rot — Use 4 lbs. per 100 gallons acre. Apply in prebloom and petal fall stages. **Shot hole** — Use 2 lbs. plus 2 gallons of oil emulsion per 100 gallons. Apply as full spray; in spring at bud swell and at petal fall.

**APRICOTS:** For control of brown rot — Use 4 lbs. per 100 gallons acre. Apply in prebloom and petal fall stages. **Shot hole** — Use 4 to 5 lbs. per 100 gallons. Apply in fall dormant periods and in spring at red bud stage.

**NECTARINES, PEACHES:** For control of brown rot — Use 4 to 5 lbs. per 100 gallons acre. Apply in preblossom, petal fall and postblossom periods as needed. **Peach blight (shot hole), peach leaf curl** — Use 4 lbs. per 100 gallons. Apply in fall dormant period. Repeat in spring at bud swell and again at petal fall.

**PLUMS:** For control of brown rot — Use 2 1/2 lbs. per 100 gallons. Apply at popcorn and full bloom stages. Limit 6 lbs. per acre. Do not apply more than 500 gallons per acre.

**PEARS:** For control of fireblight — Use 1/4 lb. per 100 gallons. Begin at 10% bloom. Repeat at 3- to 5-day intervals until all late bloom is over. Do not apply to Anjou, Seckel, or Comice. Do not apply after petal fall.

**WALNUTS:** For control of blight — Use 3 to 5 lbs. per 100 gallons. Begin in prebloom stage. Repeat at 1% bloom and 10% to 20% pistillate bloom and post bloom as necessary. If rains are forecast during bloom, apply immediately.

**BEANS:** For control of downy mildew, pod spot — Use 3 to 4 lbs. in 25 to 100 gallons acre. Begin when disease threatens. Repeat as necessary.

**BEETS:** For control of blight, leaf spot — Use 2 1/2 lbs. per 100 gallons. Begin before disease appear. Repeat at 7- to 14-day intervals during growing season.

**CUCUMBERS, SQUASH:** For control of angular leaf spot, anthracnose, downy mildew, gummy stem blight, leaf blights, leaf spots, scab and wilt — Use 4 lbs. per 100 gallons acre. Begin when plants emerge or are established, or when disease is first reported in area. Repeat at 5- to 10-day intervals through growing season. As an aid in control of bacterial wilt — Use 3 to 4 lbs. in 50 to 100 gallons acre.

**CARROTS:** For control of leaf blight, leaf spots — Use 3 to 6 lbs. in 25 to 100 gallons acre. Begin before diseases appear. Repeat at 7- to 10- day intervals.

**CELERY:** For control of bacterial blight, early blight (Cercospora), late blight (Septoria) — Use 3 to 6 lbs. in 25 to 100 gallons acre. Begin when plants are set in field or disease is first reported in area.

**EGGPLANT:** For control of anthracnose, late blight (Phytophthora), leaf spots, phomopsis blight — Use 3 to 4 lbs. in 25 to 100 gallons acre. Begin before disease is expected to appear. Repeat at 7- to 10-day intervals as necessary.

**LETTUCE:** For control of downy mildew — Use 3 to 4 lbs. in 25 to 100 gallons acre. Begin before disease is expected to appear. Repeat at 7- to 10-day intervals as necessary.

**PEPPERS:** For control of anthracnose, bacterial spot, cercospora leaf spot, phytophthora blight — Use 2 1/2 to 4 lbs. in 25 to 100 gallons acre. Begin before disease appear. Repeat at 7- to 10-day intervals. For control of damping off — Use 4 lbs. per 100 gallons. Begin when plants emerge. Repeat at 4- to 7-day intervals as necessary. Apply to plants and surrounding soil.

**POTATOES:** For control of early and late blights — Use 3 to 4 lbs. in 25 to 100 gallons acre. Begin when plants are 4 to 6 inches high or when disease first appears in area. Repeat at 4- to 10-day intervals.

**TOMATOES:** For control of anthracnose, early and late blights, septoria leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons acre. Begin when plants are 5 to 6 inches high, when fruit is set, or when diseases are first reported in area. Repeat at 4- to 10-day intervals. In areas where blight is severe, begin when plants are set in field. For control of bacterial spot — Use 2 1/2 to 6 lbs. per 100 gallons. Begin in seed bed. Repeat at 5- to 7-day intervals after first leaves appear. In the field, repeat at 5- to 7-day intervals if the disease is known to occur in the area.

**VEGETABLE SEEDLINGS:** For control of damping off (cold frames, green houses, etc.) — Use 4 1/2 lbs. in 100 gallons. Begin when plants emerge. Repeat at 4- to 7-days intervals. Spray plants and surrounding soil. Do not use on cabbage and related seedlings.

**SUGAR BEETS:** For control of leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons acre. Begin when disease appears. Repeat as necessary.