

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eye, or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NOTE TO USER

Do not apply after petal fall to young tender foliage such as that of Peaches, Almonds, and Japanese plums, nor on varieties of Pears subject to copper damage such as D'Anjou and Seckel. Do not apply as a dry powder.

This product is compatible with most other insecticides and/or fungicides EXCEPT lime sulfur and dithio-carbamates. When used in combination with oil, OBSERVE THE PRECAUTIONS ON THE OIL LABEL.

Mixing Instructions: Fill the spray tank half-full of water, begin agitation and fill spray adjacents of product. As tank continues to fill, fill required amount of this product. Do not start agitation until the spraying operation is necessary to maintain adequate agitation.

Ground Application: Use for vegetables and fruit trees at 250 gallons per acre for deciduous fruit trees and 500 gallons per acre for ornamentals. For other uses, consult the label of 15 gallons per acre.


Air Application: Use for deciduous fruit trees at 10-20 gallons per acre. For other uses, consult the label.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry place. Do not store in areas where children or animals can have access.

Disposal: Do not reuse empty containers. Dispose of empty containers in accordance with local, state, or federal regulations.

Recycling: Containers may be recycled if they are clean and free of residue. Contact your local recycling center for more information.



OXYCOP

| | |
|--|-----------|
| ACTIVE INGREDIENT | By Weight |
| Total Copper Expressed as Metallic Copper | 50.0% |
| Copper present as Copper Oxychloride Sulfate | |
| INERT INGREDIENTS | 50.0% |
| TOTAL | 100.0% |

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

HARMFUL IF SWALLOWED, DO NOT SWALLOW OR INHALE

See Additional precautions on side panel

PRACTICAL TREATMENT: (First Aid)

- IF SWALLOWED:** Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.
- IF ON SKIN:** Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.
- IF IN EYES:** Wash with plenty of water for at least 15 minutes. Get medical attention.
- IF INHALED:** Remove to fresh air. **IF NO BREATHING:** give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

ACCEPTED


JUN 22 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5367-31

PRECAUTION

MANUFACTURED FOR

MOYER CHEMICAL COMPANY



OXYCOP

| | |
|---|-----------|
| ACTIVE INGREDIENT | By Weight |
| Total Copper Expressed as Metallic Copper | 50.0% |
| Copper present as Copper Oxide and Copper Sulfate | 50.0% |
| INERT INGREDIENTS | |
| TOTAL | 100.0% |

KEEP OUT OF REACH OF CHILDREN
CAUTION
HARMFUL IF SWALLOWED, DO NOT SWALLOW OR INHALE
 See Additional precautions on side panel
PRACTICAL TREATMENT: (First Aid)
IF SWALLOWED: Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.
IF ON SKIN: Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.
IF IN EYES: Wash with plenty of water for at least 15 minutes. Get medical attention.
IF INHALED: Remove to fresh air. **IF NOT BREATHING:** give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

ALMONDS: For control of brown rot — Use 4 lbs. per 100 gallons/acre. Apply in prebloom and petal fall stages. Shot hole — Use 2 lbs. plus 2 gallons of diluent per 100 gallons. Apply as full spray, in spring at bud swell and at petal fall.

APRICOTS: For control of brown rot — Use 4 lbs. per 100 gallons/acre. Apply in prebloom and petal fall stages. Shot hole — Use 4 to 5 lbs. per 100 gallons. Apply in fall dormant periods and in spring at red bud stage.

NECTARINES, PEACHES: For control of brown rot — Use 4 to 5 lbs. per 100 gallons/acre. Apply in prebloom and petal fall stages. Peach blight (shot hole), peach leaf curl — Use 4 lbs. per 100 gallons. Apply in fall dormant period. Repeat in spring at bud swell and again at petal fall.

PLUMS: For control of brown rot — Use 2 1/2 lbs. per 100 gallons. Apply at popcorn and fall dormant stages. 1 to 1 1/2 lbs. per acre. Do not apply more than 500 gallons per acre.

PEARS: For control of fire blight — Use 1 1/2 lbs. per 100 gallons. Begin at 10% bloom. Repeat at 3 to 5 day intervals until all late bloom is over. Do not apply to Anjou, Seckel, or Comice. Do not apply after petal fall.

WALNUTS: For control of blight — Use 3 to 5 lbs. per 100 gallons. Begin in prebloom stage. Repeat at 15% bloom and 15% to 25% post-bloom and 3 post-bloom as necessary. If rains are forecast during bloom, apply immediately.

BEANS: For control of blight in new pod spot — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin when disease threatens. Repeat as necessary.

BEETS: For control of blight, leaf spot — Use 2 1/2 lbs. per 100 gallons. Begin before disease appears. Repeat at 7 to 14 day intervals during growing season.

CUCUMBERS, SQUASH: For control of angular leaf spot, anthracnose, downy mildew, gummy stem blight, leaf blights, leaf spots, stab and wilt — Use 4 lbs. per 100 gallons. Use before plants emerge or are established or when disease is first reported in area. Repeat at 5 to 10 day intervals through growing season. As an aid in control of bacterial wilt — Use 3 to 4 lbs. in 50 to 100 gallons/acre.

CARROTS: For control of leaf blight, leaf spots — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin before disease appears. Repeat at 7 to 10 day intervals.

CELERY: For control of bacterial blight, early blight (Cercospora), late blight (Septoria) — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when plants are set in field or disease is first reported in area.

EGGPLANT: For control of anthracnose, late blight (Phytophthora), leaf spots, phomopsis blight — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease is expected to appear. Repeat at 7 to 10 day intervals as necessary.

LETTUCE: For control of downy mildew — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease is expected to appear. Repeat at 7 to 10 day intervals as necessary.

PEPPERS: For control of anthracnose, bacterial spot, cercospora leaf spot, phytophthora blight — Use 2 1/2 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease appears. Repeat at 7 to 10 day intervals. For control of damping off — Use 4 lbs. per 100 gallons. Begin when plants emerge. Repeat at 4 to 7 day intervals as necessary. Apply to plants and surrounding soil.

POTATOES: For control of early and late blights — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin when plants are 4 to 6 inches high or when disease first appears in area. Repeat at 4 to 10 day intervals.

TOMATOES: For control of anthracnose, early and late blights, septoria leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when plants are 5 to 6 inches high, when fruit is set, or when disease is first reported in area. Repeat at 4 to 10 day intervals. In areas where late blight is severe. Begin when plants are set in field. For control of bacterial spot — Use 5 1/2 to 6 lbs. per 100 gallons. Begin in seed bed. Repeat at 5 to 7 day intervals after first leaves appear in the field. Repeat at 5 to 7 day intervals if the disease is known to occur in the area.

VEGETABLE SEEDLINGS: For control of damping off (cold frames, green houses, etc) — Use 4 1/2 lbs. in 100 gallons. Begin when plants emerge. Repeat at 4 to 7 day intervals during plants and seedlings. Do not use in cabbage and related seedlings.

SUGAR BEETS: For control of leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when disease threatens. Repeat as necessary.

ACCEPTED
 22 1984
 Federal Insecticide, Fungicide and Rodenticide Act.
 Index 5367-31

PRECAUCION

MANUFACTURED FOR
MOYER CHEMICAL COMPANY

BEST DOCUMENT AVAILABLE

NOTICE TO BUYER — See express warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes the risk of use and the liability of this label. Where the use of this product is prohibited by law, there shall be no liability.