

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

WARNING. Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Protective clothing, including goggles, should be worn. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not apply directly to water. Birds and wildlife from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not allow rinseate from cleaning of equipment or disposed material to enter surface or ground water.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING.** Area treated with copper oxychloride sulfate on (date of application). Do not enter treated areas without protective clothing until dusts are settled.

NOTE TO USER

Do not apply after petal fall to young tender foliage such as that of Peaches, Almonds, and Japanese plums, nor on varieties of Pears subject to copper damage such as Comice, D'Anjou and Seckel. Do not apply as a dry powder. This product is compatible with most other insecticides and/or fungicides EXCEPT lime sulfur and dithiocarbamates. When used in combination with oil, OBSERVE THE PRECAUTIONS ON THE OIL LABEL.

MIXING INSTRUCTIONS: Fill the spray tank half-full of water, begin agitation and add spray adjuvants if needed. As tank continues to fill, add required amount of this product. Constant agitation during the spraying operation is necessary to maintain adequate suspension.


STORAGE AND DISPOSAL

PRECAUTIONS: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep product dry in closed container. Do not use empty container for any other purpose.

RESIDUAL DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty into application equipment. Then dispose of empty can in a sanitary landfill or by incineration, if allowed by State and local authorities. Do not burn. If burned, stay out of fumes.



OXYCOP

ACTIVE INGREDIENT	By Weight
Total Copper Expressed as Metallic Copper	50.0%
Copper present as Copper Oxychloride Sulfate	
INERT INGREDIENTS	50.0%
TOTAL	100.0%

Some 23 previously accepted instructions

**KEEP OUT OF REACH OF CHILDREN
WARNING — AVISO
HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE**

See Additional precautions on side panel

PRACTICAL TREATMENT: (First Aid)

IF IN EYES, flush with plenty of water. Call a physician.

IF ON SKIN, wash with plenty of soap and water. Get medical attention.

IF SWALLOWED, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

EPA Reg. No. 05967-37AA

EPA Est. 11656-CA-2

6/87

PRECAUTIONAL USUARIO: Si usted no lee ingles, no use este producto. No hasta que la etiqueta le haya sido explicada apropiadamente.

JUN 25 1987

MANUFACTURED FOR

MOYER PRODUCTS, INC.

P.O. BOX 5434 • FRESNO, CALIFORNIA 93755

As directed for the particular registered use
EPA Reg. No. 5762-37

NET WEIGHT 20 LBS.

GROUND APPLICATION: For vegetable and field crops use 20 to 250 gal. per acre. For deciduous fruit crops use 60-800 gallons per acre. (1) Young fruit trees planted use a minimum of 15 gallons spray per acre.)

AIR APPLICATION: Field crops 5-20 gallons. Orchard crops 10-20 gallons. **CAUTION:** Be sure aerial application spray concentration does not harm foliage.

ALMONDS: For control of brown rot — Use 4 lbs. per 100 gallons/acre. Apply in pre-bloom and petal fall stages. Shot-hole — Use 2 lbs. plus 2 gallons of oil emulsion per 100 gallons. Apply as full spray; in spring at bud swell and at petal fall.

APRICOTS: For control of brown rot — Use 4 lbs. per 100 gallons/acre. Apply in pre-bloom and petal fall stages. Shot-hole — Use 4 to 5 lbs. per 100 gallons. Apply in fall dormant periods and in spring at red bud stage.

NECTARINES: For control of brown rot — Use 4 to 5 lbs. per 100 gallons/acre. Apply in pre-bloom, petal fall and post-bloom periods as needed. Peach blight (shot-hole), peach leaf curl — Use 4 lbs. per 100 gallons. Apply in fall dormant period. Repeat in spring at bud swell and again at petal fall.

PLUMS: For control of brown rot — Use 2 1/2 lbs. per 100 gallons. Apply at popcorn and full bloom stages. Limit 6 lbs. per acre. Do not apply more than 500 gallons per acre.

PEARS: For control of brown rot — Use 1/2 lb. per 100 gallons. Begin at 10% bloom. Repeat at 3- to 8-day intervals until all late bloom is over. Do not apply to Anjou, Bosc, or Comice. Do not apply after petal fall.

WALNUTS: For control of blight — Use 3 to 5 lbs. per 100 gallons. Begin in pre-bloom stage. Repeat at 1% bloom and 10% to 20% post-bloom and post-bloom as necessary. If rains are forecast during bloom, apply immediately.

BEANS: For control of downy mildew, pod spot — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin when disease appears. Repeat as necessary.

BEETS: For control of blight, leaf spot — Use 2 1/2 lbs. per 100 gallons. Begin before diseases appear. Repeat at 7- to 14-day intervals during growing season.

CUCUMBERS, SQUASH: For control of angular leaf spot, anthracnose, downy mildew, downy stem blight, leaf blights, leaf spots, scab and wilt — Use 4 lbs. per 100 gallons/acre. Begin when plants emerge or are established, or when disease is first reported in area. Repeat at 5- to 10-day intervals through growing season. As an aid in control of bacterial wilt — Use 3 to 4 lbs. in 50 to 100 gallons/acre.

CARROTS: For control of leaf blight — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin before diseases appear. Repeat at 7- to 10-day intervals.

CELERY: For control of bacterial blight, early blight (Carrotspora), late blight (Septoria) — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when plants are set in field or disease is first reported in area.

EGGPLANT: For control of anthracnose, late blight (Phytophthora), leaf spots, phomopsis blight — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease is expected to appear. Repeat at 7- to 10-day intervals as necessary.

LETTUCE: For control of downy mildew — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease is expected to appear. Repeat at 7- to 10-day intervals as necessary.

PEPPERS: For control of anthracnose, bacterial spot, concave leaf spot, phyt./thra blight — Use 2 1/2 to 4 lbs. in 25 to 100 gallons/acre. Begin before disease appears. Repeat at 7- to 10-day intervals. For control of damping off — Use 4 lbs. per 100 gallons. Begin when plants emerge. Repeat at 4- to 7-day intervals as necessary. Apply to plants and surrounding soil.

POTATOES: For control of early and late blights — Use 3 to 4 lbs. in 25 to 100 gallons/acre. Begin when plants are 4 to 6 inches high or when disease first appears in area. Repeat at 4- to 10-day intervals.

TOMATOES: For control of anthracnose, early and late blights, septoria leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when plants are 3 to 6 inches high, when fruit is set, or when disease is first reported in area. Repeat at 4- to 10-day intervals. In areas where blight is severe, begin when plants are set in field. For control of bacterial spot — Use 5 1/2 to 6 lbs. per 100 gallons. Begin in seed bed. Repeat at 5- to 7-day intervals after first leaves appear. In the field, repeat at 5- to 7-day intervals if the disease is known to occur in the area.

VEGETABLE SEEDLINGS: For control of damping off (cold frames, green houses, etc.) — Use 4 1/2 lbs. in 100 gallons. Begin when plants emerge. Repeat at 4- to 7-day intervals. Spray plants and surrounding soil. Do not use on cabbage and related seedlings.

SUGAR BEETS: For control of leaf spot — Use 3 to 6 lbs. in 25 to 100 gallons/acre. Begin when disease appears. Repeat as necessary.

CONDITIONS OF SALE

1. Buyer warrants that this product complies with the requirements specified and is not adulterated. The purchaser shall be responsible for the use of this product in accordance with the directions on the label. The purchaser shall be responsible for the use of this product in accordance with the directions on the label. The purchaser shall be responsible for the use of this product in accordance with the directions on the label.

2. Because of the wide range of possible uses, proper conditions and normal use of this product is not intended to be a guarantee of performance. Buyer's failure to use the recommended amount of product or failure to purchase product in the correct quantity may be cause for improper or unexpected results.

BEST AVAILABLE COPY