




DUSTING SULFUR

DIRECTIONS FOR USE

FRUIT AND NUT CROPS

ALMONDS Brown rot, scab, shot-hole—37.5 to 45 lbs. per acre. Begin when disease threatens or in pre-bloom stage. Repeat at 7 day intervals as long as necessary.

Rust—30 to 43 lbs. per acre. Apply in pre-bloom, bloom, and cover periods as necessary.

APPLES Brown rot, powdery mildew, quince rust, scab—7.5 to 59 lbs. per acre. Apply from pre-pink stage through cover. Macintosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur applications under certain conditions.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES Powdery mildew—10 to 30 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

CHERRIES Brown rot, powdery mildew, rust, scab, silver mite—20 to 39 lbs. per acre. Begin in pink stage. Repeat at bloom, petal fall, and in cover stages at 7 to 14 day intervals.

GRAPES Powdery mildew—3.8 to 40 lbs. per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other Vitis type grapes may be injured by sulfur.

NECTARINES, PEACHES Brown rot, powdery mildew, rust, scab, shot-hole, silver mite—7.5 to 60 lbs. per acre. Apply at pink, bloom, petal fall, shuck split, shuck fall, or at 10 to 14 day intervals as long as necessary. For Brown rot of fruit, apply when fruit starts to ripen.

PLUMS, PRUNES Brown rot, powdery mildew, rust, scab, shot-hole, silver mite—4 to 60 lbs. per acre. Begin at late dormant stages for shot-hole, at pink stage for other diseases. Repeat at bloom, petal fall, shuck fall, and in cover periods as necessary. For Brown rot of fruit, dust when fruit starts to ripen.

STRAWBERRIES Powdery mildew—3 to 50 lbs. per acre. Begin at first sign of disease and repeat as necessary. Note: Sulfur may injure Strawberries under certain climatic conditions.

CITRUS Thrips—25 lbs. per acre, making up to three applications in accordance with directions of the State Agricultural Experiment Station.

VEGETABLE CROPS

THE MAXIMUM LIMITATION FOR VEGETABLES IS 50 LBS. PER ACRE.

ARTICHOKES Leaf spots—22.5 to 37.5 lbs. per acre. Apply as necessary.

ASPARAGUS Rust—6 to 35 lbs. per acre. Begin after curing stops. Repeat at 7 to 10 day intervals through season.

BEANS Leaf spot, powdery mildew, rust—3.8 to 36 lbs. per acre. Begin when first true leaves form or at first sign of disease. Repeat at 7 to 10 day intervals. Sulfur may injure certain varieties of beans.

BEETS Rust—15 to 20 lbs. per acre. Apply at first sign of disease. Repeat at 7 to 10 day intervals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER Powdery mildew—6 to 25 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

CARROTS Powdery mildew, rust—10 to 50 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

CELERY Early and late blight—6 to 12 lbs. per acre. Begin as soon as plants emerge.

GARLIC Powdery mildew—18.75 to 37.5 lbs. per acre. Begin before diseases are expected to appear. Repeat as necessary.

LETTUCE Powdery mildew—10 to 15 lbs. per acre. Begin at first sign of disease. Repeat as necessary.

ONIONS Powdery mildew—6 to 50 lbs. per acre. Begin before diseases are expected to appear. Repeat as necessary.

PEAS Powdery mildew, rust—7 to 35 lbs. per acre. Begin early in season before diseases appear. Repeat at 7 to 10 day intervals.

PEPPERS Powdery mildew—7.5 to 37.5 lbs. per acre. Begin before disease appears. Repeat at weekly intervals.

POTATOES Powdery mildew (a rare disease)—4 to 10 lbs. per acre. Apply before disease occurs. Repeat as necessary.

TOMATOES Russet mite—30 to 40 lbs. per acre in early season applications.

ACTIVE INGREDIENT	By Weight
Sulfur	98.00%
INERT INGREDIENTS	
	2.00%
Total	100.00%

THIS PRODUCT IS FORMULATED FOR APPLICATION BY GROUND OR AIR EQUIPMENT

CAUTION

KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 5967-65 AA

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

NET WEIGHT 50 LBS.

FIELD CROPS

ALFALFA Powdery mildew, rust—4 to 25 lbs. per acre. Apply to stubble or when crop is not over 3 to 4 inches high or when diseases appear. Repeat at 7 to 10 day intervals or as necessary.

BARLEY, OATS, WHEAT Rusts—25 to 40 lbs. per acre. Repeat at 7 to 10 day intervals.

CLOVER Powdery mildew, rust—10 to 18 lbs. per acre. Begin when crop is in bud stage or at first sign of disease. Repeat at 7 to 10 day intervals.

COWPEAS Rust—5 to 25 lbs. per acre. Begin soon after seedlings emerge. Repeat at 7 to 10 day intervals through season.

VETCH Rust—25 to 40 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

COTTON Strawberry (pacific) mite—20 to 40 lbs. per acre. Underleaf coverage is necessary for control. Repeat as required.

ORNAMENTALS Powdery mildew, leaf spots—20 to 40 lbs. per acre. Begin when diseases appear and repeat at 5 to 10 day intervals and after rains. Avoid application to plants sensitive to sulfur.

Sulfur is listed as a safe chemical with no time limitations. However, do not allow workers to enter treated citrus groves, vineyards, peach or hickory orchards where substantial contact will be made with treated foliage until 1 day after application.

PRECAUTIONS

Sulfur may cause severe frost and ice injury to certain crops. Examples of such sensitive plants include Apple and Citrus Pears, Apricots, peaches, plums, Cucumbers, Squash, and certain varieties of Melons, Eggplants, Spinach, Walnuts, and certain Ornamentals. Many other plants such as Apples, Pears, Strawberries, and Citrus are susceptible to foliage burn with sulfur inumping. Do not make sulfur applications at such times. Do not make sulfur applications within 2 weeks of an insect spray treatment. Do not make sulfur applications within 2 weeks of growing crops for processing. Do not use product before applying sulfur.

Do not use in areas where it may cause damage to crops or to smoke.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes a risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NON-RETURNABLE