

5967-65

PRECAUTIONARY STATEMENTS: **HAZARDS TO HUMANS AND DOMESTIC ANIMALS** **CAUTION**

Harmful if swallowed. Avoid inhaling dust. Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENTS

Do not permit workers to enter treated area where substantial contact will be made with treated foliage for 1 day after application for all crops.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing, areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Caution. Area treated with sulfur on (date of application). Do not enter without appropriate protective clothing until dusts have settled. (Insert here action to take in case of accidental exposure.)"

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep product dry in closed container. Do not use empty container for any other purpose.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by the State and local authorities, by burning. If burned, stay out of smoke.



DUSTING SULFUR **ORM-C**

ACTIVE INGREDIENT	By Weight
Sulfur	98.00%
INERT INGREDIENTS	2.00%
TOTAL	100.00%

THIS PRODUCT IS FORMULATED FOR APPLICATION BY GROUND OR AIR EQUIPMENT

KEEP OUT OF REACH OF CHILDREN **CAUTION**

HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE.

See Additional precautions on side panel

PRACTICAL TREATMENT: (First Aid)

IF SWALLOWED: Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.

IF ON SKIN: Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.

IF IN EYES: Wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove to fresh air. IF NOT BREATHING: Give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

EPA REG. NO. 5967-65 AA

Product is produced at one of the following locations as stamped on bag:
 Est. No. 35296-CA-2
 Est. No. 476-CA-3

MANUFACTURED FOR
MOYER PRODUCTS, INC.
 P.O. BOX 5434, FRESNO, CALIFORNIA 93755

NET WEIGHT 50 LBS.

PRECAUTIONS

When growing crops for processing, consult the processor before apply sulfur.

Do not apply to or allow dust to drift onto sulfur sensitive crops such as asparagus or melons. Do not apply during or prior to periods of high temperature (90° F or higher) as sulfur damage may occur. Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur unless local use has proved that sulfur does not damage crops in that locality.

Sulfur may burn foliage when temperatures are excessively high.

This product is an agricultural pesticide for application by ground or air for the below indicated uses. Where multiple rates are stated, higher rates should be used when severe infections are present or are anticipated. Do not use within 2 weeks of an oil spray treatment on any crop, or within 3 to 6 weeks of an oil spray on citrus.

FRUIT AND NUT CROPS

ALMONDS: Brown rot, scab, shothole - 37.5 to 45 lbs. per acre. Begin when disease threatens or in pre-bloom stage. Repeat at 7 day intervals as long as necessary.

APPLES: Brown rot, powdery mildew, quince rust, scab - 7.5 to 15 lbs. per acre. Apply from pre-blossom stage through cover. Macintosh, Golden Delicious, Jonathon and certain other varieties may be injured by sulfur applications under certain conditions.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES: Powdery mildew - 10 to 15 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

CHERRIES: Brown rot, powdery mildew, rust, scab, silver mite - 20 to 35 lbs. per acre. Begin in pink stage. Repeat at bloom, petal fall, and in cover stages at 7 to 14 day intervals.

GRAPES: Powdery mildew - 3.5 to 40 lbs. per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur.

NECTARINES, PEACHES: Brown rot, powdery mildew, rust, scab, shothole, silver mite - 10 to 40 lbs. per acre. Apply at pink, bloom, petal fall, shuck split, shuck fall, or at 10 to 14 day intervals as long as necessary. For brown rot of fruit, apply when fruit start to ripen.

PLUMS, PRUNES: Brown rot, powdery mildew, rust, scab, shothole, silver mite - 4 to 40 lbs. per acre. Begin at late dormant stages for shothole; at pink stage for other diseases. Repeat at bloom, petal fall, shuck fall and in cover periods as necessary. For brown rot of fruit, dust when fruit starts to ripen.

STRAWBERRIES: Powdery mildew - 3 to 50 lbs. per acre. Begin at first sign of disease and repeat as necessary. Notes: Sulfur may injure strawberries under certain climatic conditions.

CITRUS: Thrips - 125 lbs. per acre making up to three applications. Timing in accordance with directions of the State Agricultural Experiment Station.

VEGETABLE CROPS

THE MAXIMUM LIMITATION FOR VEGETABLES IS 50 LBS. PER ACRE.

ARTICHOKEs: Leaf spots - 22.5 to 37.5 lbs. per acre. Apply as necessary.

ASPARAGUS: Rust - 4 to 35 lbs. per acre. Begin after cutting stops. Repeat at 7 to 10 day intervals through the season.

BEANS: Leaf spot, powdery mildew - 4 to 25 lbs. per acre. Begin when first true leaf disease. Repeat at 7 to 10 day intervals certain varieties of beans.

BEETS: Rust - 15 to 20 lbs. per acre. Repeat at 7 to 10 day intervals.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Rust - 4 to 25 lbs. per acre. Repeat at 7 to 10 day intervals.

CARROTS: Powdery mildew, rust - Begin at first sign of disease intervals.

CELERY: Early and late blight - Begin as soon as plants emerge. Repeat as long as necessary.

CARLIS: Powdery mildew - 10 lbs. per acre. Repeat before disease appears; necessary.

LETTUCE: Powdery mildew - 10 to 15 lbs. per acre. Repeat at first sign of disease.

ONIONS: Powdery mildew - 6 to 10 lbs. per acre. Repeat before disease appears; necessary.

PEAS: Powdery mildew, rust - 7 to 10 lbs. per acre. Repeat before disease appears; necessary.

PEPPERS: Powdery mildew - 7.5 to 10 lbs. per acre. Repeat before disease appears; necessary.

POTATOES: Powdery mildew - 10 to 15 lbs. per acre. Apply before disease appears; necessary.

TOMATOES: Rust mite - 30 to 40 lbs. per acre. Repeat before disease appears; necessary.

FIELD CROPS

ALFALFA: Powdery mildew, rust - Apply to stubble or when crop is in seed stage or when disease appears. Repeat as necessary.

BARLEY, OATS, WHEAT: Rust - 25 to 30 lbs. per acre. Repeat at 7 to 10 day intervals.

CLOVER: Powdery mildew, rust - Begin when crop is in bud stage. Repeat at 7 to 10 day intervals.

COFFEES: Rust - 5 to 25 lbs. per acre. Repeat at 7 to 10 day intervals.

VEGETABLES: Rust - 25 to 40 lbs. per acre. Repeat at 7 to 10 day intervals.

COTTON: Strawberry (spiral) - Underleaf coverage is necessary.

ORNAMENTS: Powdery mildew, rust - Begin when disease appears; intervals and after rains.

CONDITIONS OF SALE:

1. Moyer warrants that this product is as labeled and is not adulterated or mislabeled. When used in accordance with directions under normal conditions, it will give the results warranted, guaranteed or directed.

2. Because of the time, place, conditions and normal or abnormal storage are beyond Moyer's control, it is limited to replacement of price. In no event shall Moyer be liable for consequential damages.

ACCEPTED

SEP 30 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5967-65

BEST AVAILABLE COPY