

Reg # 5967-65

PM-21

File
198



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 23 1994

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Terri Aal
MOYER PRODUCTS, INC.
P. O. Box 5434
Fresno, CA 93755

Subject: Label Amendment Submission of 10/18/93 in Response to PR Notice 93-7
EPA Reg. No. 5967-65
DUSTING SULFUR

IMPORTANT NOTICE: This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Recycled/Recyclable
Printed with Soy/Canola
contains at least 50%

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
Compliance Division, (7203)
Worker Protection Notification
Office of Compliance Monitoring
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

BEST AVAILABLE COPY

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling as amended

AND

- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001


BEST AVAILABLE COPY

Page 4

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

BEST AVAILABLE COPY

598

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Terri Aal
MOYER PRODUCTS, INC.
BOX 5434
FRESNO CA 93755

Comment for: EPA Reg Nr.5967-65
DUSTING SULFUR

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

The requirement under the heading "Personal Protective Equipment" must state: "Applicators and other handlers must wear:". Cross out your lead-in statement and replace it with correct one.

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

The eyewear requirement for early entry in the Agricultural Use Requirements Box on your proposed labeling is missing or incorrect. Remove the incorrect requirement (if any) and add "Protective eyewear". The exact location is shown in Sections F and G on Part 2 of the Product Worksheet in Supplement Three-A to PR Notice 93-7.

BEST AVAILABLE COPY

698

MOYER

PRODUCTS

DIRECTIONS FOR USE (Con

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid inhaling dust. Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT

Mixers, loaders, applicators, and other handlers who may be exposed to the concentrate must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, for: coveralls, waterproof gloves, and shoes plus socks.

PRECAUTIONS

When growing crops for processing, consult the processor before applying sulfur. Do not apply to or allow dust to drift onto sulfur sensitive crops such as apricots or melons. Do not apply during or prior to periods of high temperature (90°F or higher) as sulfur damage may occur. Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when temperatures are excessively high.

This product is an agricultural pesticide for application by ground or air for the below indicated uses. Where multiple rates are stated, higher rates should be used when severe infections are present or are anticipated. Do not use within 2 weeks of an oil spray treatment on any crop, or within 3 to 6 weeks of an oil spray on citrus.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Keep product dry in closed container. Do not use empty container for any other purpose.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

DUSTING SULFUR ORM-C

ACTIVE INGREDIENT	By Weight
Sulfur	98.00%
INERT INGREDIENTS	2.00%
TOTAL	100.00%

THIS PRODUCT IS FORMULATED FOR APPLICATION
BY GROUND OR AIR EQUIPMENT

EPA REG. NO. 5967-65 AA

5/89

Product is produced at one of the following locations as stamped on bag:

Est. No. 35296-CA-2

Est. No. 476-CA-5

KEEP OUT OF REACH OF CHILDREN CAUTION

HARMFUL IF SWALLOWED.
DO NOT SWALLOW OR INHALE.

See additional precautions on side panel

PRACTICAL TREATMENT (First Aid):

IF SWALLOWED: Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.

IF ON SKIN: Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.

IF IN EYES: Wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove to fresh air. **IF NOT BREATHING** give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

ACCEPTED
with COMMENTS
In EPA Letter Dated

FEB 23 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
5967-65

CONDITIONS OF SALE:

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

NET WEIGHT 50 LBS.

MANUFACTURED FOR
MOYER PRODUCTS, INC.

P.O. BOX 5434, FRESNO, CALIFORNIA 93755

FRUIT AND NUT CROPS

ALMONDS: Brown rot, scab, shot-hole — 37.5 to 45 lbs. per acre. Repeat at 7 day intervals. **Rust** — 30 to 43 lbs. per acre. Apply in pre-bloom, bloom, necessary.

APPLES: Brown rot, powdery mildew, quince rust, scab — Apply from pre-pink stage through cover. Macintosh, Golden Delicious certain other varieties may be injured by sulfur applications.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES: Powdery mildew — 10 to 30 lbs. per acre. Begin when new growth appears and before blossoms open. Repeat at 10 day intervals or as necessary. **CHERRIES:** Brown rot, powdery mildew, rust, scab — 30 lbs. per acre. Begin in pink stage. Repeat at bloom, petal fall, and 10 to 14 day intervals.

GRAPES: Powdery mildew — 38 to 40 lbs. per acre. Begin 6 inches long. Repeat before blossoms open and continue as other labrusca type grapes may be injured by sulfur. **NECTARINES, PEACHES:** Brown rot, powdery mildew, rust — 10 to 60 lbs. per acre. Apply at pink, bloom, petal fall, or at 10 to 14 day intervals as long as necessary. For when fruit start to ripen.

PLUMS, PRUNES: Brown rot, powdery mildew, rust — 4 to 60 lbs. per acre. Begin at late dormant stages for other diseases. Repeat at bloom, petal fall, shuck fall and in c. For brown rot of fruit, dust when fruit starts to ripen.

STRAWBERRIES: Powdery mildew — 3 to 50 lbs. per acre of disease and repeat as necessary. Note: Sulfur may injure certain climatic conditions.

CITRUS: Thrips — 125 lbs. per acre making up to three applications in accordance with directions of the State Agricultural Experiment Station.

VEGETABLE CROPS

THE MAXIMUM LIMITATIONS FOR VEGETABLES IS 50 LBS. PER ACRE. **ARTICHOKES:** Leaf spots — 22.5 to 37.5 lbs. per acre. **ASPARAGUS:** Rust — 6 to 35 lbs. per acre. Begin after 7 to 10 day intervals through the season.

BEANS: Leaf spot, powdery mildew, rust — 38 to 36 lbs. per acre. Begin at first true leaves form or at first sign of disease. Repeat at 7 to 10 day intervals.

BEETS: Rust — 15 to 20 lbs. per acre. Apply at first sign of disease. Repeat at 7 to 10 day intervals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWERS: Powdery mildew, rust — 6 to 25 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

CARROTS: Powdery mildew, rust — 10 to 50 lbs. per acre. Repeat at 7 to 10 day intervals.

CELERY: Early and late blight — 6 to 12 lbs. per acre. Repeat at 7 to 10 day intervals as long as necessary.

GARLIC: Powdery mildew — 18.75 to 37.5 lbs. per acre. Repeat as necessary.

LETTUCE: Powdery mildew — 10 to 15 lbs. per acre. Repeat as necessary.

ONIONS: Powdery mildew — 6 to 50 lbs. per acre. Repeat as necessary.

PEAS: Powdery mildew, rust — 7 to 35 lbs. per acre. Repeat at 7 to 10 day intervals.

PEPPERS: Powdery mildew — 7.5 to 37.5 lbs. per acre. Repeat at weekly intervals.

POTATOES: Powdery mildew (a rare disease) — 4 to 12 lbs. per acre. Repeat as necessary.

TOMATOES: Rust — 30 to 40 lbs. per acre in early season.

FIELD CROPS

ALFALFA: Powdery mildew, rust — 6 to 25 lbs. per acre. Repeat at 7 to 10 day intervals or when diseases appear.

BARLEY, OATS, WHEAT: Rust — 25 to 40 lbs. per acre. Repeat at 7 to 10 day intervals.

CLOVER: Powdery mildew, rust — 12 to 18 lbs. per acre. Repeat at 7 to 10 day intervals.

COWPEAS: Rust — 5 to 25 lbs. per acre. Begin soon after 7 to 10 day intervals through the season.

VETCH: Rust — 25 to 40 lbs. per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

COTTON: Strawberry (pacific) mite — 20 to 40 lbs. per acre. Repeat as necessary.

ORNAMENTALS: Powdery mildew, leaf spots — 20 to 40 lbs. per acre. Repeat at 5 to 10 day intervals and as necessary to plants sensitive to sulfur.

BEST AVAILABLE COPY

MOYER

PRODUCTS

798
DIRECTIONS FOR USE (Continued)

CAUTIONARY STATEMENTS HUMANS AND DOMESTIC ANIMALS CAUTION

Avoid inhaling dust. Avoid contact with skin, eyes, and clothing. Immediately flush eyes and skin with plenty of water if irritation persists.

PROTECTIVE EQUIPMENT

Wear long-sleeved shirt and long pants, waterproof boots, protective eyewear.

Instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use hot water and wash PPE separately from other laundry. Do not use in enclosed areas, or aircraft in a manner prohibited in the Worker Protection Standard (WPS) (40 CFR 170.240 (d)(4-6)). The handler PPE must be used or modified as specified in the WPS.

SAFETY RECOMMENDATIONS:

Wash hands before eating, drinking, chewing gum, using the toilet. Immediately if pesticide gets inside. Then wash with soap and water. Wash the outside of clothing. Wash the outside of clothing. Wash the outside of clothing.

ENVIRONMENTAL HAZARDS

Do not apply when weather conditions favor drift. Do not contaminate water by cleaning equipment or disposing of this product only as specified on this label.

DIRECTIONS FOR USE

Follow label to use this product in a manner consistent with labeling.

Do not use in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may use this product. For any requirements specific to this product, consult the agency responsible for pesticide regulation.

CULTURAL USE REQUIREMENTS

Use in accordance with its labeling and with the restrictions, 40 CFR part 170. This Standard contains minimum requirements for agricultural workers on farms, forests, nurseries, and handlers of agricultural pesticides. It also contains specific instructions and other requirements on this label about personal protective equipment and restricted-entry interval. The requirements in this label that are more restrictive than those in the Standard apply to this product that are covered by the Worker Protection Standard.

Do not enter treated areas during the restricted entry interval (REI).

Do not enter treated areas that is permitted under the Standard and that involves contact with anything that has been treated, such as plants, soil, or water, for: coveralls, waterproof boots.

PRECAUTIONS

Before processing, consult the processor before applying sulfur to drift onto sulfur sensitive crops such as almonds, apricots, or prior to periods of high temperature (90°F or higher) or when some fruit and other plants are susceptible to injury under climatic conditions. The user is advised not to use sulfur where that sulfur does not damage crops in that locality when temperatures are excessively high.

Use sulfur as a foliar pesticide for application by ground or air for the below multiple rates are stated. Higher rates should be used when insects or diseases are anticipated. Do not use within 2 weeks of an oil spray on citrus.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not use empty container. Do not use empty container.

AL: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

AL: Completely empty bag into application equipment bag in a sanitary landfill or by incineration, or if allowed by local authorities, by burning. If burned, stay out of smoke.

DUSTING SULFUR FORM-C

ACTIVE INGREDIENT	By Weight
Sulfur	98.00%
INERT INGREDIENTS	2.00%
TOTAL	100.00%

THIS PRODUCT IS FORMULATED FOR APPLICATION
BY GROUND OR AIR EQUIPMENT

EPA REG. NO. 5967-65 AA 5/89
Product is produced at one of the following locations as stamped on bag:

Est. No. 35296-CA-2 Est. No. 476-CA-5

KEEP OUT OF REACH OF CHILDREN CAUTION

HARMFUL IF SWALLOWED.
DO NOT SWALLOW OR INHALE.

See additional precautions on side panel

PRACTICAL TREATMENT (First Aid):

IF SWALLOWED: Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.

IF ON SKIN: Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.

IF IN EYES: Wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove to fresh air. IF NOT BREATHING: give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

ACCEPTED
with COMMENTS
by EPA Letter Dated

FEB 23 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

5967-65

CONDITIONS OF SALE:

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

NET WEIGHT 50 LBS.

MANUFACTURED FOR
MOYER PRODUCTS, INC.
PO. BOX 5434, FRESNO, CALIFORNIA 93755

FRUIT AND NUT CROPS

ALMONDS: Brown rot, scab, shothole — 375 to 45 lbs per acre. Begin when disease threatens or in pre bloom stage. Repeat at 7 day intervals as long as necessary. Rust — 30 to 43 lbs per acre. Apply in pre bloom bloom, and cover periods as necessary.

APPLES: Brown rot, powdery mildew, quince rust, scab — 75 to 59 lbs per acre. Apply from pre pink stage through cover. Macintosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur applications under certain conditions.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES: Powdery mildew — 10 to 30 lbs per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

CHERRIES: Brown rot, powdery mildew, rust, scab, silver scale — 20 to 59 lbs per acre. Begin in pink stage. Repeat at bloom, petal fall, and in cover stages at 7 to 14 day intervals.

GRAPES: Powdery mildew — 38 to 40 lbs per acre. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur.

NECTARINES, PEACHES: Brown rot, powdery mildew, rust, scab, shothole, silver scale — 10 to 60 lbs per acre. Apply at pink bloom, petal fall, shuck split, shuck fall, or at 10 to 14 day intervals as long as necessary. For brown rot of fruit, apply when fruit start to open.

PLUMS, PRUNES: Brown rot, powdery mildew, rust, scab, shothole, silver scale — 4 to 60 lbs per acre. Begin at late dormant stages for shothole, at pink stage for other diseases. Repeat at bloom, petal fall, shuck fall and in cover periods as necessary. For brown rot of fruit, dust when fruit starts to ripen.

STRAWBERRIES: Powdery mildew — 3 to 50 lbs per acre. Begin at first sign of disease and repeat as necessary. Note: Sulfur may injure strawberries under certain climatic conditions.

CITRUS: Thrips — 125 lbs per acre making up to three applications. Timing in accordance with directions of the State Agricultural Experiment Station.

VEGETABLE CROPS

THE MAXIMUM LIMITATIONS FOR VEGETABLES IS 50 LBS PER ACRE.

ARTICHOKES: Leaf spots — 22.5 to 37.5 lbs per acre. Apply as necessary. ASPARAGUS: Rust — 6 to 35 lbs per acre. Begin after cutting stops. Repeat at 7 to 10 day intervals through the season.

BEANS: Leaf spot, powdery mildew, rust — 38 to 36 lbs per acre. Begin when first true leaves form or at first sign of disease. Repeat at 7 to 10 day intervals. Sulfur may injure certain varieties of beans.

BEETS: Rust — 15 to 20 lbs per acre. Apply at first sign of disease. Repeat at 7 to 10 day intervals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Powdery mildew — 6 to 25 lbs per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

CARROTS: Powdery mildew, rust — 10 to 50 lbs per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

CELERY: Early and late blight — 6 to 12 lbs per acre. Begin as soon as plants emerge. Repeat at 7 to 10 day intervals as long as necessary.

GARLIC: Powdery mildew — 18.75 to 37.5 lbs per acre. Begin before diseases are expected to appear. Repeat as necessary.

LETTUCE: Powdery mildew — 10 to 15 lbs per acre. Begin at first sign of disease. Repeat as necessary.

ONIONS: Powdery mildew — 6 to 59 lbs per acre. Begin before diseases are expected to appear. Repeat as necessary.

PEAS: Powdery mildew, rust — 7 to 35 lbs per acre. Begin early in season before diseases appear. Repeat at 7 to 10 day intervals.

PEPPERS: Powdery mildew — 75 to 37.5 lbs per acre. Begin before disease appears. Repeat at weekly intervals.

POTATOES: Powdery mildew (a rare disease) — 4 to 12 lbs per acre. Apply before disease occurs. Repeat as necessary.

TOMATOES: Rust — 30 to 40 lbs per acre in early season applications.

FIELD CROPS

ALFALFA: Powdery mildew, rust — 6 to 25 lbs per acre. Apply to stubble or when crop is not over 3 to 6 inches high or when diseases appear. Repeat at 7 to 10 day intervals or as necessary.

BARLEY, OATS, WHEAT: Rust — 25 to 40 lbs per acre. Repeat at 7 to 10 day intervals.

CLOVER: Powdery mildew, rust — 12 to 18 lbs per acre. Begin when crop is in bud stage or at first sign of disease. Repeat at 7 to 10 day intervals.

COWPEAS: Rust — 5 to 25 lbs per acre. Begin soon after seedlings emerge. Repeat at 7 to 10 day intervals through the season.

VETCH: Rust — 25 to 40 lbs per acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

COTTON: Strawberry (spiral) mite — 20 to 40 lbs per acre. Underleaf coverage is necessary to control. Repeat as necessary.

ORNAMENTALS: Powdery mildew, leaf spots — 20 to 40 lbs per acre. Begin when diseases appear and repeat at 5 to 10 day intervals and after rains. Avoid application to plants sensitive to sulfur.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid inhaling dust. Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENTS

Do not permit workers to enter treated area where substantial contact will be made with treated foliage for 1 day after application for all crops.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should specify protective clothing, areas or fields that cannot be entered without this clothing, period of time area is to be vacated, and statement of practical treatment in case of exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "Caution—Area treated with Sulfur-on (date of application). Do not enter without appropriate protective clothing until dusts have settled. (Insert here action to be taken in case of accidental exposure)."

PRECAUTIONS

When growing crops for processing, consult the processor before applying sulfur. Do not apply to or allow dust to drift onto sulfur sensitive crops such as apricots or melons. Do not apply during or prior to periods of high temperature (90°F or higher) as sulfur damage may occur. Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when temperatures are excessively high.

This product is an agricultural pesticide for application by ground or air for the below indicated uses. Where multiple rates are stated, higher rates should be used when severe infections are present or are anticipated. Do not use within 2 weeks of an oil spray treatment on any crop, or within 3 to 6 weeks of an oil spray on citrus.

FRUIT AND NUT CROPS

ALMONDS: Brown rot, scab, shot-hole — 37.5 to 45 lbs. per acre. Begin when disease threatens or in pre-bloom stage. Repeat at 7 day intervals as long as necessary. Rust — 30 to 43 lbs. per acre. Apply in pre-bloom, bloom, and cover periods as necessary.

APPLES: Brown rot, powdery mildew, quince rust, scab — 7.5 to 59 lbs. per acre. Apply from pre-pink stage through cover. Macintosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur applications under certain conditions.

Continued on right panel

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Keep product dry in closed container. Do not use empty container for any other purpose.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

MOYER

PRODUCTS

DUSTING SULFUR ORM-C

ACTIVE INGREDIENT	By Weight
Sulfur	98.00%
INERT INGREDIENTS	2.00%
TOTAL	100.00%

THIS PRODUCT IS FORMULATED FOR APPLICATION
BY GROUND OR AIR EQUIPMENT

EPA REG. NO. 5967-65 AA

5/89

Product is produced at one of the following locations as stamped on bag:

Est. No. 35296-CA-2

Est. No. 476-CA-5

KEEP OUT OF REACH OF CHILDREN CAUTION

HARMFUL IF SWALLOWED.
DO NOT SWALLOW OR INHALE.

See additional precautions on side panel.
PRACTICAL TREATMENT (First Aid):

IF SWALLOWED: Administer water freely and induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately.

IF ON SKIN: Rinse off material and wash skin with soap and water. Get medical attention if irritation persists.

IF IN EYES: Wash with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove to fresh air. **IF NOT BREATHING:** give artificial respiration preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

CONDITIONS OF SALE:

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

NET WEIGHT 50 LBS.

MANUFACTURED FOR
MOYER PRODUCTS, INC.
PO BOX 5434, FRESNO, CALIFORNIA 93755

DIRECTIONS FOR

BLACKBERRIES, BOYSENBERRIES:

Powdery mildew — 10 to 30 lbs. per acre and before blossoms open. Repeat at 14 day intervals.

CHERRIES: Brown rot, powdery mildew — 10 to 30 lbs. per acre. Begin in pink stage. Repeat at 14 day intervals.

GRAPES: Powdery mildew — 38 to 40 lbs. per acre. Repeat before blossoms open. 6 inches long. Repeat before blossoms open. Other Vitis type grapes may be injured.

NECTARINES, PEACHES: Brown rot, mildew — 10 to 60 lbs. per acre. Apply at fall, or at 10 to 14 day intervals as long when fruit start to ripen.

PLUMS, PRUNES: Brown rot, powdery mildew — 4 to 60 lbs. per acre. Begin at late dormancy. Repeat at bloom, petal fall, for brown rot of fruit, dust when fruit start to ripen.

STRAWBERRIES: Powdery mildew — of disease and repeat as necessary. Note: tan climatic conditions.

CITRUS: Thrips — 125 lbs. per acre in accordance with directions of the State.

VEGETABLES

THE MAXIMUM LIMITATIONS FOR VEGETABLES:

ARTICHOKES: Leaf spots — 22.5 to 30 lbs. per acre.

ASPARAGUS: Rust — 6 to 35 lbs. per acre. 7 to 10 day intervals through the season.

BEANS: Leaf spot, powdery mildew, rust. First true leaves form or at first sign of disease. May injure certain varieties of beans.

BEETS: Rust — 15 to 20 lbs. per acre. 7 to 10 day intervals.

BROCCOLI, BRUSSELS SPROUTS: Mildew — 6 to 25 lbs. per acre. Begin at intervals.

CARROTS: Powdery mildew, rust — 10 to 15 lbs. per acre. Repeat at 7 to 10 day intervals.

CELERY: Early and late blight — 6 to 10 lbs. per acre. Repeat at 7 to 10 day intervals.

GARLIC: Powdery mildew — 18.75 to 30 lbs. per acre. Repeat as necessary.

LETTUCE: Powdery mildew — 10 to 15 lbs. per acre. Repeat as necessary.

ONIONS: Powdery mildew — 6 to 50 lbs. per acre. Repeat as necessary.

PEAS: Powdery mildew, rust — 7 to 35 lbs. per acre. Repeat at 7 to 10 day intervals.

PEPPERS: Powdery mildew — 7.5 to 30 lbs. per acre. Repeat at weekly intervals.

POTATOES: Powdery mildew (a rare disease occurs). Repeat as necessary.

TOMATOES: Rust — 30 to 40 lbs. per acre.

FIELD CROPS

ALFALFA: Powdery mildew, rust — 6 to 10 lbs. per acre. Repeat at 7 to 10 day intervals or as necessary.

BARLEY, OATS, WHEAT: Rusts — 2 to 10 lbs. per acre. Repeat at 7 to 10 day intervals.

CLOVER: Powdery mildew, rusts — 12 to 15 lbs. per acre. Repeat at first sign of disease.

COWPEAS: Rust — 5 to 25 lbs. per acre. Repeat at 7 to 10 day intervals through the season.

VETCH: Rust — 25 to 40 lbs. per acre. Repeat at 7 to 10 day intervals.

COTTON: Strawberry (pacific) mite — 1 to 2 lbs. per acre. Repeat as necessary.

ORNAMENTALS: Powdery mildew, leaf diseases appear and repeat at 5 to 10 day intervals to plants sensitive to sulfur.

BEST AVAILABLE COPY