

CHLORDANE 40-W

ACTIVE INGREDIENTS
Technical Chlordane*

By Weight
40.0%
60.0%
100.0%

TOTAL

*Equivalent to 24% Octachloro 4,7-Methanotetrahydroindane and 16% related compounds.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing of dust or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Do not store or transport with feed or food. Keep children and pets off treated area until material is watered into the soil and grass is dry. Forage crops treated with this material should not be fed to dairy animals or to animals being finished for slaughter. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

This product will kill fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions. — NON-RETURNABLE —

Caution: Keep Out of Reach of Children

EPA REG. NO. 5967-69

NET WEIGHT 5 LBS.

MANUFACTURED BY

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

Moyer Chemical Co. — Chlordane 40-W

DIRECTIONS FOR MIXING

Mix in water with agitation; agitate solution while spraying. Apply to row and field crops in 20 to 100 gallons of water per acre by ground spray rig or 5 to 15 gallons of water per acre by airplane. Dosages for tree crops are for use with conventional sprayers applying 100 gallons or more per acre. For use with concentrate sprayers applying 40 to 90 gallons per acre, adjust dosage to apply the same amount of actual material per acre as would be applied by a conventional sprayer.

Repeat treatment as necessary. Do not apply to food crops within 30 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, wiping or other effective means.

DIRECTIONS FOR USE

CHLORDANE 40-W is a wettable powder formulation for use in control of certain insects infesting crops, lawns, turf, parks, cemeteries, nursery stock, ornamental plants and buildings.

LAWNS, TURF, PARKS, CEMETERIES, NURSERY STOCK AND ORNAMENTAL PLANTS — To control ants, armyworms, crickets, grasshoppers and spiders, apply 2½ lbs. of Chlordane 40-W per acre. For chiggers and European earwigs, use 2½ to 3¼ lbs. per acre on lawns or turf. For chinch bugs use 5 to 6 lbs. per acre on lawns and turf. For sod webworm use 5 to 7½ lbs. per acre. On lawn, soil or turf uses, the material should be washed into the soil or turf following the application. On foliage applications, do not allow dairy animals or animals being fattened for slaughter to graze on the treated area. Do not feed treated forage to these animals.

BLACKBERRIES, LOGANBERRIES, RASPBERRIES AND STRAWBERRIES — To control strawberry root weevil, white-fringed beetle, white grubs and wireworms, apply 25 lbs. of Chlordane 40-W per acre in sufficient water for coverage. Apply to the soil and work into the ground prior to or at time of planting.

BUILDING INTERIORS — To control ants, crickets, roaches and silverfish apply as a spray in water at the rate of 2 lbs. of this material in 5 gallons of water. Use a coarse wet spray and make applications on a "spot" basis to the cracks, surface or other areas where the insects are seen. Do not apply to large areas of premises. Avoid food contamination. Repeat treatment when necessary.

GRAPES: Darkling Ground Beetles, Bud Beetles — 8 to 10 lbs. per acre in water to cover. Do not apply after edible parts start to form.

LETTUCE — Leafminer, Lygus Bugs. Use 3 to 4 lbs. per acre in water to cover. Do not apply after edible parts start to form.

Use designated amount of product in the following gallons of water per acre unless otherwise noted: GROUND APPLICATION — Vegetable and field crops—30 to 250 gallons. Deciduous Fruit Crops—60 to 800 gals. (On young fruit plantings—Use a minimum of 15 gals. water per acre.) Where recommended by AIR APPLICATION — Field Crops—3 to 20 gals. Orchard Crops—10 to 20 gals.

3 73

ACCEPTED