

STORE IN COOL, DRY PLACE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTION, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION — Vegetable and Field Crops: 100 to 200 (On young fruit plantings)
Deciduous Fruit Crops: 50 to 100 (On young fruit plantings)
Citrus: 100 to 200 (On young fruit plantings)

Where Recommended for A.R. APPLICATION — Field Crops: 100 to 200
A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly soluble materials such as lime or Bordeaux.
MIXING DIRECTIONS: Add to spray tank when it is full with agitator running and maintain agitation until spraying is completed.

ALFALFA: Grasshoppers, Leafhoppers, Ligus Bugs, Alfalfa Webworms, Armyworms, Alfalfa Mite, Spider Mites, Mites, Aphids, Aphis, and other insects. By A.R.

BEANS (Dry or Field), SOYBEANS, LIMA BEANS AND PEAS for Processing: Alfalfa Mite, Aphids, Spider Mites, Lygus Leaf Miner, GREEN BEANS: Aphids, Spider Mites, Looper, Nymphs, Leafhoppers, Mites, and other insects. Large Looper, Mite Bugs.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, PALE COLLARDS: Imported Cabbage worm, Diamond-Back Caterpillars, Aphids, Looper.

CELERY: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners.

COTTON: Lygus, Bollworms, Loopers and For Suppression of Mites. For HEAVY POPULATION OF LISTED INSECTS.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetle, Aphids, Spider Mites, Loopers, Thrips, Armyworms, Pickworms, Leaf Miner.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Beetle Beetles, Flea Beetles, Leaf Miners, Whiteflies.

GRAPES: Dirosiphia, Leafhoppers.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm.
MELONS (Cantaloupe, Honeydew, Muskmelon, Watermelon), PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miner.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: Aphis, Salt Brown Scale, Citrus Mite, Citrus Cutworms, Leaf Roller (Tortrix), Thrips (Bloom).

PEACHES (Western States Only): Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only).

RICE: Grasshoppers.

SPINACH, CHARD, TURNIP GREENS: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miner.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest).

SUGAR BEETS: Spider Mites, Leafhoppers.

TOBACCO: Hornworm.

TOMATO: Fruit Flies (Drosophila), Field and Green House, Tomato Fruit Worm, Hornworm, Leaf Mites, Flea Beetles, Mites.

WALNUTS (CALIF. ONLY): Aphids.

FOREST AND SHADE TREES, CONIFERS (Alder, Fir, Douglas Fir, Hemlock, Juniper, Pine, Spruce, Larch, Fir, Redwood, Red Spruce, Mite, Aphid, Spittlebug, Zimmerman Moth, Juniper Webworm, Ips, Bark Beetle, Spruce Budworm, Adult Sawfly, Western Hemlock Bark Borer): Broadleaf Trees (Ash, Birch, Bass, Walnut, Box Elder, Dogwood, Elm, Evergreen, Pear, Flowering Dogwood, Gum, Magnolia, Maple, Oak, Sycamore, Willow): Suppression of Two Spine and Red Spider Mites, Aphids, Leaf Miners, Thrips, Flea Leaf Beetles, Citrus Greening (California), Spring and Fall Caterpillars, Fall Webber, Oak Webber, and Tortoise Moth.

GREENHOUSE PLANTS, TREATMENT OF ROSES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Aphis, Whitefly, Aphid, Leafhopper, Mite, Flea Beetle.

FRUIT FLIES (Drosophila): In and around Food Processing Plants, Loading Docks, Cellars and Refuse Areas. Control Mites.

WARNING: Do not apply to plants in bloom.

CAUTION: Do not use in areas where children or livestock are present.

DANGER: Harmful if inhaled. Avoid contact with eyes, nose, and mouth. Wash thoroughly after handling.

CLOVER MITES, ROACHES, FLEAS, TICKS, AND OTHER INSECTS: Do not apply to plants in bloom.

CAUTION: Do not use in areas where children or livestock are present.

DO NOT USE: Harmful if inhaled. Avoid contact with eyes, nose, and mouth.

DANGER: Harmful if inhaled. Avoid contact with eyes, nose, and mouth.

DO NOT STORE: Do not store in areas where children or livestock are present.

DO NOT APPLY: Do not apply to plants in bloom.

DO NOT PERMIT WORK: Do not permit work in areas where children or livestock are present.

CONTACT WILL BE MAINTAINED: Contact will be maintained in areas where children or livestock are present.

DAYS HAVE LAPSED: Days have lapsed in areas where children or livestock are present.