

ACCEPTED

MAR 07 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5967-85

# ENDOSULFAN-SULFUR DUST

## DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is an agricultural pesticide for application by ground or air for the following uses:

**GRAPES:** For control of Grape Leafhoppers, Powdery Mildew — Use 25 - 35 lbs. per acre. Do not use on Concord variety of Grapes as severe injury is likely to occur. Apply when insects first appear and repeat every 5 to 7 days as necessary. Mildew control should begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Do not apply within 7 days of harvest.

**PEACHES:** For control of Green Peach Aphid, Peach Twig-Borer and Powdery Mildew—Use 50 - 60 lbs. per acre. Apply to give thorough coverage to all plant parts. For Green Peach Aphid—Apply when insects first appear. Repeat every 5 to 7 days as necessary to secure control. For Mildew control—Apply at pink, bloom, petal fall, shuck split or at 10 to 14 days intervals as long as necessary. For Peach Twig-Borer—Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 30 days of harvest, nor make more than 2 applications during the fruiting period.

**TOMATOES:** For control of Colorado Potato Beetles, Flea Beetles, Green Peach Aphid and Whitefly—Use 12 lbs. per acre. For control of Cabbage Looper—Use 17 lbs. per acre. For control of Stink Bugs and Tomato Hornworms—Use 15 - 25 lbs. per acre. For control of Yellow Striped Armyworm—use 25 lbs. per acre. Apply when insects first appear and repeat at 10 to 14 day intervals until control is achieved. Do not apply within 1 day of harvest.

**STRAWBERRIES:** For control of Strawberry Aphid—Use 15 to 30 lbs. per acre. For control of Meadow Spittlebug—Use 15 to 25 lbs. per acre. For control of Cyclamen Mites—Use 50 lbs. per acre. Dust crown of each plant thoroughly. Begin dusting when insects first appear. Repeat applications until control is achieved. Do not apply at intervals of less than 35 days when fruit is present nor within 4 days of picking.

# NO. 4-25

ACTIVE INGREDIENTS	By Weight
Endosulfan* .....	4.0%
Sulfur .....	25.0%
INERT INGREDIENTS .....	71.0%
TOTAL.....	100.0%

\*Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide.

## CAUTION:

### KEEP OUT OF REACH OF CHILDREN

SEE OTHER PRECAUTIONS ON SIDE PANEL

NET WEIGHT 50 LBS.

EPA REGISTRATION NO. 5967-85 AA

EPA EST. 5967-CA-1

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Buyer assumes all risk of use and or handling when such use and/or handling is contrary to label instructions. — NON-RETURNABLE —

1 75

MANUFACTURED BY

## MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

## CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

In case of contact with skin or eyes, flush with plenty of water: FOR EYES, get medical attention.

Wash with soap and water after handling and before eating or smoking: Wear clean clothing.

Do not store or transport near feed or food.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated areas the day of application should wear protective clothing.

Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when temperatures are excessively high.

Do not allow workers to enter treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes 1 day (24 hrs.), Peaches 1 day (24 hrs.).

Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

**HIGHLY TOXIC TO FISH AND WILDLIFE.** Birds feeding on treated areas may be killed. Do not contaminate any body of water. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**THIS PRODUCT IS HIGHLY TOXIC TO BEES** and should not be applied when bees are actively visiting the area.