

# DIRECTIONS FOR USE:

This product is an agricultural pesticide for the following uses:

**ALFALFA (SEED):** Armyworms, Cutworms, Alfalfa Caterpillar—2 to 3 pints per acre. Lygus Bugs—3 to 3.2 pints per acre. Leafhoppers, Thrips—1 to 2 pints per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**GRASS, LAWN AND TURF:** Armyworms, Chiggers, Cutworms—2 to 3 pints per 100 gallons of water per acre. Non-agricultural areas only. Keep children and pets off treated areas until material is washed into soil and grass is dry.

**CORN:** Armyworms, Corn Earworm (in whorl), Cutworms, Webworms—3 to 6 pints per acre. Grasshoppers, Crickets—1 pint per 100 lbs. of grain. Apply 10 to 20 lbs. per acre. **Chinch Bug, Flea Beetle**—3 to 6 pints per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. No time limitations on use of grain.

**COTTON:** Boll Worm, Cutworms, Armyworms, Grasshoppers, Flea Beetles—3 to 4 pints per acre. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**SORGHUM:** Armyworms, Cutworms—2 to 3 pints per acre. Grasshoppers—1 pint per 100 lbs. grain. Apply 10 to 20 lbs. per acre. Do not apply within 40 days of harvest. Do not apply more than once after heads start to form. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not ensile treated forage.

**SMALL GRAINS: BARLEY, OATS, RYE, WHEAT:** Armyworms, Cutworms—2 to 3 pints per acre. Grasshoppers—1 pint per 100 lbs. of grain. Apply 10 to 20 lbs. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on use of grain, on barley, rye and wheat.

**BEANS: DRY, GREEN AND LIMA:** Cutworms, Darkling Ground Beetle—3 to 4 pints per acre. Salt Marsh Caterpillar—4 pints per acre. Cucumber Beetle, Lygus Bugs—3 to 6 pints per acre. Grasshoppers—3 to 6 pints per acre. No limitations on use of shelled beans as human food. Do not apply to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER:** Cabbage Looper, Imported Cabbage Worm, Diamond Back Moth Larvae, Armyworms, Cutworms—3 to 4 pints per acre. Do not apply to Broccoli, Brussels Sprouts, or Cauliflower after edible parts begin to form or within 30 days of harvest.

**CARROTS:** Cutworms, Flea Beetles, Grasshoppers, Lygus Bugs (seed crop)—3 to 5 pints per acre. No time limitations. Do not feed treated tops to dairy animals or animals being finished for slaughter.

**CABBAGE:** Cabbage Looper, Imported Cabbage Worm, Diamond Back Moth Larvae, Armyworms, Cutworms—3 to 4 pints per acre. Do not apply after heads begin to form.

**CITRUS: ORANGES, LEMONS, GRAPEFRUIT, TANGERINES:** Citrus Cutworms—3 to 5 quarts per acre applied as an outside coverage spray in 100 to 400 gallons of water by conventional orchard sprayer. Do not apply within 30 days of harvest. Do not apply during bloom period. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**CELERY:** Cutworms, Celery Leaf Tier, Vegetable Weevil—4 to 5 pints per acre. Do not apply after plants start to bunch or after plants are half mature.

**COLLARDS, KALE:** Armyworms, Imported Cabbage Worm, Cutworms, Grasshoppers—3 to 4 pints per acre. Do not apply within 28 days of harvest. Cabbage Looper, Flea Beetle—3 to 4 pints per acre. Do not apply within 35 days of harvest.

**EGGPLANT, PEPPERS:** Armyworms, Cucumber Beetle, Cutworms, Flea Beetle—3 to 3.5 pints per acre. Do not apply within 5 days of harvest.

**KOHLRABI:** Cutworms, Armyworms, Imported Cabbage Worm, Cabbage Looper—3 to 4 pints per acre. Do not apply after edible parts start to form or within 30 days of harvest.

**LETTUCE HEAD and LEAF:** Armyworms, Cabbage Looper, Salt Marsh Caterpillar, Cucumber Beetle—3 to 5 pints per acre. Grasshoppers—4 pints per acre. Do not apply after seeding stage on leaf lettuce. Do not apply to head lettuce after heads begin to form.

**OKRA:** Cutworms, Leafminer—3 to 5 pints per acre. Do not apply after pods begin to form.

**ONION (DRY ONLY):** Thrips—1 to 2 pints per acre. Do not apply to green or spring onions.

**PEAS:** Armyworms, Lygus Bugs—3 to 4 pints per acre. Do not apply after pods begin to form. Do not use as food for within 7 days of harvest. Do not use as feed for dairy animals or animals being finished for slaughter.

**PUMPKIN:** Flea Beetle, Leafminer, Salt Marsh Caterpillar, Webworm—3 to 4 pints per acre. Do not apply after pods begin to form.

**RUTABAGA:** Flea Beetle, Leafminer—3 to 4 pints per acre. Do not use treated tops for food or feed.

**TOMATOES:** Armyworms, Imported Cabbage Worm, Cabbage Looper, Flea Beetle, Salt Marsh Caterpillar, Webworm—3 to 4 pints per acre. Do not apply after fruit begins to form.

**SPINACH:** Armyworms, Imported Cabbage Worm, Cabbage Looper, Flea Beetle, Salt Marsh Caterpillar, Webworm—3 to 4 pints per acre. Do not apply after leaves begin to form.

**STRAWBERRY:** Flea Beetle, Leafminer—3 to 4 pints per acre. Do not apply after fruit begins to form.

**LADINO CLOVER:** Flea Beetle, Leafminer—3 to 4 pints per acre. Do not apply after leaves begin to form.

**ORNAMENTALS:** Big Leaf Beetle, Flea Beetle, Leafminer—3 to 4 pints per acre. Do not use treated plants for food or feed.

ACCEPTED

af Tier, Vegetable Weevil—4 to 5 pints per acre. Do not  
ch or after plants are half mature.

as. Imported Cabbage Worm, Cutworms, Grasshoppers—  
within 28 days of harvest. Cabbage Looper, Flea Beetle—  
apply within 35 days of harvest.

orms, Cucumber Beetle, Cutworms, Flea Beetle—2 to 3  
within 5 days of harvest.

worms Imported Cabbage Worm, Cabbage Looper—  
apply after edible parts start to form or within 30 days

Armyworms, Cabbage Looper, Salt Marsh Caterpillar,  
per acre. Grasshoppers—4 pints per acre. Do not apply  
lettuce. Do not apply to head lettuce after heads begin

3 to 5 pints per acre. Do not apply after pod begin

Do not apply to green or spring

lygids, Bugs—1 to 2 pints per acre. Do not apply after  
are to be used as food, or within 7 days of harvest if  
food. Do not feed treated vines to dairy animals or animals

worm, Leaf Miner, Pepper Weevil—1 to 2 pints per acre  
harvest

Beetles—3 to 4 pints per acre. Do not use treated tops


ato Fruit Worm, Hornworms, Tomato Pin Worm, Cabbage  
Thrips—3 pints per acre. Do not apply

ported Cabbage Worm, Cutworms, Leaf Miners, Thrips—  
apply after seedling stage.

rawberry Weevil—2 to 3 pints per acre. Do not apply

ONLY: Cutworms, Clover Leaf Weevil, Armyworms—3 to  
only. Do not feed treated forage to livestock.

ankerworm, Fall Webworm, Tent Caterpillar—2 to 3 pints  
Bug—3 to 4 pints per 100 gals. water.



ACCEPTED  
59  
MA

# TOXAPHENE SPRAY 8-E

ACTIVE INGREDIENTS	By Weight
Toxaphene* .....	71.00%
Xylene .....	22.72%
INERT INGREDIENTS .....	6.28%
TOTAL .....	100.00%

(\*Technical Chlorinated Camphene Chlorine content 67% to 69%)

**WARNING:**  
**KEEP OUT OF REACH OF CHILDREN**  
**SEE RIGHT PANEL FOR REQUIRED**  
**WARNING STATEMENTS.**

**EPA REGISTRATION NO. 5967-95-ZA**  
**EPA EST. 5967-CA-1**

MANUFACTURED BY  
**MOYER CHEMICAL COMPANY**

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

## DIRECTIONS

Mix in water with agitation  
spraying. USE DESIGNATED  
UCT IN THE FOLLOWING  
PER ACRE UNLESS OTHER

**Vegetable and field crops**  
by ground, 5 to 10 gallons  
20 to 250 gallons by gro  
air. **Tree Fruits** — 40 to  
10 to 20 gallons by air.  
gallons by ground, 10 to  
**young fruit trees**, use a  
by ground per acre.

## USE PRECA

This product is toxic to the  
body of water. Birds, fish, etc.  
killed. Do not apply within 6  
from treated areas. Do not  
pour. Do not enter streams  
and spread of water. A  
on this label.

## DIRECTIONS FOR MIXING:

Mix in water with agitation; agitate solution while spraying. USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS OF WATER PER ACRE UNLESS OTHERWISE NOTED:

**Vegetable and field crops** — 20 to 250 gallons by ground, 5 to 10 gallons by air. **Small fruits** — 20 to 250 gallons by ground, 7 to 10 gallons by air. **Tree Fruits** — 40 to 800 gallons by ground, 10 to 20 gallons by air. **Citrus** — 100 to 3,000 gallons by ground, 10 to 20 gallons by air. **On young fruit trees**, use a minimum of 20 gallons by ground per acre.

## USE PRECAUTIONS:

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from treated areas. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not re-use container.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## WARNING

DO NOT TAKE INTERNALLY  
DO NOT INHALE  
DO NOT GET ON SKIN

**May Be Fatal If Swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing.** Wash thoroughly after using. Keep away from children and domestic animals. Do not store or transport near feed or food. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not use on young plants. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

— NON-RETURNABLE —

NET CONTENTS

GALLONS

