

ACCEPTED

6-25-74

5967-96

# MOYER

## TOXAPHENE 40 W

ACTIVE INGREDIENTS	BY WEIGHT
*Toxaphene	40.00%
Xylene	12.59%
INERT INGREDIENTS	47.41%
TOTAL	100.00%

\*Technical Chlorinated Camphene (Chlorine content 67%-69%)

### WARNING: KEEP OUT OF REACH OF CHILDREN

Read entire label. Use strictly in accordance with cautions and directions, and with applicable state and federal regulations.

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

Burn bag immediately when empty. Stay out of smoke. Do not re-use container.

May be fatal if swallowed. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. Do not breathe spray mist. Avoid contamination of feed and food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Warranty: This product is sold as is without warranty, expressed or implied concerning use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and/or handling is contrary to label instructions.

NON RETURNABLE —

EPA REGISTRATION NO. 5967-96 AA

EPA EST. 5967-CA.1

NET WEIGHT 5 LBS.

MANUFACTURED BY

### MOYER CHEMICAL COMPANY

BAY SHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

REGISTERED  
6-25-74  
UNDER THE  
FEDERAL  
INSECTICIDE  
ACT OF 1947  
5967-96

**TOXAPHENE 40 W**  
**THIS PRODUCT IS AN AGRICULTURAL INSECTICIDE**  
**FOR THE FOLLOWING USES:**

**BEANS (DRY, GREEN, LIMA):** Armyworms, Bean Beetles, Lygus Bugs, Serpentine Leaf Miners, Thrips—use 5 to 10 lbs. per acre in 30 to 200 gallons of water. Air application—use 10 to 20 gallons of water. Do not apply to Green or Snap Beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter. No limitations on shelled beans as human food.

**BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI:** Cabbage Looper, Corn Earworms, Cutworms, Diamond-Back Moth Larvae—use 5 to 10 lbs. per acre in 30 to 200 gallons of water. Flea Beetles—use 7½ lbs. per acre in 30 to 200 gallons of water. By air—use 8 lbs. in 10 to 20 gallons of water. Do not apply after edible parts begin to form or within 30 days of harvest.

**CABBAGE:** Cabbage Looper, Cutworms, Corn Earworms, Diamond-Back Moth Larvae, Vegetable Weevil—use 6 to 10 lbs. per acre in 30 to 200 gallons of water. Flea Beetle—use 7½ lbs. per acre in 30 to 200 gallons of water. By air—use 10 to 20 gallons of water. Do not apply after heads begin to form.

**CARROTS:** Cutworms, Lygus Bugs, (Seed crop only)—use 3 to 5 lbs. per acre in 30 to 200 gallons of water. Flea Beetle—use 7½ lbs. per acre in 30 to 200 gallons of water. By air—use 10 to 20 gallons of water. No time limitations. Do not feed treated tops to dairy animals or animals being finished for slaughter.

**CELERY:** Celery Leaf Tier, Cutworms, Vegetable Weevil—use 10 lbs. per acre in 30 to 200 gallons of water. By air—use 10 to 20 gallons of water. Do not apply after plants start to bunch or after plants are half mature.

**COLLARDS, KALE:** Armyworms, Cutworms, Imported Cabbage Worm—use 5 lbs. per acre in 30 to 200 gallons of water. By air—use 10 to 20 gallons of water. Do not apply within 28 days of harvest.

**Cabbage Looper, Flea Beetle, Salt Marsh Caterpillar—use 10 lbs. per acre in 30 to 200 gallons of water. By air—use 10 to 20 gallons of water. Do not apply within 35 days of harvest.**

**CORN:** Corn Earworm (in whorl), Chinch Bugs, Cutworms, Flea Beetles, Webworm—use 3 to 5 lbs. per acre in 30 to 200 gallons of water. Armyworms—use 6½ lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitations on use of grain.

**COTTON:** Armyworms, Boll Worms, Cutworms, Flea Beetles, Grasshoppers—use 7½ to 15 lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. Do not apply after bolls open when 10 lbs. or more are applied.

**EGGPLANT, PEPPERS:** Armyworms, Blister Beetles, Colorado Potato Beetles, Cutworms, Flea Beetles—use 5 to 7½ lbs. per acre in 30 to 100 gallons of water. By air—use 5 to 20 gallons of water. Do not apply within 5 days of harvest.

**LETTUCE:** Cabbage Looper, Cutworms, Diamond-Back Moth Larvae, Flea Beetles—use 7½ to 12 lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not apply after seedling stage on leaf lettuce. Do not apply after heads begin to form on head lettuce.

**LEEKs, ONIONS, SHALLOTS:** Cutworms, Thrips—use 7½ lbs. per acre for Shallots; 10 lbs. per acre for Leeks; 12½ lbs. per acre for Onions—in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not apply to green or spring Onions. Do not apply after seedling stage for Leeks and Shallots.

**PARSNIPS:** Cutworms, Flea Beetles—use 7 to 12½ lbs. in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not use tops for food or feed.

**PEARS:** Pear Psylla—use 2½ to 3 lbs. per 100 gallons of water; apply 400 to 600 gallons of water per acre. Do not apply within 40 days of harvest, or after second cover spray.

**PEAS:** Cutworms, Lygus Bugs—use 7½ to 12½ lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not apply after pods begin to form if pods are to be used as food; OR: within 7 days of harvest if pods are not to be used as food. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**RUTABAGAS:** Cutworms, Flea Beetles—use 5 to 10 lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not use treated tops for food or feed.

**SPINACH:** Armyworms, Cutworms, Imported Cabbage Worm, Leaf Miners, Thrips—use 5 to 10 lbs. per acre in 30 to 200 gallons of water. By air—use 5 to 20 gallons of water. Do not apply more than 5 lbs. after seedling stage. If using 5 lbs. or less per acre — do not apply within 21 days of harvest. Do not apply more than once per season.

**STRAWBERRIES:** Cutworms, Strawberry Weevil—use 5 to 7½ lbs. per acre in 75 to 400 gallons of water. By air—use 10 to 20 gallons of water. Do not apply after fruit begin to form.

**TOMATOES:** Armyworms, Cabbage Looper, Hornworms, Russet Mites, Serpentine Leaf Miners, Tomato Fruit Worms, Tomato Pinworms—use 5 to 12½ lbs. per acre in 30 to 200 gallons of water. Do not apply more than 5 lbs. within 1 (one) day of harvest; Do not apply more than 12½ lbs. within 3 days of harvest.