

BEN SUL SPRAY

DIRECTIONS FOR USE:

THIS PRODUCT IS AN AGRICULTURAL FUNGICIDE FOR THE BELOW INDICATED USES:

ALMONDS: Scab, Brown Rot—6 lbs. per 100 gals. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

APPLES, PEARS: Scab, Powdery Mildew—5 lbs. or 4 lbs. plus 1 gal. liquid Lime-Sulfur per 100 gals. Apply as a full coverage spray at pre-bloom, calyx and early cover spray. Under extreme disease conditions, 7 to 8 lbs. may be needed.

GRAPES: Powdery Mildew—5 lbs. per 100 gals. Apply as needed to maintain control. Always follow local recommendations for best results.

NECTARINES, PEACHES: Powdery Mildew, Brown Rot—5 lbs. per 100 gals. Apply at pink bud for Brown Rot control. For Mildew, apply at petal fall and as needed to maintain control until pit hardening. Check local recommendations.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines): For Citrus Flat Mite, Citrus Rust Mite, Citrus Thrips, Yuma Spider Mites—Apply at least 60 lbs. in 400 to 1500 gals. of water. Apply when pests first appear and repeat as necessary. Do not apply during or preceding high temperatures. Do not apply within two months of an oil application.

STRAWBERRIES: For Mildew, use 5 to 10 lbs. per acre. By Air: Apply in 10 to 20 gals. of water. By Ground: Apply in 200 to 500 gals. of water.

STRAWBERRIES are sensitive to sulfur, especially during periods of high temperature. Do not apply during hot weather, or when it is anticipated that hot weather will follow application. It is suggested that you contact your local Cooperative Agricultural Extension Service for further advice.

NET WEIGHT 50 LBS.


(CONTAINS BENTONITE)

ACTIVE INGREDIENTS

Sulfur By Weight 81.25%

INERT INGREDIENTS 18.75%

TOTAL 100.00%



WARNING:

Keep Out of Reach of Children

EPA REG. NO. 5967-101 AA

EPA EST. 5967-CA-1

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY

P.O. BOX 945

SAN JOSE, CALIFORNIA

IMPORTANT

This product is a mixture of Bentonite and Sulfur in the correct proportions to ensure the best spreading and sticking qualities.

Do not permit workers to enter the treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes 1, Peaches and Nectarines 1, Citrus 1.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

WARNING

Certain crops and other plants are susceptible to injury from Sulfur. Do not use Sulfur on any crop unless local conditions have proven its use to be safe. Do not allow this material to drift onto neighboring crops. Avoid contamination of feed and foodstuffs. Do not use during or immediately prior to periods of extreme high temperatures (90° F or higher), as foliage damage may result. Do not apply within 3 weeks of an oil spray or injury may result. Use only in accordance with local recommendations.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS OF WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION — Deciduous Fruit Crops, Grapes and Strawberries—60 to 800 gals, Citrus—400 to 3,000 gals. (On young fruit planting — Use minimum of 15 gals. water per acre.) Where recommended for AIR APPLICATION—Strawberries—10 to 20 gals. Orchard and Vineyard Crops—10 to 20 gals.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. — NON-RETURNABLE —