

300
17

JUL 17 1985

Siemer & Associates, Inc.
4672 W. Jennifer
Suite 103
Fresno, CA 93711

Attention: S. R. Siemer

Gentlemen:

Subject: Cu Zn
EPA Registration No. 5967-111
Application Submission Dated April 2, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted at the next printing.

- 1. Delete "or wetlands" from the environmental hazards section. It is no longer required.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

BEST AVAILABLE COPY

Harry M. Jacoby
Harry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (YS-767)

Enclosure

RD/PHB:JOB:88667;Notes:RD-39;Kendrick:898-1270;7/11/85;Del:7/23/85;Kim

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

REENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled.

Mix in water with agitation; agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted: Vegetable and Field Crops — 20 to 250 gallons by ground, 5 to 10 gallons by air. Small Fruits — 20 to 250 gallons by ground, 7 to 10 gallons by air. Tree Fruits — 40 to 800 gallons by ground, 10 to 20 gallons by air. Citrus — 100 to 3,000 gallons by ground, 10 to 20 gallons by air. On young fruit trees, use a minimum of 20 gallons by ground per acre.

STORAGE AND DISPOSAL

STORAGE: Keep product dry in closed container. Do not use empty container for any other purpose.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food or feed by storage or disposal.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 17 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

5967-111

ACTIVE INGREDIENT

Copper (in Basic Copper Sulfate)
INERT INGREDIENTS (Including

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

See Additional precautions on
PRACTICAL TREATMENT

IF SWALLOWED: Administer first aid by sticking finger down throat. Get professional medical assistance.

IF ON SKIN: Rinse off material with water. Get medical attention if irritation persists.

IF IN EYES: Wash with plenty of water. Get medical attention.

IF INHALED: Remove to fresh air. Get medical attention if respiration is affected. Never give anything by mouth to an unconscious person.

EPA REG. NO. 05967-111 AA

PR

AL USUARIO: Si usted usa este producto hasta que se agote completamente.

TO THE USER: If you use this product until it is completely explained to you.)

MANU

MOYER CHEI

P.O. BOX 3434, F

NET W

MOYER

CHEMICALS

Zn

By Weight

(as) expressed as metallic	19.0%
g 19% Zinc as elemental)	81.0%
TOTAL	100.0%

REACH OF CHILDREN

CAUTION

DO NOT SWALLOW OR INHALE

precautions on side panel

TREATMENT: (First Aid)

water freely and induce vomiting by it. Repeat until vomit fluid is clear. Pro-
nce should be secured immediately.

rial and wash with soap and water. Get
tion persists.

ty of water for at least 15 minutes. Get

h air. IF NOT BREATHING; give artificial
outh to mouth. Get medical attention.
outh to an unconscious person.

VA EPA EST. 35296-CA-2

PRECAUCIÓN

ted no lee ingles, no use este
la etiqueta le haya sido ex-
3.

you cannot read English, do not
until the label has been fully

UFACTURED FOR

CHEMICAL COMPANY

FRESNO, CALIFORNIA 93755

4-85

WEIGHT 8 LBS.

ALMONDS: For control of Shothole blight — Use 13.95 lbs. per 100 gals. of water. Use in dormant, petal fall and jacket sprays.

APPLES: For control of Fireblight — Use 1. to 1.5 lbs. per 100 gals. of water. Begin at 10% bloom. Repeat at 5-day intervals until late bloom is over.

APRICOTS: For control of Brown rot — Use 11 to 13 lbs. per 100 gals. of water. Apply in red bud to pre-jacket fall stages. For control of Shothole — Use 14 to 22 lbs. per 100 gals. of water. Apply as a fall or winter dormant spray, applying before the first rains. Repeat in red bud to popcorn stages.

CITRUS: (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos): For control of Brown rot — Use 1.32 to 4 lbs. plus 3 lbs. Zinc Sulfate and 4 lbs. hydrated lime in 100 gals. of water. In California, for groves to be fumigated or where copper injury is a problem, use the lower rate. Where the groves are not to be fumigated or where copper injury is not a problem, use the higher rate. Apply at first indication of fall rains, or at first sign of disease.

PEACHES, NECTARINES: For control of Leaf curl and shot-hole — Use 11 to 16 lbs. per 100 gals. of water; or in California, use 34 to 44 lbs. per acre in sufficient water for good coverage; 25 to 50 lbs. per 20 gals. of water per acre by air — apply in dormant season.

PEARS: For control of Fireblight — Use 1.4 lbs. per 100 gals. of water. Apply 10% to 20% bloom, full bloom and at 5- to 7-day intervals during bloom. Do not use on Comice, Anjou or Seckel varieties.

STRAWBERRIES: Downy mildew, leaf blight, leaf scorch, leaf spot — Use 6.75 lbs. per 100 gals. of water. Begin after leaves form. Repeat at 10 to 14-day intervals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: For control of Downy mildew and Leaf spots — Use 6 to 8 lbs. per 100 gals. of water. Begin when plants are above ground in plant bed or before diseases appear. Repeat at 7- to 10-day intervals in plant bed and field.

CELERY: For control of early blight, late blight, and leaf spots — Use 6 to 13 lbs. per 100 gals. of water. Begin in plant bed or field when plants are 6 inches high or when diseases first appear. Repeat at 5- to 14-day intervals as necessary.

LETTUCE: For control of Downy mildew — Use 6 to 8 lbs. per 100 gals. of water. Begin before disease appears. Repeat at 7- to 10-day intervals.

ONIONS: For control of Downy mildew — Use 6 to 8 lbs. per 100 gals. of water. Begin when plants are 4 to 6 inches high. Repeat at 7-day intervals.

POTATOES: For control of early and late blight — Use 6 to 13 lbs. per 100 gals. of water. Begin when plants emerge, or when they are 4 to 6 inches high; or when diseases appear. Repeat at 7- to 10-day intervals or more frequently under severe late blight conditions (rainy or foggy periods).

TOMATOES: For control of Anthracnose, bacterial spot, early or late blight, or septoria — Use 6 to 13 lbs. per 100 gals. of water. Begin in plant bed or field before diseases appear. Repeat at 4- to 14-day intervals. For ground use only.

CONDITIONS OF SALE:

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product.

2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

BEST AVAILABLE COPY