

BEST AVAILABLE COPY

MOYER

PRODUCTS

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION

Caution causes eye irritation. Harmful if swallowed. May cause drowsiness and dizziness in some individuals. Avoid contact with bare skin. Avoid contact with clothing. Avoid breathing dust. Avoid contact with food. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wastes.

Cu Zn

ACTIVE INGREDIENTS	By Weight
Copper (as Basic Copper Sulfate)	19.0
Expressed as metallic	
INERT INGREDIENTS (including	81.0
10% Zinc as elemental)	
TOTAL	100.0

**KEEP OUT OF REACH OF
CHILDREN**

CAUTION

**HARMFUL IF SWALLOWED.
DO NOT SWALLOW OR INHALE**

See additional precautions on side panel

PRACTICAL TREATMENT (First Aid)

IF IN EYES: Flush with plenty of water. Call a physician.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove to fresh air. IF NOT BREATHING, give artificial respiration, preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container. Keep container tightly closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: In compliance with local, state, and federal regulations, this container must be disposed of in a sanitary and safe manner as required by state and local authorities. Do not reuse or refill if banned, stay out of stream.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until fumes have dried. Certain states may require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Warning or oral warnings must be given to workers who are expected to be in treated areas. In an area about to be treated with this product, when oral warnings are given, warnings shall be given in all language customarily understood by workers. Written or oral warnings must include the following information: CAUTION. Area treated with Copper Basic Copper Sulfate on (date of application). Do not enter without appropriate protective clothing and aprons have dried. Required Protective Clothing: long sleeved shirt and long legged pants.

MIXING DIRECTIONS

Mix in water with agitation, apply with sprayer. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

Vegetable and Field Crops — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

Small Fruits — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

Tree Fruit — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

Citrus — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

On young fruit trees — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

ALMONDS — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

POTATOES — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

PEACHES, NECTARINES — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

PLUMS — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

APRICOTS — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

CHERRY — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

PEAR — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

CITRUS — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

ORANGE — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

LEMON — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

LIME — 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre. Use 20 gal. of water per acre.

It is not a problem to use the higher rate. Apply at first indication of fall rains or at first sign of disease.

PEACHES, NECTARINES — For control of LEAF CURL and SHOTHOLE — Use 10 to 16 lbs. per 100 gal. of water. In California use 34 to 44 lbs. per 100 gal. of water for good coverage. 25 to 50 lbs. per 20 gal. of water per acre by air — apply in dormant season.

PLUMS — For control of FIREBLIGHT — Use 4 to 6 lbs. per acre in sufficient water for good coverage. Use 3 to 4 lbs. per acre in 20 gal. water by air. Apply 100 to 200 bloom, full bloom and 3 to 7 day intervals during bloom. Do not use on (certain) (certain) varieties.

STRAWBERRIES — For control of LEAF SPOT — Use 4 to 6 lbs. per acre. Begin after leaves form. Repeat at 10 to 15 day intervals.

BRIGHT LEAF SCORCH, LEAF SPOT — Use 4 to 6 lbs. per acre. Begin after leaves form. Repeat at 10 to 15 day intervals.

BRUSSELS SPROUTS, CABBAGE — For control of DOWNY MILDEW and LEAF SPOTS — Use 6 to 8 lbs. per acre. Begin when plants are above ground in plant bed or before diseases appear. Repeat at 7 to 10 day intervals in plant bed and field.

CELERY — For control of EARLY BLIGHT, LATE BLIGHT, and LEAF SPOTS — Use 6 to 10 lbs. per acre. Begin in plant bed or field when plants are 6 inches high or when diseases first appear. Repeat at 5 to 10 day intervals as necessary.

LETTUCE — For control of DOWNY MILDEW — Use 4 to 6 lbs. per acre. Begin before disease appears. Repeat at 7 to 10 day intervals.

ONIONS — For control of DOWNY MILDEW — Use 4 to 6 lbs. per acre. Begin when plants are 6 to 8 inches high. Repeat at 7 day intervals.

POTATOES — For control of EARLY and LATE BLIGHT — Use 6 to 10 lbs. per acre. Begin when plants are 6 inches high or when they are 4 to 6 inches high. When diseases appear. Repeat at 7 to 10-day intervals as necessary. Under severe late blight and/or severe early blight, repeat more frequently.

TOMATOES — For control of ANTHRACNOSE, BACTERIAL SPOT, EARLY or LATE BLIGHT or SEPTORIA — Use 6 to 10 lbs. per acre. Begin in plant bed or field before diseases appear. Repeat at 4 to 14 day intervals. For ground use only.

CONDITIONS OF SALE

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such a authorization must be in writing) to make any other warranty, guarantee or promise concerning this product.

2. Because the time, place, rate of application, weather conditions and normal natural conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price in no event shall Moyer be liable for indirect or consequential damages.

NET WEIGHT 16 LBS.

MANUFACTURED FOR
MOYER PRODUCTS, INC.
P.O. BOX 5434
FRESNO, CALIFORNIA 93755