

BEST AVAILABLE COPY**MOYER****PRODUCTS****PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Caution: Causes eye irritation. Harmful if absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with the skin, eyes, or clothing. Avoid breathing dust. Protective clothing, including goggles, should be worn. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wastes.

Cu Zn

ACTIVE INGREDIENTS	By Weight
Copper (in Basic Copper Sulfate)	19.0%
expressed as metallic	
INERT INGREDIENTS (including	
19% Zinc as elemental)	81.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE

See additional precautions on side panel

PRACTICAL TREATMENT. (First Aid):

IF IN EYES: Flush with plenty of water. Call a physician.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove to fresh air. If NOT SPEAKING, give artificial respiration, preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry, well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NET WEIGHT 16 LBS.

MANUFACTURED FOR
MOYER PRODUCTS, INC.
P.O. BOX 5434
FRESNO, CALIFORNIA 93755

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Certain states may require more restrictive re-entry intervals for various crops treated with this product; consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information: CAUTION: Area treated with Copper (in Basic Copper Sulfate) on (date of application). Do not enter without appropriate protective clothing until sprays have dried. Required Protective Clothing: long sleeved shirt and long legged pants.

MIXING DIRECTIONS

Mix in water with agitation. Agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted.

Vegetable and Field Crops — 20 to 250 gallons by ground; 5 to 20 gallons by air.

Small Fruits — 20 to 250 gallons by ground; 7 to 10 gallons by air.

Tree Fruits 40 to 800 gallons by ground; 20 gallons by air.

Citrus 100 to 1,500 gallons by ground; 20 gallons by air.

On young fruit trees use a minimum of 20 gallons by ground per acre.

ALMONDS: For control of SHOTHOLE BLIGHT — Use 14 lbs. per 100 gals. of water. Use in dormant petal fall and jacket sprays.

APPLES: For control of FIREBLIGHT

Use 4 to 6 lbs. per acre in sufficient water for good coverage. Use 4 to 6 lbs. per acre in 20 gal. of water by air. Begin at 10% bloom. Repeat at 3 day intervals until late bloom is over.

APRICOTS: For control of BROWN ROT

Use 33 to 52 lbs. per acre in sufficient water for good coverage. Use 25 to 39 lbs. per acre in 20 gal. by air. Apply in bud to pre-jacket fall stages. For control of SHOTHOLE — Use 42 to 88 lbs. per acre in sufficient water for good coverage. Use 32 to 66 lbs. per acre in 20 gal. by air. Apply as a fall or winter dormant spray, applying before the first rains. Repeat in bud to popcorn stages.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos) For control of BROWN ROT — Use 13 to 4 lbs. plus 3 lbs. Zinc Sulfate and 4 lbs. hydrated lime in 100 gals. of water. In California, for groves to be fumigated or where copper injury is a problem, use the lower rate. Where the groves are not to be fumigated or where copper

injury is not a problem, use the higher rate. Apply at first indication of fall rains, or at first sign of disease.

PEACHES, NECTARINES: For control of LEAF CURL and SHOTHOLE — Use 11 to 16 lbs. per 100 gals. of water, or in California, use 34 to 44 lbs. per acre in sufficient water for good coverage. 25 to 50 lbs. per 20 gals. of water per acre by air — apply in dormant season.

PEARS: For control of FIREBLIGHT

Use 4 to 9 lbs. per acre in sufficient water for good coverage. Use 3 to 6 lbs. per acre in 20 gal. water by air. Apply 10% to 20% bloom, full bloom and 5 to 7 day intervals during bloom. Do not use on Comice, Anjou or Seckel varieties.

STRAWBERRIES: DOWNY MILDEW, LEAF

BLIGHT, LEAF SCORCH, LEAF SPOT — Use 4 to 6 lbs. per acre. Begin after leaves form. Repeat at 10 to 14 day intervals.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE

CALIFLOWERS: For control of DOWNY MILDEW

and LEAF SPOTS — Use 6 to 8 lbs. per acre. Begin when plants are above ground in plant bed or before seedlings appear. Repeat at 7 to 10 day intervals in plant bed and field.

CELERY: For control of EARLY BLIGHT, LATE BLIGHT, and LEAF SPOTS — Use 6 to 13 lbs. per acre. Begin in plant bed or field when plants are 6 inches high or when disease first appears. Repeat at 5 to 10 day intervals as necessary.

LETTUCE: For control of DOWNY MILDEW — Use 4 to 8 lbs. per acre. Begin before disease appears. Repeat at 7 to 10 day intervals.

ONIONS: For control of DOWNY MILDEW — Use 4 to 8 lbs. per acre. Begin when plants are 4 to 6 inches high. Repeat at 7 day intervals.

POTATOES: For control of EARLY and LATE BLIGHT — Use 6 to 13 lbs. per acre. Begin when plants emerge or when they are 4 to 6 inches high, or when disease appears. Repeat at 7 to 10-day intervals or more frequently under severe late blight conditions (rainy or foggy periods).

TOMATOES: For control of ANTHRACNOSE, BACTERIAL SPOT, EARLY or LATE BLIGHT, or SEPTORIA — Use 6 to 13 lbs. per acre.

Begin in plant bed or field before diseases appear. Repeat at 4 to 14 day intervals. For ground use only.

CONDITIONS OF SALE

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product.

2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

12-89