

Reg # 5967-111

PM - 22

File  
178



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

FEB 23 1994

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Terri Aal  
MOYER PRODUCTS, INC.  
P. O. Box 5434  
Fresno, CA 93755

Subject: Label Amendment Submission of 10/11/93 in Response to PR Notice 93-7  
EPA Reg. No. 5967-111  
MOYER CU ZN

**IMPORTANT NOTICE:** This Letter Requires You to Submit a Notification to the Office of Compliance Monitoring within 20 Days.

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

**SERIOUS ERROR(S) ON SUBMITTED LABELING:**

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

**YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING.** Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

**YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.**



Recycled/Recyclable  
Printed with Soy/Plant-based Ink  
contains at least 50% recycled content

Each notification must include the following information:

1. Registrant's name and address.
2. EPA Registration number of the product for which the labeling was submitted.
3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

1. Quantity of the product that has been sold or distributed.
2. EPA Producing Establishment number.
3. Location of the product, if known.
4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director  
Compliance Division, (7203)  
Worker Protection Notification  
Office of Compliance Monitoring  
U.S. Environmental Protection Agency  
401 M Street SW  
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

**WHAT THIS ACCEPTANCE MEANS:**

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

**WHAT YOU NEED TO DO NEXT:**

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling as amended

**AND**

- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (7505C)  
U.S. Environmental Protection Agency  
401 M Street, SW  
Washington, D.C. 20460-0001


4 3 8

Page 4

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Sincerely,

  
Jim Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

598

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division

TERRI AAL  
MOYER PRODUCTS, INC.  
BOX 5434  
FRESNO CA 93755

Comment for: EPA Reg Nr.5967-111  
MOYER CU ZN

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

The Agricultural Use Requirements section must be located in a clearly separate box on the product labeling -- with lines or other graphic indicators (such as contrasting color) to separate the section from the surrounding text.

The following statement on your proposed label is not required for this product: "Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas." Delete it from the Agricultural Use Requirements box.

The eyewear requirement for early entry in the Agricultural Use Requirements Box on your proposed labeling is missing. Add "Protective eyewear". The exact location is shown in Sections F and G on Part 2 of the Product Worksheet in Supplement Three-A to PR Notice 93-7.

The 72 hour REI exception for arid areas is not needed on this label.

# MOYER PRODUCTS

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Causes eye irritation. Harmful if absorbed through the skin or inhaled. May cause skin sensitization reactions in certain individuals. Avoid contact with the skin, eyes, or clothing. Avoid breathing dust.

#### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear.

Follow manufacturer's instructions for cleaning and maintaining personal protective equipment (PPE). If no such instructions for washable, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product contains clay which contains crystalline silica (0.5%) a naturally occurring component of sand and inorganic soils. Crystalline silica has been classified as a 2A carcinogen by the International Agency for Research on Cancer (IARC).

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or County, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and to the extent the various Federal, State, and local laws, rules, and regulations regarding the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, certification, notification, and emergency assistance. It also contains specific instructions and cautions pertaining to the use of this product in greenhouses, in confined areas, and in structures used for housing livestock. The requirements in this label only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval. REI of 12 hours.

Do not enter or allow worker entry into treated areas that are under the restricted entry interval. REI of 12 hours.

Do not enter or allow worker entry into treated areas that are under the restricted entry interval. REI of 12 hours.

Do not enter or allow worker entry into treated areas that are under the restricted entry interval. REI of 12 hours.

#### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a cool, dry, well-ventilated area.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag in to application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## Cu Zn

#### ACTIVE INGREDIENTS

Copper (in Basic Copper Sulfate)  
expressed as metallic

By Weight

19.0%

INERT INGREDIENTS (including  
19% Zinc as elemental)

81.0%

TOTAL

100.0%

## KEEP OUT OF REACH OF CHILDREN CAUTION

HARMFUL IF SWALLOWED.  
DO NOT SWALLOW OR INHALE

See additional precautions on side panel

#### PRACTICAL TREATMENT: (First Aid):

**IF IN EYES:** Flush with plenty of water. Call a physician.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove to fresh air. IF NOT BREATHING, give artificial respiration, preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

EPA Reg. No. 5967-111AA

EPA Est. No. 2925-CA 1

5.92

NET WEIGHT 16 LBS.

MANUFACTURED FOR  
**MOYER PRODUCTS, INC.**  
P.O. BOX 5434  
FRESNO, CALIFORNIA 93755

#### MIXING DIRECTIONS

Mix in water with agitation, agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted.

**Vegetable and Field Crops** — 20 to 250 gallons by ground, 5 to 20 gallons by air.

**Small Fruits** — 20 to 250 gallons by ground, 7 to 10 gallons by air.

**Tree Fruits** — 40 to 800 gallons by ground, 20 gallons by air.

**Citrus** — 100 to 1,500 gallons by ground, 20 gallons by air. Or young fruit trees use a minimum of 20 gallons by ground per acre.

**ALMONDS:** For control of SHOTHOLE BLIGHT — Use 14 lbs. per 100 gals. of water. Use in dormant, petal fall and jacket sprays.

**APPLES:** For control of FIREBLIGHT — Use 4 to 6 lbs. per acre in sufficient water for good coverage. Use 4.5 lbs. per acre in 20 gal. of water by air. Begin at 10% bloom. Repeat at 5-day intervals until late bloom is over.

**APRICOTS:** For control of BROWN ROT BLOSSOM BLIGHT — Use 33 to 52 lbs. per acre in sufficient water for good coverage. Use 25 to 39 lbs. per acre in 20 gal. by air. Apply in red bud to pre-jacket fall stages. For control of SHOTHOLE — Use 42 to 66 lbs. per acre in sufficient water for good coverage. Use 32 to 66 lbs. per acre in 20 gal. by air. Apply as a fall or winter dormant spray, applying before the first rains. Repeat in red bud to petal fall stages.

**CITRUS:** (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos). For control of BROWN ROT — Use 13 to 4 lbs. plus 3 lbs. Zinc Sulfate and 4 lbs. hydrated lime in 100 gals. of water. In California, for groves to be fumigated or where copper injury is a problem, use the lower rate. Where the groves are not to be fumigated or where copper injury is not a problem, use the higher rate. Apply at first indication of fall rains, or at first sign of disease.

**PEACHES, NECTARINES:** For control of LEAF CURL and SHOTHOLE — Use 11 to 16 lbs. per 100 gals. of water or in California, use 34 to 44 lbs. per acre in sufficient water for good coverage. 25 to 50 lbs. per 20 gals. of water per acre by air — apply in dormant season.

**PEARS:** For control of FIREBLIGHT — Use 4 1/2 to 5 1/2 lbs. per acre in sufficient water for good coverage. Use 3 to 4 lbs. per acre in 20 gal. water by air. Apply 10% to 20% bloom. Full bloom and 5- to 7-day intervals during bloom. Do not use on Comice, Anjou or Seckel varieties.

**STRAWBERRIES:** DOWNY MILDEW, LEAF BLIGHT, LEAF SCORCH, LEAF SPOT — Use 4 to 6 1/2 lbs. per acre. Begin after leaves form. Repeat at 10- to 14-day intervals.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** For control of DOWNY MILDEW and LEAF SPOTS — Use 6 to 8 lbs. per acre. Begin when plants are above ground in plant bed or before diseases appear. Repeat at 7- to 10-day intervals in plant bed and field.

**CELERY:** For control of EARLY BLIGHT, LATE BLIGHT and LEAF SPOTS — Use 6 to 12 lbs. per acre. Begin in plant bed or field when plants are 6 inches high or when diseases first appear. Repeat at 5- to 14-day intervals as necessary.

**LETTUCE:** For control of DOWNY MILDEW — Use 4 to 6 lbs. per acre. Begin before disease appears. Repeat at 7- to 10-day intervals.

**ONIONS:** For control of DOWNY MILDEW — Use 6 to 8 lbs. per acre. Begin when plants are 4 to 6 inches high. Repeat at 7-day intervals.

**POTATOES:** For control of EARLY BLIGHT, LATE BLIGHT, and SCAB — Use 6 to 13 lbs. per acre. Begin when plants are 4 to 6 inches high. Repeat at 7- to 10-day intervals. For control of SCAB — Use 6 to 13 lbs. per acre. Begin when plants are 4 to 6 inches high. Repeat at 7- to 10-day intervals.

**TOMATOES:** For control of SPOT, EARLY or LATE BLIGHT — Use 6 to 13 lbs. per acre. Begin in plant bed or field when plants are 6 inches high or when diseases first appear. Repeat at 7- to 10-day intervals.

**FOR THE FOLLOWING CROPS:** Use 4 lbs. per 100 gallons of water for control of disease and repeat at 7- to 10-day intervals.

CROP	DISEASE
Auraria	Xanthom
Alfalfa	Alternaria
Asparagus	Cercospora
Begonia	Xanthom
Bulbs (Easter Lily, Tulip, Gladiolus)	Anthracnose
Carnation	Alternaria
Chrysanthemum	Septoria
Cotoneaster	Botrytis
Euonymus	Botrytis
Ivy	Xanthom
Phlox	Botrytis
Pyracantha	Fire
Rose	Powdery
Yucca (Adam's Needle)	Cercospora

1. Moyer warrants that the products specified and used on this label when used under normal conditions of use will be effective in controlling the diseases and pests named. Because the time of application and the amount of product applied are beyond Moyer's control, replacement of product or refund of money shall not be made.

2. Moyer warrants that the products specified and used on this label when used under normal conditions of use will be effective in controlling the diseases and pests named. Because the time of application and the amount of product applied are beyond Moyer's control, replacement of product or refund of money shall not be made.

BEST AVAILABLE COPY

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear.

Follow manufacturer's instructions for cleaning and maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product contains clay which contains crystalline silica (0.5%). a naturally occurring component of sand and inorganic soils. Crystalline silica has been classified as a 2A carcinogen by the International Agency for Research on cancer (IARC).

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 14 hours.

The REI is 12 hours in outdoor areas where average rainfall is less than 25 inches a year.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gaiters, shoes plus socks.

#### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry, well-ventilated area.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag in to application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACTIVE INGREDIENTS	By Weight
Copper (in Basic Copper Sulfate) expressed as metallic	19.0%
INERT INGREDIENTS (including 19% Zinc as elemental)	81.0%
TOTAL	100.0%

### KEEP OUT OF REACH OF CHILDREN

## CAUTION

HARMFUL IF SWALLOWED.  
DO NOT SWALLOW OR INHALE

See additional precautions on side panel

**PRACTICAL TREATMENT: (First Aid):**

**IF IN EYES:** Flush with plenty of water. Call a physician.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove to fresh air. IF NOT BREATHING, give artificial respiration, preferably mouth to mouth. Get medical attention. Never give anything by mouth to an unconscious person.

EPA Reg No 5967-111AA

EPA Est No 2935-CA-1

5/92

**NET WEIGHT 16 LBS.**

MANUFACTURED FOR  
**MOYER PRODUCTS, INC.**  
P.O. BOX 5434  
FRESNO, CALIFORNIA 93755

**BEST AVAILABLE COPY**

### MIXING DIRECTIONS

Mix in water with agitation, agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted:

**Vegetable and Field Crops** — 20 to 250 gallons by ground, 5 to 20 gallons by air.

**Small Fruits** — 20 to 250 gallons by ground, 7 to 10 gallons by air.

**Tree Fruits** — 40 to 800 gallons by ground, 20 gallons by air.

**Citrus** — 100 to 1500 gallons by ground, 20 gallons by air. On young fruit trees, use a minimum of 20 gallons by ground per acre.

**ALMONDS:** For control of SHOTHOLE BLIGHT — Use 14 lbs. per 100 gals. of water. Use in dormant, petal fall and jacket sprays.

**APPLES:** For control of FIREBLIGHT — Use 4 to 6 lbs. per acre in sufficient water for good coverage. Use 4.5 lbs. per acre in 20 gal. of water by air. Begin at 10% bloom. Repeat at 5-day intervals until late bloom is over.

**APRICOTS:** For control of BROWN ROT BLOSSOM BLIGHT — Use 33 to 52 lbs. per acre in sufficient water for good coverage. Use 25 to 39 lbs. per acre in 20 gal. by air. Apply in red bud to pre-jacket fall stages. For control of SHOTHOLE — Use 42 to 88 lbs. per acre in sufficient water for good coverage. Use 32 to 66 lbs. per acre in 20 gal. by air. Apply as a fall or winter dormant spray, applying before the first rains. Repeat in red bud to popcorn stages.

**CITRUS:** (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos) For control of BROWN ROT — Use 13 to 4 lbs. plus 3 lbs. Zinc Sulfate and 4 lbs. hydrated lime in 100 gals. of water in California, for groves to be fumigated or where copper injury is a problem, use the lower rate. Where the groves are not to be fumigated or where copper injury is not a problem, use the higher rate. Apply at first indication of fall rains, or at first sign of disease.

**PEACHES, NECTARINES:** For control of LEAF CURL and SHOTHOLE — Use 11 to 16 lbs. per 100 gals. of water, or in California, use 34 to 44 lbs. per acre in sufficient water for good coverage, 25 to 50 lbs. per 20 gals. of water per acre by air — apply in dormant season.

**PEARS:** For control of FIREBLIGHT — Use 4 1/2 to 5 1/2 lbs. per acre in sufficient water for good coverage. Use 3 to 4 lbs. per acre in 20 gal. water by air. Apply 10% to 20% bloom, full bloom and 5- to 7-day intervals during bloom. Do not use on Comice, Anjou or Seckel varieties.

**STRAWBERRIES:** DOWNY MILDEW, LEAF BLIGHT, LEAF SCORCH, LEAF SPOT — Use 4 to 6 1/4 lbs. per acre. Begin after leaves form. Repeat at 10- to 14-day intervals.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** For control of DOWNY MILDEW and LEAF SPOTS — Use 6 to 8 lbs. per acre. Begin when plants are above ground in plant bed or before diseases appear. Repeat at 7- to 10-day intervals in plant bed and field.

**CELERY:** For control of EARLY BLIGHT, LATE BLIGHT, and LEAF SPOTS — Use 6 to 13 lbs. per acre. Begin in plant bed or field when plants are 6 inches high or when diseases first appear. Repeat at 5- to 14-day intervals as necessary.

**LETTUCE:** For control of DOWNY MILDEW — Use 4 to 8 lbs. per acre. Begin before disease appears. Repeat at 7- to 10-day intervals.

**ONIONS:** For control of DOWNY MILDEW — Use 6 to 8 lbs. per acre. Begin when plants are 4 to 6 inches high. Repeat at 7-day intervals.

**POTATOES:** For control of EARLY and LATE BLIGHT — Use 6 to 13 lbs. per acre. Begin when plants emerge or when they are 4 to 6 inches high or when diseases appear. Repeat at 7- to 10-day intervals or more frequently under severe late blight conditions (rainy or foggy periods).

**TOMATOES:** For control of ANTHRACNOSE, BACTERIAL SPOT, EARLY or LATE BLIGHT, or SEPTORIA — Use 6 to 13 lbs. per acre. Begin in plant bed or field before diseases appear. Repeat at 4- to 14-day intervals. For ground use only.

### ORNAMENTALS

**SYCAMORE:** Anthracnose — Apply as a full cover spray using 5 1/2 to 7 1/2 lbs. per 100 gallons of water. Make first application at bud break and second application 7 to 10 days later at 10% leaf expansion.

**FOR THE FOLLOWING ORNAMENTALS:** Use 3 to 4 lbs. per 100 gallons of water. Begin application at first sign of disease and repeat at 7- to 14-day intervals as needed.

CROP	DISEASES
Auraria	Xanthomonas & Cercospora leafspots, Alternaria
Azalea*	Cercospora leafspot, Botrytis blight, Phytophthora dieback and Powdery mildew
Begonia	Xanthomonas leafspot
Bulbs (Easter Lily, Tulip, Gladiolus)	Anthrachnose, Botrytis blight
Carnation*	Alternaria blight, Pseudomonas leafspot, and Botrytis blight
Chrysanthemum*	Septoria leafspot and Botrytis blight
Coroneaster	Botrytis blight
Euconymus	Botrytis blight, Anthracnose
Ivy*	Xanthomonas leafspot
Pachysandra	Botrytis leaf blight
Pennwinkle	Phomopsis stem blight
Philodendron	Bacterial leafspot
Pyracantha	Fireblight, Scab
Rose*	Powdery mildew, Black spot
Yucca (Adam's needle)	Cercospora and Septoria leafspot

\*Discoloration of foliage and/or blooms have been noted on some varieties. To prevent residues on commercial plants, do not spray just before the selling season.

NOTE: Phytotoxicity may depend on varietal differences. If unfamiliar with the use of CuZn, apply the recommended rate to a few plants and observe after 7 to 14 days for symptoms of phytotoxicity. With specific plants not listed above we recommend a preliminary trial to determine plant tolerance.

### CONDITIONS OF SALE:

1. Moyer warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with the directions under normal conditions of use. No one other than an officer of Moyer is authorized (and such authorization must be in writing) to make any other warranty, guarantee or direction concerning this product.
2. Because the time, place, rate of application, weather conditions and normal or abnormal conditions of use or storage are beyond Moyer's control, Moyer's liability is limited to replacement of product or refund of purchase price. In no event shall Moyer be liable for indirect or consequential damages.

**BEST AVAILABLE COPY**