DIBROM®-SEVIN® DUST NO. 4-5



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

This product is an agricultural pesticide for application by ground or air for the below indicated uses.

Begin applications at first sign of insects. Repeat as necessary.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, **CABBAGE:** Aphids, Diamond-Back Moths — Use 40 lbs. per acre. Do not apply within 3 days of harvest.

PEPPERS: Aphids, Leafminers—Use 25 lbs. per acre. Do not apply within 1 day of harvest.

GRAPES: Leafhoppers, Leafrollers — Use 40 lbs. per acre. Drosophila, Thrips—Use 25 lbs. per acre. Do not apply within 4 days of harvest.

LETTUCE, SPINACH, TURNIP GREENS: Aphids, Armyworms - Use 40 lbs. per acre. On HEAD LETTUCE—Do not apply within 3 days of harvest. Do not apply within 14 days of harvest on LEAF LETTUCE, SPINACH, and TURNIP GREENS.

PEACHES: Mites, Oriental Fruit Moth — Use 40 Ibs. per acre. Drosophila — Use 50 lbs. per acre. Codling Moths — Use 80 lbs. per acre, apply at control is achieved. Orange Tortrix — Use 80 lbs. ACCEAPTED per acre. Do not apply within 4 days of harvest.



ACTIVE INGREDIENTS

**Naled Carbaryl (1-naphthyl N-methylcarban INERT INGREDIENTS

TOTAL...

****1,2-dibromo-2,2-dichloroethyl dimethyl phosphate.** U.S. PAT. NO. 2,971,882

DIBROM® is a registered trademark of Chevron Chemical Company. SEVIN® is a registered trademark of Union Carbide Corporation for carbaryl.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin and eyes. May cause irritation of eyes, nose and throat. Wash thoroughly after using. Do not store or transport near feed or food.


NOTE TO PHYSICIAN

Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Atropine is also the antidote for Naled.

SEE OTHER PRECAUTIONS ON SIDE PANEL

DATE OF MANUFACTURE

USE WITHIN DAYS OF MANUFACTURE





	By Weigh
• • • • • • • • • • • • • • • • • • •	4.0%
mate)	5.0%
· · · · · · · · · · · · · · · · · · ·	91.0%
••••••••••••••••••••••••••••••••••••••	100.0%

SAN JOSE, CALIFORNIA



USE PRECAUTIONS

THIS PRODUCT IS TOXIC TO FISH, BIRDS AND OTHER WILDLIFE. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

May corrode metal dust equipment; clean equipment after use.

KEEP PESTICIDE IN ORIGINAL CONTAINER. SUNLIGHT.

BURN BAG IMMEDIATELY WHEN EMPTY-STAY OUT OF SMOKE.

Do not permit workers to enter the treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes 2; Peaches 2. STORE IN COOL, DRY PLACE, OUT OF DIRECT

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and/or handling is contrary to label instructions.

--- NON-RETURNABLE ----