

ENDOSULFAN DUST NO. 5

DIRECTIONS FOR USE:

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container.

This product is an agricultural pesticide for application by ground or air for the following uses:

ARTICHOKES: Artichoke Aphid, Green Peach Aphid, Artichoke Plume Moth—Apply at 20 to 30 lbs. per acre when aphids first appear and repeat as required. Dust plants thoroughly. Do not apply after buds start to form.

BROCCOLI, CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER: Cabbage Aphid, Cabbage Looper, Imported Cabbageworm, Diamond-Back Moth Larvae, Cross-Striped Cabbageworm, Flea Beetle, Harlequin Bug, Stink Bug—20 lbs. per acre. Do not apply to Broccoli and Cabbage within 7 days of harvest and to Brussels Sprouts and Cauliflower within 14 days of harvest.

CUCUMBERS, PUMPKINS, SUMMER and WINTER SQUASH: Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer—20 lbs. per acre. For Vine Borer, apply weekly to flower buds, stems and vines beginning when moths first appear. May be applied up to day of harvest.

GRAPES: Grape Leafhoppers—20 to 30 lbs. per acre. Do not use on Concord variety of grapes as severe injury is likely to occur. Do not apply within 7 days of harvest.

LETTUCE (Head): Cabbage Looper, Green Peach Aphid, Diamond-Back Moth Larvae, Imported Cabbage Worm—20 lbs. per acre. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove wrapper leaves. Do not feed crop refuse to livestock.

LETTUCE (Leaf): Cabbage Looper, Green Peach Aphid, Diamond-Back Moth Larvae, Imported Cabbage Worm—20 lbs. per acre. Do not apply within 14 days of harvest. Do not make more than 2 applications per season.

MELONS: Aphids, Cucumber Beetle, Striped Cabbage Flea Beetle—20 to 25 lbs. per acre. Cabbage Looper, Omnivorous Leaf Roller—25 to 30 lbs. per acre. May be applied up to day of harvest.

ORNAMENTALS: Aphids, White Fly—Apply dust thoroughly to plants. On Chrysanthemums, best results will be obtained if applied before plants flower. Do not use on White Birch.

PEACHES: Cat-Facing Insects, Green Peach Aphid—40 to 60 lbs. per acre. Apply to give thorough coverage to all plant parts. Do not apply within 30 days of harvest, nor make more than 2 applications during the fruiting period.

PEPPERS: Green Peach Aphid, Flea Beetles, Hornworms, Pepper Maggot—20 to 30 lbs. per acre and do not apply within 1 day of harvest.

POTATOES: Flea Beetle, Colorado Potato Beetle, Leafhopper, Aphids, Potato Psyllid—20 lbs. per acre. No time limitations. Do not plant root crops other than carrots, potatoes and sweet potatoes as follow-up crops.


STRAWBERRIES: Strawberry Aphid, Meadow Spittlebug—2 to 20 lbs. per acre. Do not apply within 4 days of picking. Do not apply at intervals of less than 15 days, or more than twice during any 35-day period when fruit is present. Cyclamen Mite—75 lbs. per acre. Dust crown of each plant thoroughly. Do not apply at intervals of less than 35 days when fruit is present nor within 4 days of picking.

TOMATOES: Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Whitefly—20 lbs. per acre. Do not use within 1 day of harvest. Hornworm—20 lbs. per acre. Do not apply to tomatoes within 1 day of picking.

WALNUTS: Walnut Aphid—40 lbs. per acre. Apply when aphids first appear and repeat as required. Do not apply after husks split. Do not graze livestock in treated groves.

ACTIVE INGREDIENTS	By Weight
*Endosulfan	5.0%
INERT INGREDIENTS	95.0%
TOTAL	100.0%

*Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide.



CAUTION:

Keep Out of Reach of Children

SEE OTHER PRECAUTIONS ON THE SIDE PANELS

EPA REGISTRATION NO. 5967-115 AA

EPA EST. 5967-CA-1

11/74
SULFAMEN
 MANUFACTURED BY

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

CAUTION:

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; For eyes — get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Avoid contamination of feed and foodstuffs.

WORKERS ENTERING TREATED AREAS WITHIN 24 HOURS OF APPLICATION SHOULD WEAR PROTECTIVE CLOTHING.

USE PRECAUTIONS:

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply to or allow drift onto areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply under condition involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Do not apply to or allow dust to drift onto sulfur sensitive crops such as apricots or melons. Do not apply during or prior to periods of high temperature (90° F or higher) as sulfur damage may occur. Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur unless local use has proved that sulfur does not damage crops in that locality.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

— NON-RETURNABLE —

NET WEIGHT 50 LBS.