

[1032]

ETHION 25 WP

THIS PRODUCT IS AN AGRICULTURAL PESTICIDE FOR THE BELOW INDICATED USES:

CROP	PEST	LBS. PER 100 GALLONS	REMARKS	LIMITATIONS
Almonds	European Red Mite Two-Spotted Mite	1-2 LBS.	Apply 200 to 350 gallons per acre.	Do not apply within 30 days of harvest. Do not apply more than once during fruiting season. Do not graze livestock in treated orchards.
Apples	European Red Mite Two-Spotted Mite	1 LB.	Apply 400 to 800 gallons per acre.	Do not apply more than 3 times after first cover. Do not apply within 30 days of harvest. Do not use on Wealthy or Melba types as injury is likely to occur.
Cotton	Spider Mites Lygus Bugs and Other Mirids	4 to 6 lbs. per acre in sufficient water to give uniform coverage to all plant surfaces.	Workers entering fields within 24 hours should wear protective clothing.	Do not apply after bolls open. Do not feed treated foliage to livestock.
Grapes	Grape Leafhopper Two-Spotted Mite Willamette Mite	1 LB.	Apply 200 to 300 gallons per acre.	Do not apply within 30 days of harvest. Do not apply more than twice in the fruiting season.
Melons	Leafminers	2 to 4 lbs. per acre.	Apply 10 to 100 gals.	Do time limitation.
Peaches	European Red Mite Two-Spotted Mite	1-2 LBS.	Apply 200 to 350 gallons per acre.	Do not apply more than twice in fruiting season or within 30 days of harvest.
Pears	European Red Mite Two-Spotted Mite	1 LB.	Apply 400 to 660 gallons per acre.	Do not apply within 30 days of harvest. Do not apply more than 3 times after first cover spray or 20 days before harvest, do not apply more than twice after first cover spray.
	Pear psylla Codling moth	2 LBS.	Apply 500 gals. per acre.	
Prunes	European Red Mite Two-Spotted Mite European Fruit Lecanium scales	1 LB. 1½ LBS.	Apply 460 gallons per acre.	Do not apply within 21 days of harvest. Do not apply more than 3 times during fruiting season. Do not apply more than 460 gals. of spray per acre to prunes at the higher rate.
Walnuts	Spider Mites	1-2 LBS.	Apply 250 to 500 gallons per acre.	Do not apply after husk split. Do not apply more than twice during fruiting season. Do not graze livestock in treated groves.

Do not permit workers to enter treated area where substantial contact will be made with foliage until 14 days after application on Grapes, Peaches and Nectarines

EPA REGISTRATION NO. 5967-118 AA

EPA EST. 5967-CA-1

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. —NON RETURNABLE—

NET WEIGHT 4 LBS.

ACCEPTED

5967-118

[272]

MOYER**ETHION 25 WP****ACTIVE INGREDIENTS**

*Ethion	By Weight
INERT INGREDIENTS	25.00%
TOTAL	75.00%
	100.00%

*O,O,O',O'-Tetraethyl S,S'-methylene bisphosphorodithioate

*U.S. Patent No. 2,873,228

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDES IN ORIGINAL CONTAINERS. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WARNING**Do Not Inhale, Do Not Get on Skin, Do Not Take Internally.**

WARNING — MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing dust and spray mist. Wear dust respirator and goggles. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not contaminate feed and foodstuffs. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Do not store or transport near feed or food.

ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

WARNING: Keep Out of Reach of Children**EPA REGISTRATION NO. 5967-118 AA**

EPA EST. 5967-CA-1

8/74

MANUFACTURED BY

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY

P. O. BOX 945

SAN JOSE, CALIFORNIA

[1022]

ETHION 75 WP

THIS PRODUCT IS AN AGRICULTURAL P*

THE BELOW INDICATED USES:

CROP	PEST	LBS. PER 100 GALL.	MARKS	LIMITATIONS
Almonds	European Red Mite Two-Spotted Mite	1-2 LBS.	Apply 200 to 350 gallons per acre.	Do not apply within 30 days of harvest. Do not apply more than once during fruit- ing season. Do not graze livestock in treated orchards.
Apples	European Red Mite Two-Spotted Mite	1 LB.	Apply 400 to 600 gallons per acre.	Do not apply more than 3 times after first cover. Do not apply within 30 days of harvest. Do not use on Wageningen or Melba types as injury is likely to occur.
Cotton	Spider Mites Lycus Bugs and Other Mirids	4 to 6 lbs. per acre in sufficient water to give uniform coverage to all plant surfaces.	Workers entering fields within 24 hours should wear protec- tive clothing.	Do not apply after bolls open. Do not feed treated foliage to livestock.
Grapes	Grape Leafhopper Two-Spotted Mite Willamette Mite	1 LB.	Apply 200 to 300 gallons per acre.	Do not apply within 30 days of harvest. Do not apply more than twice in the fruiting season.
Melons	Leafminers	2 to 4 lbs. per acre.	Apply 10 to 100 gals.	Do time limitation.
Peaches	European Red Mite Two-Spotted Mite	1-2 LBS.	Apply 200 to 350 gal- lons per acre.	Do not apply more than twice in fruiting season or within 30 days of harvest.
Pears	European Red Mite Two-Spotted Mite Pear psylla Coding moth	1 LB. 2 LBS.	Apply 400 to 600 gal- lons per acre. Apply 500 gals. per acre.	Do not apply within 30 days of harvest. Do not ap- ply more than 3 times after first cover spray or 20 days before harvest, do not ap- ply more than twice after first cover spray.
Prunes	European Red Mite Two-Spotted Mite European Fruit Lecanium scales	1 LB. 1 1/2 LBS.	Apply 400 gallons per acre.	Do not apply within 21 days of harvest. Do not ap- ply more than 3 times dur- ing fruiting season. Do not apply more than 400 gals. of spray per acre to prunes at the higher rate.
Walnuts	Spider Mites	1-2 LBS.	Apply 250 to 500 gallons per acre	Do not apply after husk split. Do not apply more than twice during fruiting season. Do not graze live- stock in treated groves.


Do not permit workers to enter treated area where substantial contact will be made with foliage until 14 days after application on Grapes, Peaches and Nectarines.

EPA REGISTRATION NO. 5967-118 AA

EPA EST. 5967-CA-1

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. —NON-RETURNABLE—

NET WEIGHT 4 LBS.



[292]



ETHION 25 WP

ACTIVE INGREDIENTS		By Weight
*Ethion		25.00%
INERT INGREDIENTS		75.00%
TOTAL		100.00%

*O O O O - Tetraanthyl S,S'-methylene bisphosphorodithioate *P. No. 2873 228

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDES IN ORIGINAL CONTAINER - DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS

WARNING

Do Not Inhale, Do Not Get on Skin, Do Not Take Internally.

WARNING - MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing dust and spray mist. Wear dust respirator and goggles. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. For eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Do not contaminate feed and foodstuffs. For commercial application wear rubber gloves, protective clothing, rubber overshoes and goggles. Do not store or transport near feed or food.

ATROPINE IS ANTIDOTAL. P. 2873 228 is an antitoxin and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected humans. For all animals or other application for food storage crop, about to be harvested.

This product is toxic to fish, birds, and other wildlife. Keep out of lake, streams, or ponds. Do not apply where birds, fish, or other wildlife are trapped, when weather conditions favor drift from areas treated. Do not contaminate water by leakage of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and other beneficial insects. Do not apply to the area.

WARNING: Keep Out of Reach of Children

EPA REGISTRATION NO. 5967-118 AA
EPA EST. 5967 CA-1

R 74

MANUFACTURED BY

MOYER CHEMICAL COMPANY

210 BAYSHORE HIGHWAY

P. O. BOX 945

FAIR OAKS, CALIF. 94534

