

DIAZINON® DUST NO. 4

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER. BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS FOR USE:

Apply when air is calm. Apply when insects appear. Repeat as necessary. This product is formulated for application by ground or air. Make thorough coverage of plants with a uniform deposit of dust.

BEANS (Pole and Snap): For control of Black Bean Aphids, Leafminers, Mexican Bean Beetles — Use 25 to 35 lbs. per acre. Do not apply within 7 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI: For control of Aphids, Diamond Back Moths, Imported Cabbage Worm — Use 20 to 25 lbs. per acre. Apply when insects first appear, repeat as necessary. Do not apply within 7 days of harvest on Cabbage or within 5 days of harvest on Cauliflower or Broccoli.

CELERY: For control of Aphids, Flea Beetles — Use 10 to 12 lbs. per acre. Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.

CLOVER: For control of Aphids, Leafhoppers — Use 15 lbs. per acre. Do not apply within 7 days of cutting or grazing by animals being finished for slaughter.

CORN: For control of Corn Earworm — Use 30 to 40 lbs. per acre. Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days of treatment.

CUCUMBERS: For control of Aphids, Leafhoppers, Thrips — Use 30 to 40 lbs. per acre. Do not apply within 7 days of harvest.

GRAPES: For control of Pacific Mite, Leafhoppers — Use 20 to 25 lbs. per acre. Do not use within 10 days of harvest.

LETTUCE: For control of Aphids, Dipterous Leafminers — Use 10 to 12 lbs. per acre. Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.

LIMA BEANS: For control of Black Bean Aphids, Leafminers, Leafhoppers, Mexican Bean Beetles — Use 25 to 35 lbs. per acre. Do not use within 7 days of harvest.

MELONS (Cantaloupes, Muskmelons, Watermelons, Summer Squash, Winter Squash): For control of Aphids, Leafhoppers, Leafminers, Thrips — Use 25 to 35 lbs. per acre. Do not use within 3 days of harvest. On Summer Squash — Do not apply within 7 days of harvest.

ONIONS (Including Green Onions): For control of Onion Thrips — Use 20 to 30 lbs. per acre. Do not apply within 10 days of harvest.

PEACHES, NECTARINES: For control of Scale Crawlers, Clover Mite, Pacific Mites — Use 40 to 60 lbs. per acre. Do not use within 20 days of harvest.

PEPPERS: For control of Aphids — Use 25 to 40 lbs. per acre. Apply when insects first appear, repeat as necessary. Do not apply within 5 days of harvest.

SPINACH: For control of Aphids, Dipterous Leafminers — Use 10 to 12 lbs. per acre. Apply when insects first appear, repeat as necessary. Do not apply within 10 days of harvest.


STRAWBERRIES: For control of Aphids, Two-Spotted Mites — Use 20 to 30 lbs. per acre. Apply when insects first appear, repeat as necessary. Thorough underleaf coverage is essential for good control. Do not apply within 5 days of harvest.

TOMATOES: For control of Aphids, Dipterous Leafminers, Cucumber Beetles — Use 30 to 40 lbs. per acre. Do not apply within 3 days of harvest. If applied within 1 day of harvest — do not use more than 15 to 20 lbs.

ORNAMENTALS (Azalea, Camellia, Carnation, Chrysanthemum, Gladioli, Rose): For control of Aphids, Carnation Bud Mites, Carnation Shoot Mites, Whiteflies — Use 35 to 60 lbs. per acre. Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pitta, and Gardenia, as injury to plants may occur.

ACTIVE INGREDIENT	By Weight
*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	4.0%
INERT INGREDIENTS	96.0%
TOTAL.....	100.0%

Diazinon® is the registered trademark of CIBA-Geigy Corp.



CAUTION:

Keep Out of Reach of Children

SEE OTHER PRECAUTIONS ON THE SIDE PANELS

EPA REGISTRATION NO. 5967-122 AA
EPA EST. 5967-CA-1

11/74

MANUFACTURED BY

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA



CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

In case of contact with skin or eyes, flush with plenty of water: **FOR EYES**, get medical attention.

Wash with soap and water after handling and before eating or smoking: Wear clean clothing.

Do not store or transport near feed or food.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed.

Keep out of lakes, streams, ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

USE PRECAUTIONS:

Do not permit workers to enter treated area where substantial contact will be made with treated foliage until following indicated days have lapsed after application: Grapes, 5; Peaches and Nectarines, 5.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

— NON-RETURNABLE —

NET WEIGHT 50 LBS.