DIBROM®-SEVIN® SULFUR DUST

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

THIS PRODUCT IS AN AGRICULTURAL PESTI-CIDE FOR APPLICATIONS BY GROUND OR AIR FOR THE BELOW INDICATED USES.

BEGIN APPLICATIONS AT FIRST SIGN OF IN-SECTS. REPEAT AS NECESSARY.

GRAPES: For Leafhoppers, Red Banded Leafrollers, Powdery Mildew—use 40 lbs. per acre. For Drosophila, Thrips—use 25 lbs. per acre.

For Powdery Mildew: Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur.

Do not apply within 4 days of harvest.

PEACHES: For Mites. Oriental Fruit Moth, Powdery Mildew—use 40 lbs. per acre. For Drosphila—use 50 lbs. per acre. For Orange Tortrix—use 80 lbs. per acre. For Orange Tortrix and Oriental Fruit Moth apply at petal fall and every 10 to 14 days thereafter until control is achieved. For Powdery Mildew begin at first sign of disease and repeat as necessary. Do not apply within 4 days of harvest.

NET WEIGHT 50 LBS.

4

NO. 4-5-25

ACTIVE INGREDIENTS Naled*	•••••	By Weight 4.0%
Carbaryl**		, ,
Sulfur	•••••	25.0%
INERT INGREDIENTS		
	TOTAL	100.0%

*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate
U.S. PAT. NO. 2,971,882

DIBROM® is a registered trademark of Chevron Chemic

**1-naphthyl N-methylcarbamate

SEVIN® is a registered trademark of Union Carbide Corporation for the second secon

USE WITHIN 60 DAYS OF MFG.

CAUTION: KEEP OUT OF REACH OF CHIEDRE

2.75

EPA REGISTRATION NO. 5967-123 AA EPA EST. 5967-CA-1

MANUFACTURED BY

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY

P.O. BOX 945

SAN JOSE, CALIFORNIA

H

197

CAUTION:

AVOID EYE CONTACT. HARMFUL IF SWALLOWED OR INHALED. MAY CAUSE IRRITATION OF EYES, NOSE AND THROAT. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN. Wash thoroughly after using. Do not store or transport near feed or food.

May corrode metal dust equipment; clean equipment after use. Do not apply when weather conditions favor drift of spray from areas treated.

USE PRECAUTIONS:

THIS PRODUCT WILL KILL FISH AND WILDLIFE. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT RE-USE EMPTY CONTAINER. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DO NOT FERMIT WORKERS TO ENTER THE TREATED AREA where substantial contact will be made with treated foliage until following indicated days have lapsed after application Grapes 2, Peaches 2.

BURN BAG IMMEDIATELY WHEN EMPTY — STAY OUT OF SMOKE!

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

- NON-RETURNABLE -