

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. _____

CAUTION

MAY BE HARMFUL IF SWALLOWED

AVOID BREATHING OF SPRAY MIST.

Before using this product, be sure the spray tank is clean and contains just enough water to cover the agitator shaft.

With the agitator in motion (120 R.P.M. or more) add the amount of spray oil required for a full tank. Pump this mixture through the overflow and back into the tank for a minute or two. The mixture should turn white, indicating good emulsification. Then fill the tank with water.

If an injector is used, first add 25 to 50 gallons of water to tank with engine running to provide good pressure. Then put injector suction tube into the measured amount of spray oil, opening injector valve to allow the oil to be sucked into the tank. Then fill the tank with water.

DO NOT USE THIS MATERIAL IF IT DOES NOT EMULSIFY.

ENVIRONMENTAL CAUTIONS

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

Fish and other Aquatic life may be killed by this product. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

— NON-RETURNABLE —

30 GALLONS



DORMANT SOLUBLE

ACTIVE INGREDIENT	By Weight
Petroleum Oil	99.00%
INERT INGREDIENTS	1.00%
TOTAL.....	100.00%

Classification — Dormant Heavy
Unsulphonated Residue — 70% Minimum

CAUTION:

Keep Out of Reach of Children

EPA REGISTRATION NO. 5967-127 AA

EPA EST. 5967-CA-1

9-74

SEE SIDE PANEL FOR CAUTION STATEMENT

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

KEEP THIS
READ ENTIRE
CAUTIONS
AND FEDERAL
KEEP PEST
CONCENTRATE

DORMANT SOLUBLE
used on the
full dormant

BROWN APPL
Cherries, Pe
Use 2 1/2 to

BROWN ALM
SCALE: Appl
and other d
spray.

LEAFROLLER
gals. per 10

STRAWBERRY
wintering Sp
to 3 pts. of a
NITRO KILL
indicated be

By Ground —

By Air — ap
of spray solu

Use 400 to 8
ages, consult

TO OBTAIN
ATTENTION

Use efficient
or immediate
or other unsu

Plants should
their moisture
Agricultural B
area

DORMANT SOLUBLE

ACTIVE INGREDIENT	By Weight
Petroleum Oil	99.00%
INERT INGREDIENTS	1.00%
TOTAL.....	100.00%

Classification — Dormant Heavy
Unsulphonated Residue — 70% Minimum

CAUTION:

Keep Out of Reach of Children

EPA REGISTRATION NO. 5967-127 AA

EPA EST. 5967-CA-1

9 74

SEE SIDE PANEL FOR CAUTION STATEMENT

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE CALIFORNIA

KEEP THIS DRUM IN THE SHADE.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE IN FOOD OR DRINK CONTAINER.

DIRECTIONS FOR USE

DORMANT SOLUBLE OIL is a prepared emulsive dormant oil to be used on the crops appearing on this label when these are in the full dormant period.

BROWN APRICOT SCALE AND BLACK SCALE: Apricots, Almonds, Cherries, Peaches, Prunes, Nectarines and other deciduous fruits. Use 2½ to 3½ gals. per 100 gals. of spray.

BROWN ALMOND MITE EGGS, ITALIAN PEAR SCALE, SAN JOSE SCALE: Apricots, Almonds, Cherries, Peaches, Prunes, Nectarines and other deciduous fruits. Use 3½ to 4 gals. to 100 gals. of spray.

LEAFROLLER EGGS: Apples, Apricots, Pears, Prunes. Use 4 to 5 gals. per 100 gals. of spray.

STRAWBERRIES: To Burn Back Dormant Plants and Reduce Overwintering Spider Mite Population. Use 5 to 10 gals. per acre plus 2 to 3 pts. of a General Dinitro Weedkiller or 2 to 3 qts. of MOYER'S NITRO KILL WEEDKILLER. Add enough water to obtain solutions indicated below:

By Ground — apply a minimum 75 gals. of spray solution per acre.

By Air — apply a minimum of 10 gals. and not more than 20 gals. of spray solution.

Use 400 to 800 gallons of water per acre; for concentrate spray dosages, consult your Cooperative Agricultural Extension Service.

TO OBTAIN BEST RESULTS WITH THIS PRODUCT, SPECIAL ATTENTION SHOULD BE GIVEN TO THE FOLLOWING:

Use efficient equipment of the proper type, avoid spraying during or immediately prior to hot or freezing weather, hot dry winds, rain or other unsuitable conditions, avoid over-spraying or double-spraying.

Plants should be sprayed only when in vigorous condition and when their moisture condition is suitable. Consult with your Cooperative Agricultural Extension Service for maximum dosage rate in your local area.