

# MALATHION-SULFUR DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

## DIRECTIONS FOR USE:

Apply when air is calm. Apply when insects or diseases appear. Repeat as necessary. This product is formulated for application by ground or air. Make thorough coverage of plants with a uniform deposit of dust.

**BEANS:** For the control of Aphids, Mexican Bean Beetles, Leafhoppers, Lygus Bugs, Spider Mites and Thrips — Use 45 lbs. per acre. Apply at first sign of insects and repeat at regular intervals as necessary to maintain control. For Powdery Mildew — apply at same rate — Begin at first sign of disease and every 7 to 10 days as necessary to maintain control. Do not apply within 1 day of harvest.

**GRAPES:** For the control of Aphids, Orange Tortrix, Spider Mites and Thrips — Use 20 to 30 lbs. per acre. Apply at first sign of insects. Repeat regularly at 2-week intervals or as necessary. For Powdery Mildew — apply at same rate — Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. Concord and other labrusca type grapes may be injured by sulfur. Do not apply within 3 days of harvest.

**PEAS:** For the control of Aphids and Leafhoppers — Use 50 lbs. per acre. Apply at first sign of insects and repeat at regular intervals as necessary to maintain control. For Powdery Mildew — apply at same rate — Begin at first sign of disease and repeat every 7 to 10 days as necessary to maintain control. Do not apply within 7 days of harvest.

**STRAWBERRIES:** For the control of Aphids, Lygus Bugs, Strawberry Leafroller, Thrips and White Flies — Use 50 lbs. per acre. Apply at first sign of insects and repeat as necessary. For Powdery Mildew — apply at same rate — Begin at first sign of disease and repeat as necessary to maintain control. Sulfur may burn Strawberries when temperatures are excessively high. Do not apply within 3 days of harvest.

# NO. 4-25

ACTIVE INGREDIENTS	By Weight
Malathion* .....	4.0%
Sulfur .....	25.0%
INERT INGREDIENTS .....	71.0%
TOTAL.....	100.0%

\*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

SEE OTHER PRECAUTIONS ON THE SIDE PANELS

## CAUTION:

### KEEP OUT OF REACH OF CHILDREN

EPA REGISTRATION NO. 5967-131 AA

ACCEPTED  
1174

FEB 6 1975

EPA EST. 5967-CA-1

MANUFACTURED BY  
**MOYER CHEMICAL COMPANY**

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

5967-131

## CAUTION:

HARMFUL IF SWALLOWED, MAY CAUSE IRRITATION OF EYES, NOSE AND THROAT. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN. Wash thoroughly after use. Do not store or transport near feed or food.

Do not apply to or allow dust to drift onto sulfur sensitive crops such as apricots or melons. Do not apply during or prior to periods of high temperature (90° F or higher) as sulfur damage may occur. Some fruit and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur unless local use has proved that sulfur does not damage crops in that locality.

Sulfur may burn foliage when temperatures are excessively high.

## THIS PRODUCT IS TOXIC TO FISH.

Do not contaminate lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment of residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**SULFUR** is listed as a safe chemical with no time limitations. However: Do not allow workers to enter treated vineyard where substantial contact will be made with treated foliage until 1 day after application.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET WEIGHT 50 LBS.

## DIRECTIONS FOR USE:

This material is an agricultural pesticide for the following uses:

### ORNAMENTALS:

For the control of Aphids, Mealybugs, Spider Mite, White Flies and Thrips, use one pint per 100 gallons of water. Apply so that all leaf surfaces receive thorough coverage. Repeat application as often as necessary to maintain adequate control. Injury may occur to Boston, Maidenhair or Pteris Ferns and to certain species of Crassula.

NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

— NON-RETURNABLE —

NET CONTENTS ..... GALLONS

# Malathion Spray 8 Miscible

ACTIVE INGREDIENT	By Weight
Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate) .....	83.0%
INERT INGREDIENTS .....	17.0%
TOTAL .....	100.0%

THIS MATERIAL CONTAINS 8 POUNDS OF MALATHION PER GALLON

**CAUTION: Keep Out of Reach of Children**

**SEE RIGHT PANEL FOR PRECAUTIONS**

**EPA REGISTRATION NO. 5967-139 AA**

EPA EST. 5967-CA-1

11, 74

MANUFACTURED BY

**MOYER CHEMICAL COMPANY**

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

Do not  
apply  
containing

Harmful  
contact  
with  
skin

**EI**

This  
product  
applied.  
Do not  
disperse  
the liquid

This  
treatment  
may  
cause  
irritation

DO NOT  
OPEN