

THIS PRODUCT IS FORMULATED FOR USE BY GROUND APPLICATION FOR THE FOLLOWING:

DIRECTIONS FOR USE:

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER: Root Maggots — Broadcast Application: Apply 40-60 lbs. per acre over the entire seedbed and disk into the top 3-4 inches before seeding.

Furrow Application: Apply 20 lbs. per acre directly in the furrow on the seed.

ONIONS: Onion Maggots: Apply 20 lbs. per acre in the furrow with the seed at the time of planting.

Wireworms: Apply 40-80 lbs. per acre as a pre-plant broadcast treatment. Thoroughly disk or harrow to a depth of 4-8 inches.

RADISHES: Root Maggots: Apply 30-40 lbs. per acre (6-8 ozs. per 1,000 linear feet of row) in the seed furrow at planting time.

TURNIPS: Root Maggots: Apply 20 lbs. per acre in the seed furrow at planting time.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, ENDIVE (ESCAROLE), LETTUCE:

Cutworms: As a broadcast soil treatment; apply 40-80 lbs. per acre immediately before planting. Work into the top 2-3 inches of soil.

For subterranean cutworms—DIAZINON NO. 5G should be worked into the top 3-6 inches of soil.

Wireworms: Broadcast Application: Apply 40-80 lbs. per acre on the soil as a preplant broadcast treatment. Thoroughly disk or harrow into the upper 4-8 inches.

Mole Crickets: As a broadcast soil treatment to the above crops — apply 20 lbs. per acre. Make application 1-2 days prior to sowing of seed or transplanting, and mix in the top 1 or 2 inches of soil.

LAWN INSECTS: Lawn Chinch Bugs: Apply 1½-2 lbs. per 1,000 sq. ft. of lawn (6½-10 lbs. per 5,000 sq. ft. of lawn). Application should be repeated in 7-10 days if necessary.

For longer residual control, and in lawns of thicker growth, such as St. Augustine lawns, use 2-4 lbs. per 1,000 sq. ft. of lawn (10-20 lbs. per 5,000 sq. ft. of lawn) Following application, brush the lawn to assure that all granules reach the soil surface.

White Grubs, such as Japanese Beetle Larvae, European Chafer and Southern Chafer: Apply 2½ lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For grub control — make application any time between late July and early October.

CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN AND EYES. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. WASH THOROUGHLY AFTER HANDLING.

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from treated areas. Do not apply where run-offs is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Do not apply while bees are working in the field.

Burn empty bags. Stay out of smoke.

**DATE OF MANUFACTURE
DO NOT USE AFTER 60 DAYS OF MANUFACTURE.**

Dispose of spillage in accordance with good environmental sanitation practices.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

SPECIMEN
NET WEIGHT 50 LBS.

ACCEPTED

-15-75

FEDERAL INSECTICIDE

AND FUNGICIDE ACT

1947

15 U.S.C. 136

15 U.S.C. 136a

15 U.S.C. 136b

15 U.S.C. 136c

15 U.S.C. 136d

15 U.S.C. 136e

15 U.S.C. 136f

15 U.S.C. 136g

15 U.S.C. 136h

15 U.S.C. 136i

15 U.S.C. 136j

15 U.S.C. 136k

15 U.S.C. 136l

15 U.S.C. 136m

15 U.S.C. 136n

15 U.S.C. 136o

15 U.S.C. 136p

15 U.S.C. 136q

15 U.S.C. 136r

15 U.S.C. 136s

15 U.S.C. 136t

15 U.S.C. 136u

15 U.S.C. 136v

15 U.S.C. 136w

15 U.S.C. 136x

15 U.S.C. 136y

15 U.S.C. 136z

15 U.S.C. 136aa

15 U.S.C. 136ab

15 U.S.C. 136ac

15 U.S.C. 136ad

15 U.S.C. 136ae

15 U.S.C. 136af

15 U.S.C. 136ag

15 U.S.C. 136ah

15 U.S.C. 136ai