

1 (of 2)

MALATHION 25-W

ACTIVE INGREDIENT	By Weight
Malathion*	25.00%
INERT INGREDIENTS	75.00%
TOTAL	100.00%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate

READ THE ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. KEEP PESTICIDES IN ORIGINAL CONTAINERS. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

COMPLETELY EMPTY BAG AND DISPOSE OF WASTE PESTICIDE BY BURYING. BURN BAG IMMEDIATELY AND STAY OUT OF SMOKE.

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed or foodstuffs.

NOTE TO PHYSICIANS: May cause cholinesterase inhibition. Atropine is antidotal.

USE PRECAUTIONS:

Burn empty bags. Keep out of smoke. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish, shrimp, crab and other aquatic organisms. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

— NON-RETURNABLE —

NET WEIGHT 5 LBS.
EPA REGISTRATION NO. 5967-140 AA
EPA EST. 5967-CA-1

MANUFACTURED BY
MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P.O. BOX 945 SAN JOSE, CALIFORNIA

2 (of 2)

MALATHION 25-W

DIRECTIONS FOR USE:

Mix in water with agitation; agitate solution while spraying. Dosages are calculated for conventional sprayers applying 100 gallons or more per acre as specified below. For use with concentrate sprayers applying 40 to 90 gallons per acre, adjust dosage to apply the same amount of actual material as would be applied with a conventional sprayer.

APPLES: To control Aphids and Mealybugs, use 2 to 2½ lbs. per 100 gals. Use 500-600 gals. per acre. Do not apply within 3 days of harvest. May injure foliage on some varieties.

CITRUS FRUITS: To control California Red, Yellow, Purple, Citricola, Black, Brown Soft Scale and Aphids, use 2½ to 3½ lbs. per 100 gals. To control Fuller's Rose Weevil, use 1¾ lbs. per 100 gals. Apply 400 to 2,000 gals. per acre. Do not apply within 7 days of harvest. Do not apply during full bloom. Do not permit workers to enter treated citrus groves where substantial contact will be made with treated foliage until 1 day after application.

PEARS: To control Aphids, Mealybugs and Psylla, use 2 lbs. per 100 gals. Do not use within 1 day of harvest. Use 500 to 800 gals. per acre

WALNUTS: To control Walnut Aphid, use 4 to 6 lbs. per acre; to control Walnut Husk Fly, use 3 lbs. per acre. Apply in 200-300 gals. water by air blast sprayer.

ORNAMENTALS: Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle, Four-lined Leaf Bug, Tarnished Plant Bug, Leafhoppers—2½ lbs. Soft Brown Scale, Monterey Pine Scale, Black Scale, Juniper Scale, Florida Red Scale—6 lbs. Pine Needle Scale—4 lbs. Oak Kermes, Flatheaded Scale—3 lbs. Tent Caterpillar—2 lbs. Locust Broomworms—4 lbs. Repeat as necessary. To control Listed Scale—Apply 1 lb. per 100 gal. spray at time scale crawlers have settled on foliage. Do not use on Boston, Maidenhair and Pteris ferns, and some species of ornamental plants.

ACCEPTED