

WETTABLE SULFUR

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container.

Burn bag immediately when empty — Stay out of smoke

DIRECTIONS FOR USE:

This product is an agricultural pesticide for application by ground or air for the below indicated uses:

ALMONDS: For control of Scab—Use 6 lbs. per 100 gallons. Begin when disease threatens or in pre-bloom stage. Repeat at 7-day intervals as long as necessary.

APPLES: For control of Scab—Use 3 to 4 lbs. plus 2 gals. liquid lime sulfur in 100 gals. For control of Mildew—Use 5 lbs. per 100 gals. Start at pre-pink through calyx periods and include in cover sprays. *Note:* McIntosh, Golden Delicious, Jonathan, Belleflower and certain other varieties may be injured by sulfur applications under certain conditions.

CHERRIES: For control of Brown Rot and Powdery Mildew—Use 6 lbs. per 100 gals. Begin in pink stage. Repeat at bloom, petal fall and in cover stages at 7 to 10 day intervals. Do not spray when pollination is taking place.

GRAPES: For control of Powdery Mildew—Use 6 lbs. plus wetting agent. Begin when new shoots are 6 inches long. Repeat before blossoms open and continue as necessary. *Note:* Concord and other labrusca type grapes may be injured by sulfur.

ORANGES, LEMONS, GRAPEFRUIT, TANGERINES: For control of Six-Spotted Mites, Citrus Rust Mites, and Citrus Flat Mites—Use 4 lbs. per 100 gals. For control of Brown Rot: Use 5 to 7 lbs. per 100 gals. Consult your local Cooperative Agricultural Extension Service for recommendations for your local area.

PEACHES, NECTARINES: For control of Brown Rot and Rust—Use 4 lbs. per 100 gals. Apply at pink, bloom, petal fall, shuck split, shuck fall and at 10 to 14 day intervals as long as necessary. For Brown Rot—apply as fruit starts to ripen.

ACTIVE INGREDIENT	By Weight
Sulphur	95.00%
INERT INGREDIENTS	5.00%
TOTAL.....	100.00%

**CAUTION:
KEEP OUT OF REACH OF CHILDREN**

USE PRECAUTIONS:

Sulfur and sulfur compounds should not be applied in mid-day when the temperature is above 75-80° F. in arid sections or when temperature is above 85-90° F. in sections where there is normal rainfall and humidity. Do not combine sulfur with foliar spray oil or use sulfur within 50 days before or after foliar oil applications. Rinse equipment and container and dispose of wastes in non-crop land away from water supplies.

Do not permit workers to enter treated area where substantial contact will be made with treated foliage until following days have lapsed after application: Grapes, 1; Peaches and Nectarines, 1; Citrus, 1.

Do not re-use container. Burn bag after using. Keep out of smoke.


NOTICE TO BUYER—Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

— NON-RETURNABLE —

DIRECTIONS FOR MIXING

Mix in water with agitation; agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted: Vegetable and Field Crops—20 to 250 gallons by ground, 5 to 10 gallons by air. Small Fruits—20 to 250 gallons by ground, 7 to 10 gallons by air. Tree Fruits—40 to 800 gallons by ground, 10 to 20 gallons by air. Citrus—100 to 3,000 gallons by ground, 10 to 20 gallons by air. On young fruit trees, use a minimum of 20 gallons by ground per acre.

4 75



EPA REGISTRATION NO. 5967-143 AA

EPA EST. NO. _____

NET WEIGHT 50 LBS.

MANUFACTURED FOR

MOYER CHEMICAL COMPANY

1310 BAYSHORE HIGHWAY P. O. BOX 945 SAN JOSE, CALIFORNIA

