

AC

5967-144

Jun. 26, 1975

UNDER M...

DIRECTIONS FOR USE:

ALFALFA: For control of Aphids, Leafhoppers, Spider Mites, Lygus Bugs, Armyworms, use 1/2 to 1 pint per acre. Do not apply within 7 days of harvest or pasturing.

APPLES: For control of Woolly Apple Aphid, Rosy Aphid, Green Aphid, use 3/4 pint per 100 gals. water per acre. Do not apply within 3 days of harvest. May injure foliage on some varieties.

BEANS: For control of Aphids, Spider Mites, Lygus Bugs, Bean Beetles, use 1/2 to 1 3/4 pints per acre. Do not apply within 1 day of harvest.

BEETS: For control of Aphids, use 3/4 to 2 1/2 pints per acre. Do not apply within 7 days of harvest.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE, COLLARDS: For control of Aphids, Cabbage Looper, use 1/2 to 2 1/2 pints per acre. Do not apply within 3 days of harvest on Broccoli. Do not apply within 7 days of harvest on Brussels Sprouts, Cabbage, Cauliflower and Kale.

CARROTS: For control of Aphids, Leafhoppers, use 3/4 to 2 pints per acre. Do not apply within 7 days of harvest.

CELERY: For control of Aphids, Spider Mites, use 3/4 to 1 1/2 pints. Do not apply within 7 days of harvest.

COTTON: For control of Aphids, Leafhoppers, Whiteflies, Thrips, Lygus Bugs, Leafworms, Leaf-perforators, use 3/4 to 4 pints per acre. No limitations.

CUCUMBERS SQUASH: For control of Aphids, Spider Mites, Cucumber Beetles, use 3/4 to 1 1/2 pints per acre. Do not apply within 1 day of harvest.

GRAPES: For control of Mealy Bugs, Leafhoppers, Spider Mites, use 1 to 2 3/4 pints per acre. Do not apply within 3 days of harvest.

LETTUCE: For control of Aphids, Leafhoppers, Cabbage Loopers, use 1 to 1 1/2 pints per acre. Do not apply within 7 days of harvest on Head Lettuce or 14 days of harvest on Leaf Lettuce.

MELONS: For control of Leafhoppers, Aphids, Cucumber Beetles, Spider Mites, Aphids, use 3/4 to 1 3/4 pints per acre. Do not apply within 1 day of harvest.

ORNAMENTALS: For control of Aphids, Leafhoppers, Mealy Bugs, Thrips, Whiteflies, use 3/4 pint per 100 gals. water per acre. For control of Soft Brown Scale, Monterey Pine Scale, use 1 pint per 100 gals. water per acre. May cause injury to Boston, Maidenhair and Pteris Ferns, and some species of Crassula.

PEARS: For control of Aphids, European Red Mite, Mealy Bugs, use 3/4 pints per 100 gals. water per acre. Do not apply within 1 day of harvest.

POTATOES: For control of Aphids, Leafhoppers, Mealy Bugs, use 1/2 to 3 pints per acre. No limitations.

PRUNES: For control of Leafhoppers, Aphids, Thrips, European Red Mite, use 1/2 pint per 100 gals. water per acre. Do not apply within 3 days of harvest.

STRAWBERRIES: For control of Aphids, Lygus Bugs, Whiteflies, Leafrollers, use 3/4 to 1 1/2 pints per acre. Do not apply within 3 days of harvest.

SUGAR BEETS: For control of Aphids, Spider Mites, Leafhoppers, use 3/4 to 2 pints per acre. Do not apply within 7 days of harvest if tops are to be used for food or feed.

TOMATOES: For control of Aphids, Spider Mites, Vinegar Flies, use 1/2 to 2 pints per acre. For Fruit-worm, use 3 1/2 pints per acre. Do not apply within 5 days of harvest.

WALNUTS: For control of Walnut Aphid, use 3/4 pint per 100 gals. water. No time limitations.

See Right Panel for Other Directions For Use and Mixing Directions.

MOYER

CHEMICAL

MALATHION SPRAY NO. 8-E

ACTIVE INGREDIENTS	By Weight
Malathion*	81.00%
Xylene	7.70%
INERT INGREDIENTS	11.30%
TOTAL	100.00%

*Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate)
This material contains 8 lbs. of Malathion per gallon.

EPA REGISTRATION NO. 5967-144 AA
EPA EST. 5967-CA-1

CAUTION: Keep Out of Reach of Children

1/75
SEE RIGHT PANEL FOR ADDITIONAL CAUTION STATEMENTS

Manufactured by

MOYER CHEMICAL COMPANY

1316 BAYSHORE HIGHWAY P. O. BOX 945, SAN JOSE, CALIFORNIA

CORN of harv
CORN apply
ENDIV
FIGS: 1
100 ga
MUSTA
7 days
ONION
of harv
RADIS
DIREC
amount
Field (C
by gro
Citrus-
of 20 g
Do not
foliage

Do no
stroy e

Harmfu
feed at

This pro
occur.
by clea
This pro
tion me
DO NI
NOTIC
produc
when st

MOYER

CHEMICAL

MALATHION SPRAY NO. 8-E

ACTIVE INGREDIENTS	By Weight
Malathion*	81.00%
Xylene	7.70%
INERT INGREDIENTS	11.20%
TOTAL.....	100.00%

*Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate)
This material contains 8 lbs. of Malathion per gallon.

EPA REGISTRATION NO. 5967-144 AA
EPA EST. 5967-CA-1

CAUTION: Keep Out of Reach of Children

1/75
SEE RIGHT PANEL FOR ADDITIONAL CAUTION STATEMENTS

Manufactured by

MOYER CHEMICAL COMPANY

1310 BAY SHORE HIGHWAY, P. O. BOX 945, SAN JOSE, CALIFORNIA

DIRECTIONS FOR USE:

CORN: For control of Aphids, Corn Earworm, use $\frac{3}{4}$ to 1 pint per acre. Do not apply within 5 days of harvest.

CORN, Field, Pop: For control of Corn Earworm, Thrips, Aphids, use $\frac{1}{2}$ to 1 pint per acre. Do not apply within 5 days of harvest.

ENDIVE: For control of Aphids, use 1 to 2 pints per acre. Do not apply within 7 days of harvest.

FIGS: For control of Dried Fruit Beetle, Drosophila Fly, use $\frac{1}{2}$ to 2 pints to 1 to 2 gals. molasses per 100 gals. of water per acre. Maximum dosage $2\frac{1}{2}$ pints per acre. Do not apply within 3 days of harvest.

MUSTARD GREENS, SPINACH: For control of Aphids, use 1 to 2 pints per acre. Do not apply within 7 days of harvest.

ONIONS, GARLIC: For control of Thrips, Aphids, use 1 to 2 pints per acre. Do not apply within 3 days of harvest.

RADISHES: For control of Aphids, use 1 to 2 pints per acre. Do not apply within 7 days of harvest.

DIRECTIONS FOR MIXING: Mix in water with agitation; agitate while spraying. Use designated amount of product in the following gallons of water per acre, unless otherwise noted: Vegetable and Field Crops—20 to 250 gallons by ground, 5 to 10 gallons by air. Small Fruits—20 to 250 gallons by ground, 7 to 10 gallons by air. Tree Fruits—40 to 800 gallons by ground, 10 to 20 gallons by air. Citrus—100 to 3,000 gallons by ground, 10 to 20 gallons by air. On Young Fruit Trees, use a minimum of 20 gallons by ground per acre.

Do not permit workers to enter treated vineyards where substantial contact will be made with treated foliage until one day after application.

USE PRECAUTIONS:

Do not allow spray to drift onto adjacent crops. Do not apply while bees are working in the field. Destroy empty container by puncturing and burying to prevent re-use.

CAUTION:

Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Avoid contamination of feed and foodstuffs. Wash after handling.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment on residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

NOTICE TO BUYER — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and for handling of this material when such use and/or handling is contrary to label instructions. — **NON-RETURNABLE** —

NET CONTENTS.....GALLONS