

TIME-SAVER

SANNI - RINSE

Odorless Sanitizing Agent

Phenol Coefficient 25 for *S. Typhosa* at 20° C.

Active Ingredient - Alkyl* dimethyl benzyl ammonium chlorides

*(50% C14, 40% C12, 10% C10) 10%

Inert Ingredients 90%

This product fulfills the criteria of Appendix I of the Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 550 ppm of hardness calculated as CaCO_3 when tested by the AOAC Germicidal and Detergent Sanitizers Official Method.

E.P.A. REG. NO. 5991-1

WARNING: Keep out of reach of Children.


Causes eye or skin irritation. Do not get in eyes, on skin or on clothing.

HARMFUL IF SWALLOWED.

AVOID CONTAMINATION OF FOODS.

SEE ANTIDOTE STATEMENT ON SIDE PANEL

Manufactured By
TIME CHEMICAL INC.
CHICAGO • ATLANTA • NEW YORK



DIRECTIONS

As Final Sanitizing Rinse for Eating Utensils, Dishes and Drinking Glasses

- Wash articles in warm water containing a good detergent other than soap. **DO NOT USE SOAP.**
- Rinse with clear water.
- Place in sanitizing solution 200 p.p.m. SANNI-RINSE for 2 minutes or for the contact time specified by the governing Sanitary Code.

Food Processing Plants

Thoroughly clean and rinse all surfaces prior to sanitizing. Sanitize by wiping, spraying or misting with a solution of 200 p.p.m. SANNI-RINSE.

SANNI-RINSE is not to be used as a preservative for foods.

Hospitals and Household Use

For the general disinfection of floors, walls, furniture, bath tubs, stall showers, toilet seats, first clean with a good detergent. Then disinfect with a SANNI-RINSE solution of 400 p.p.m.

For specific disinfection against the organism *Pseudomonas Aeruginosa*, it is necessary for use solution to contain 800 p.p.m. active agent.

Milk Farms and Milk Processing Plants

First clean equipment and utensils thoroughly with a good detergent. Then sanitize with a SANNI-RINSE solution of 200 p.p.m. A further rinse in clean running water is recommended.

As A Deodorant Solution

For waste containers, soiled materials or surfaces, use 1 to 2 ounces of SANNI-RINSE to each gallon of water.

CONCENTRATION CHART

Active Agents	Sanni-Rinse
200 p.p.m.	1 ounce to 4 gallons
400 p.p.m.	1 ounce to 2 gallons
800 p.p.m.	1 ounce to 1 gallon

NO - RINSE REGULATIONS

SANNI-RINSE when used at 200 p.p.m. or less necessitates no water rinse.

SANNI-RINSE has been authorized for use by most leading health authorities for sanitizing of eating and drinking utensils in restaurants, hotels, bars, taverns, soda fountains and camps.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use. If swallowed, promptly drink a large quantity of water. Avoid alcohol. Call a physician immediately.

Rinse empty container thoroughly with water and discard it.

Contents . . . 1 gallon